

Multiscale Analysis in Momentum Space for Quasi-periodic Potential in Dimension Two

Yu. Karpeshina, R. Shterenberg

September 11, 2012

Abstract

We consider a polyharmonic operator $H = (-\Delta)^l + V(\vec{x})$ in dimension two with $l \geq 2$, l being an integer, and a quasi-periodic potential $V(\vec{x})$. We prove that the absolutely continuous spectrum of H contains a semiaxis and there is a family of generalized eigenfunctions at every point of this semiaxis with the following properties. First, the eigenfunctions are close to plane waves $e^{i\langle \vec{\lambda}, \vec{x} \rangle}$ at the high energy region. Second, the isoenergetic curves in the space of momenta $\vec{\lambda}$ corresponding to these eigenfunctions have a form of slightly distorted circles with holes (Cantor type structure). A new method of multiscale analysis in the momentum space is developed to prove these results.

Contents

1	Introduction	4
2	Preliminary Remarks	11
3	Step I	13
3.1	Operator $H^{(1)}$	13
3.2	Perturbation Formulas	13
3.3	Geometric Considerations	17
3.4	Isoenergetic Surface for Operator $H^{(1)}$	18
3.5	Preparation for Step II. Construction of the Second Nonresonant Set . . .	20
3.5.1	Model Operator for Step II	20
3.5.2	Estimates for the Resolvent of the Model Operator	22
3.5.3	Resonant and Nonresonant Sets for Step II	29

4	Step II	31
4.1	Operator $H^{(2)}$. Perturbation Formulas	31
4.2	Isoenergetic Surface for Operator $H^{(2)}$	37
4.3	Preparation for Step III - Geometric Part. Properties of the Quasiperiodic Lattice	38
4.3.1	General Lemmas	38
4.3.2	Lattice Points in the Nonresonant Set	40
4.3.3	Lattice Points in the Resonant Set	41
4.4	Preparation for Step III - Analytic Part	46
4.4.1	Model Operator for Step III	46
4.4.2	Resonant and Nonresonant Sets for Step III	55
5	Step III	56
5.1	Operator $H^{(3)}$. Perturbation Formulas	56
5.2	Isoenergetic Surface for Operator $H^{(3)}$	66
5.3	Preparation for Step IV	66
5.3.1	Properties of the Quasiperiodic Lattice. Continuation	66
5.3.2	Model Operator for Step IV	72
5.3.3	Resonant and Nonresonant Sets for Step IV	78
6	STEP IV	79
6.1	Operator $H^{(4)}$. Perturbation Formulas	79
6.2	Isoenergetic Surface for Operator $H^{(4)}$	81
7	Induction	82
7.1	Inductive formulas for r_n	82
7.2	Preparation for Step $n + 1$, $n \geq 4$	82
7.2.1	Properties of the Quasiperiodic Lattice. Induction	82
7.2.2	Model Operator for Step $n + 1$	86
7.2.3	Resonant and Nonresonant Sets for Step $n + 1$	90
7.3	Operator $H^{(n+1)}$. Perturbation Formulas	91
7.4	Isoenergetic Surface for Operator $H^{(n+1)}$	93
8	Isoenergetic Sets. Generalized Eigenfunctions of H	93
8.1	Construction of Limit-Isoenergetic Set	93
8.2	Generalized Eigenfunctions of H	94
9	Proof of Absolute Continuity of the Spectrum	98
9.1	Operators $E_n(\mathcal{G}'_n)$, $\mathcal{G}'_n \subset \mathcal{G}_n$	98
9.2	Sets $\mathcal{G}_\infty$ and $\mathcal{G}_{\infty,\lambda}$	104
9.3	Projections $E(\mathcal{G}_{\infty,\lambda})$	105
9.4	Proof of Absolute Continuity	107

10 Appendices	107
10.1 Appendix 1. Proof of Lemma 3.12	107
10.2 Appendix 2. Proof of Lemma 3.13	108
10.3 Appendix 3	112
10.4 Appendix 4	116
10.5 Appendix 5	118
10.6 Appendix 6. On Application of Bezout Theorem	119
10.7 Appendix 7. On the Proof of Geometric Lemmas Allowing to Deal with Clusters instead of Boxes	120

1 Introduction

We study an operator

$$H = (-\Delta)^l + V(\vec{x}) \quad (1)$$

in two dimensions, where l is an integer, $l \geq 2$, $V(\vec{x})$ is a quasi-periodic potential being a trigonometric polynomial:

$$V = \sum_{\mathbf{s}_1, \mathbf{s}_2 \in \mathbb{Z}^2, 0 < |\mathbf{s}_1| + |\mathbf{s}_2| \leq Q} V_{\mathbf{s}_1, \mathbf{s}_2} e^{2\pi i \langle \mathbf{s}_1 + \alpha \mathbf{s}_2, \vec{x} \rangle}, \quad 1 \leq Q < \infty. \quad (2)$$

We assume that the irrationality measure μ of α is finite: $\mu < \infty$, or in other words, that α is not a Liouville number¹.

The one-dimensional situation $d = 1$, $l = 1$ is thoroughly investigated in discrete and continuum settings, see e.g. [1]–[13] and references there. It is known that a one-dimensional quasi-periodic Schrödinger operator demonstrates spectral and transport properties which are not close to those of a periodic operator. The spectrum of the quasi-periodic operator is, as a rule, a Cantor set, while in the periodic case, it has a band structure. In the periodic case the spectrum is absolutely continuous, while in the quasi-periodic case, it can have any nature: absolutely continuous, singular continuous and pure point. The transition between different types of spectrum can happen even with a small change of a coefficient in a quasi-periodic operator [8]. The mechanism of the difference in spectral behavior between periodic and quasi-periodic cases can be explained by a phenomenon which is known as resonance tunneling in quantum mechanics. It is associated with small denominators appearing in formal series of perturbation theory. Since the spectrum of the one-dimensional Laplacian is thin (multiplicity 2), resonance tunneling can produce an effect strong enough to destroy the spectrum. If a potential is periodic, then resonance tunneling produces gaps in the spectrum near the points $\lambda_n = (\pi n/a)^2$, $n \in \mathbb{Z}$, a being the period of the potential. If the potential is quasi-periodic, then it can be thought as a sort of combination of infinite number of periodic potentials, each of them producing gaps near its own λ_n -s. Since the set of all λ_n -s can be dense, the number of points surrounded by gaps can be dense too. Thus, the spectrum gets a Cantor like structure. The properties of the operator in the high energy region for the continuum case $d = 1$ are studied in [1]–[4], [7]. The KAM method is used to prove absolute continuity of the spectrum and existence of quasiperiodic solutions at high energies.

There are important results on the density of states, spectrum, localization concerning the quasi-periodic operators in $\mathbb{Z}^d$ and, partially, in $\mathbb{R}^d$, $d > 1$, e.g. [14]–[24]. However, it is still much less known about (1) than about its one-dimensional analog. The properties of the spectrum in the high energy region, existence of extended states and quantum transport are still the wide open problems in the multidimensional case.

Here we study properties of the spectrum and eigenfunctions of (1) in the high energy region. We prove the following results for the case $d = 2$, $l \geq 2$.

¹ Note, that $\mu \geq 2$ for any irrational number α .

1. The spectrum of the operator (1) contains a semiaxis.

This is a generalization of a renown Bethe-Sommerfeld conjecture, which states that in the case of a periodic potential, $l = 1$ and $d \geq 2$, the spectrum of (1) contains a semiaxis. There is a variety of proofs for the periodic case, the earliest one is [25]. For a limit-periodic periodic potential, being periodic in one direction, the conjecture is proved in [26]. For a general case of limit-periodic potential the conjecture is proven in [27]–[29]. Here we present the first proof of (a generalized) Bethe-Sommerfeld conjecture for a quasi-periodic potential.

2. There are generalized eigenfunctions $\Psi_\infty(\vec{x}, \vec{x})$, corresponding to the semi-axis, which are close to plane waves: for every $\vec{x}$ in an extensive subset $\mathcal{G}_\infty$ of $\mathbb{R}^2$, there is a solution $\Psi_\infty(\vec{x}, \vec{x})$ of the equation $H\Psi_\infty = \lambda_\infty\Psi_\infty$ which can be described by the formula:

$$\Psi_\infty(\vec{x}, \vec{x}) = e^{i\langle \vec{x}, \vec{x} \rangle} (1 + u_\infty(\vec{x}, \vec{x})), \quad (3)$$

$$\|u_\infty\|_{L_\infty(\mathbb{R}^2)} = O(|\vec{x}|^{-\gamma_1}), \quad \gamma_1 > 0, \quad (4)$$

where $u_\infty(\vec{x}, \vec{x})$ is a quasi-periodic function, namely a point-wise convergent series of exponentials $e^{2\pi i \langle \mathbf{n} + \alpha \mathbf{m}, \vec{x} \rangle}$, $\mathbf{n}, \mathbf{m} \in \mathbb{Z}^2$. The eigenvalue $\lambda_\infty(\vec{x})$, corresponding to $\Psi_\infty(\vec{x}, \vec{x})$, is close to $|\vec{x}|^{2l}$:

$$\lambda_\infty(\vec{x}) = |\vec{x}|^{2l} + O(|\vec{x}|^{-\gamma_2}), \quad \gamma_2 > 0. \quad (5)$$

The “non-resonant” set $\mathcal{G}_\infty$ of vectors $\vec{x}$, for which (3) – (5) hold, is an extensive Cantor type set: $\mathcal{G}_\infty = \bigcap_{n=1}^\infty \mathcal{G}_n$, where $\{\mathcal{G}_n\}_{n=1}^\infty$ is a decreasing sequence of sets in $\mathbb{R}^2$. Each $\mathcal{G}_n$ has a finite number of holes in each bounded region. More and more holes appear when n increases, however holes added at each step are of smaller and smaller size. The set $\mathcal{G}_\infty$ satisfies the estimate:

$$|\mathcal{G}_\infty \cap \mathbf{B}_R| = |\mathbf{B}_R| (1 + O(R^{-\gamma_3})), \quad \gamma_3 > 0, \quad (6)$$

where $\mathbf{B}_R$ is the disk of radius R centered at the origin, $|\cdot|$ is the Lebesgue measure in $\mathbb{R}^2$.

3. The set $\mathcal{D}_\infty(\lambda)$, defined as a level (isoenergetic) set for $\lambda_\infty(\vec{x})$,

$$\mathcal{D}_\infty(\lambda) = \{\vec{x} \in \mathcal{G}_\infty : \lambda_\infty(\vec{x}) = \lambda\},$$

is proven to be a slightly distorted circle with infinite number of holes. It can be described by the formula:

$$\mathcal{D}_\infty(\lambda) = \{\vec{x} : \vec{x} = \kappa_\infty(\lambda, \vec{v})\vec{v}, \quad \vec{v} \in \mathcal{B}_\infty(\lambda)\}, \quad (7)$$

where $\mathcal{B}_\infty(\lambda)$ is a subset of the unit circle S_1 . The set $\mathcal{B}_\infty(\lambda)$ can be interpreted as the set of possible directions of propagation for almost plane waves (3). The set $\mathcal{B}_\infty(\lambda)$ has a Cantor type structure and an asymptotically full measure on S_1 as $\lambda \rightarrow \infty$:

$$L(\mathcal{B}_\infty(\lambda)) \underset{\lambda \rightarrow \infty}{=} 2\pi + O(\lambda^{-\gamma_4/2l}), \quad \gamma_4 > 0, \quad (8)$$

here and below $L(\cdot)$ is a length of a curve. The value $\varkappa_\infty(\lambda, \vec{\nu})$ in (7) is the “radius” of $\mathcal{D}_\infty(\lambda)$ in a direction $\vec{\nu}$. The function $\varkappa_\infty(\lambda, \vec{\nu}) - \lambda^{1/2l}$ describes the deviation of $\mathcal{D}_\infty(\lambda)$ from the perfect circle of the radius $\lambda^{1/2l}$. It is proven that the deviation is asymptotically small:

$$\varkappa_\infty(\lambda, \vec{\nu}) \underset{\lambda \rightarrow \infty}{=} \lambda^{1/2l} + O(\lambda^{-\gamma_5}), \quad \gamma_5 > 0. \quad (9)$$

4. The branch of the spectrum of the operator (1) corresponding to the generalized eigenfunctions $\Psi_\infty(\vec{\mathcal{Z}}, \vec{x})$ is absolutely continuous.

To prove the results listed above we suggest a method which can be described as *multiscale analysis in the space of momenta*. This is a development of the method, which is used in [27]–[29] for the case of limit-periodic potentials. The essential difference is that in [27]–[29] we constructed a modification of KAM method, where the space variable $\vec{x}$ still plays some role (e.g. in the uniform in $\vec{x}$ approximation of a limit-periodic potential by periodic ones), while in the present situation all considerations are happening in the space of the dual variable $\vec{\mathcal{Z}}$. The KAM method in [27]–[29] was motivated by [30]–[32], where the method is used for periodic problems. Multiscale analysis which we apply here is deeply analogous to the original multiscale method developed in [33] (see also [19], [20]) for the proof of localization. The essential difference is that in [33], [19], [20] the multiscale procedure is constructed with respect to space variable $\vec{x}$ to prove localization, while we construct a multiscale procedure in the space of momenta $\vec{\mathcal{Z}}$ to prove delocalization.

Here is a brief description of the iteration procedure which leads to the results described above. Indeed, let $\vec{\mathcal{Z}} \in \mathbb{R}^2$. We consider a set of finite linear combinations of plane waves $e^{i\langle \vec{\mathcal{Z}} + 2\pi(\mathbf{n} + \alpha\mathbf{m}), \vec{x} \rangle}$, $\mathbf{n}, \mathbf{m} \in \mathbb{Z}^2$. The set is invariant under action of the differential expression (1). Let $H(\vec{\mathcal{Z}})$ be a matrix describing action of (1) in the linear set of the exponents. Obviously,

$$H(\vec{\mathcal{Z}}) = H_0(\vec{\mathcal{Z}}) + V, \quad H_0(\vec{\mathcal{Z}})_{(\mathbf{n}, \mathbf{m}), (\mathbf{n}', \mathbf{m}')} = |\vec{\mathcal{Z}} + 2\pi(\mathbf{n} + \alpha\mathbf{m})|_{\mathbb{R}^2}^2 \delta_{(\mathbf{n}, \mathbf{n}')} \delta_{(\mathbf{m}, \mathbf{m}')},$$

$$V_{(\mathbf{n}, \mathbf{m}), (\mathbf{n}', \mathbf{m}')} = V_{\mathbf{n} - \mathbf{n}', \mathbf{m} - \mathbf{m}'}.$$

Next, we consider an expanding sequence of finite sets Ω_n in the space $\mathbb{Z}^2 \times \mathbb{Z}^2$ of indices $(\mathbf{n}, \mathbf{m})$: $\Omega_n \subset \Omega_{n+1}$, $\lim_{n \rightarrow \infty} \Omega_n = \mathbb{Z}^2 \times \mathbb{Z}^2$. Let P_n be the characteristic projection of set Ω_n in the space $\ell^2(\mathbb{Z}^2 \times \mathbb{Z}^2)$. We consider a sequence of finite matrices $H^{(n)}(\vec{\mathcal{Z}}) = P_n H(\vec{\mathcal{Z}}) P_n$. Each matrix corresponds to a finite dimensional operator in $\ell^2(\mathbb{Z}^2 \times \mathbb{Z}^2)$, given that the operator acts as zero on $(I - P_n)\ell^2$. For each n we construct a “non-resonant” set $\mathcal{G}_n$ in

the space $\mathbb{R}^2$ of momenta $\vec{\mathcal{Z}}$, such that: if $\vec{\mathcal{Z}} \in \mathcal{G}_n$, then $H^{(n)}(\vec{\mathcal{Z}}) = P_n H(\vec{\mathcal{Z}}) P_n$ has an eigenvalue $\lambda_n(\vec{\mathcal{Z}})$ and its spectral projector $\mathcal{E}_n(\vec{\mathcal{Z}})$ which can be described by perturbation formulas with respect to the previous operator $H^{(n-1)}(\vec{\mathcal{Z}})$. If $\vec{\mathcal{Z}} \in \cap_{n=1}^\infty \mathcal{G}_n$ then $\lambda_n(\vec{\mathcal{Z}})$ and $\mathcal{E}_n(\vec{\mathcal{Z}})$ have limits. The linear combinations of the exponentials, corresponding to the projectors $\mathcal{E}_n(\vec{\mathcal{Z}})$, have a point-wise limit in $\vec{x}$, the limit being a generalized eigenfunction of (1). The generalized eigenfunction is close to the plane wave $e^{i\langle \vec{\mathcal{Z}}, \vec{x} \rangle}$ in the high energy region.

Each matrix $H^{(n)}$ is considered as a perturbation of a matrix $\hat{H}^{(n)}$, the latter has a block structure, i.e., consists of a variety of blocks $H^{(s)}(\vec{\mathcal{Z}} + 2\pi(\mathbf{n} + \alpha\mathbf{m}))$, $s = 1, \dots, n-1$, and, naturally, some diagonal terms. Blocks with different indices (s) have sizes of different orders of magnitude (the size increasing with s). Thus we have a multiscale structure in the definition of $\hat{H}^{(n)}$. We use $\hat{H}^{(n)}(\vec{\mathcal{Z}})$ as a starting operator to construct perturbation series for $H^{(n)}(\vec{\mathcal{Z}})$. At a step n we apply our knowledge of spectral properties of $H^{(s)}(\vec{\mathcal{Z}} + 2\pi(\mathbf{n}' + \alpha\mathbf{m}'))$, $s = 1, \dots, n-1$, $\mathbf{n}', \mathbf{m}' \in \mathbb{Z}^2$, obtained in the previous steps, to describe spectral properties of $H^{(n)}(\vec{\mathcal{Z}} + 2\pi(\mathbf{n} + \alpha\mathbf{m}))$, $\mathbf{n}, \mathbf{m} \in \mathbb{Z}^2$ and to construct $\mathcal{G}_n$.

At step one we use a regular perturbation theory and elementary geometric considerations to prove the following results. There is a set $\mathcal{G}_1 \subset \mathbb{R}^2$ such that: if $\vec{\mathcal{Z}} \in \mathcal{G}_1$, then the operator $H^{(1)}(\vec{\mathcal{Z}})$ has a single eigenvalue close to the unperturbed one:

$$\lambda^{(1)}(\vec{\mathcal{Z}}) \underset{|\vec{\mathcal{Z}}| \rightarrow \infty}{=} |\vec{\mathcal{Z}}|^{2l} + O(|\vec{\mathcal{Z}}|^{-\gamma_2}), \quad \gamma_2 > 0. \quad (10)$$

A normalized eigenvector $\mathbf{u}^{(1)}$ is also close to the unperturbed one: $\mathbf{u}^{(1)} = \mathbf{u}^{(0)} + \tilde{\mathbf{u}}^{(1)}$, where $(\mathbf{u}^{(0)})_{(\mathbf{n}, \mathbf{m})} = \delta_{\mathbf{n}, \mathbf{0}} \delta_{\mathbf{m}, \mathbf{0}}$ and the l^1 -norm of $\tilde{\mathbf{u}}^{(1)}$ is small: $\|\tilde{\mathbf{u}}^{(1)}\|_{l^1} < |\vec{\mathcal{Z}}|^{-\gamma_1}$, $\gamma_1 > 0$. It follows that:

$$\Psi_1(\vec{\mathcal{Z}}, \vec{x}) = e^{i\langle \vec{\mathcal{Z}}, \vec{x} \rangle} + \tilde{u}_1(\vec{\mathcal{Z}}, \vec{x}), \quad \|\tilde{u}_1\|_{L^\infty(\mathbb{R}^2)} \underset{|\vec{\mathcal{Z}}| \rightarrow \infty}{=} O(|\vec{\mathcal{Z}}|^{-\gamma_1}), \quad \gamma_1 > 0, \quad (11)$$

where $\Psi_1(\vec{\mathcal{Z}}, \vec{x})$, $\tilde{u}_1(\vec{\mathcal{Z}}, \vec{x})$ are the linear combinations of the exponentials corresponding to vectors $\mathbf{u}^{(1)}$ and $\tilde{\mathbf{u}}^{(1)}$, respectively. It is shown that function $\Psi_1(\vec{\mathcal{Z}}, \vec{x})$ satisfies the equation for eigenfunctions with a good accuracy:

$$-\Delta \Psi_1 + V \Psi_1 = |\vec{\mathcal{Z}}|^{2l} \Psi_1 + f_1, \quad \|f_1\|_{L^\infty(\mathbb{R}^2)} \underset{|\vec{\mathcal{Z}}| \rightarrow \infty}{=} O(|\vec{\mathcal{Z}}|^{-\gamma_6}), \quad \gamma_6 > 0. \quad (12)$$

Relation (10) is differentiable:

$$\nabla \lambda^{(1)}(\vec{\mathcal{Z}}) \underset{|\vec{\mathcal{Z}}| \rightarrow \infty}{=} 2l |\vec{\mathcal{Z}}|^{2l-2} \vec{\mathcal{Z}} + O(|\vec{\mathcal{Z}}|^{-\gamma_7}), \quad \gamma_7 > 0. \quad (13)$$

Next, we construct a sequence $\mathcal{G}_n$, $n \geq 2$, such for any $\vec{\mathcal{Z}} \in \mathcal{G}_n$ the operator $H^{(n)}(\vec{\mathcal{Z}})$ has a single eigenvalue $\lambda^{(n)}(\vec{\mathcal{Z}})$ in a super exponentially small neighborhood of $\lambda^{(n-1)}(\vec{\mathcal{Z}})$:

$$\lambda^{(n)}(\vec{\mathcal{Z}}) \underset{|\vec{\mathcal{Z}}| \rightarrow \infty}{=} \lambda^{(n-1)}(\vec{\mathcal{Z}}) + O(|\vec{\mathcal{Z}}|^{-|\vec{\mathcal{Z}}|^{\gamma_8 n}}), \quad \gamma_8 > 0. \quad (14)$$




Figure 1: Isoenergetic curve $\mathcal{D}_1(\lambda)$




Figure 2: Isoenergetic curve $\mathcal{D}_2(\lambda)$

Similar estimates hold for the eigenvectors and the corresponding functions $\Psi_n(\vec{\mathcal{Z}}, \vec{x})$:

$$\Psi_n(\vec{\mathcal{Z}}, \vec{x}) = \Psi_{n-1}(\vec{\mathcal{Z}}, \vec{x}) + \tilde{u}_n(\vec{\mathcal{Z}}, \vec{x}), \quad \|\tilde{u}_n\|_{L_\infty(\mathbb{R}^2)} \underset{k \rightarrow \infty}{=} O(|\vec{\mathcal{Z}}|^{-|\vec{\mathcal{Z}}|^{\gamma_9 n}}), \quad \gamma_9 > 0. \quad (15)$$

$$-\Delta \Psi_n + V \Psi_n = \lambda^{(n)}(\vec{\mathcal{Z}}) \Psi_n + f_n, \quad \|f_n\|_{L_\infty(\mathbb{R}^2)} \underset{|\vec{\mathcal{Z}}| \rightarrow \infty}{=} O(|\vec{\mathcal{Z}}|^{-|\vec{\mathcal{Z}}|^{\gamma_{10} n}}), \quad \gamma_{10} > 0. \quad (16)$$

Formula (14) is differentiable with respect to $\vec{\mathcal{Z}}$:

$$\nabla \lambda^{(n)}(\vec{\mathcal{Z}}) \underset{|\vec{\mathcal{Z}}| \rightarrow \infty}{=} \nabla \lambda^{(n-1)}(\vec{\mathcal{Z}}) + O(|\vec{\mathcal{Z}}|^{-|\vec{\mathcal{Z}}|^{\gamma_8 n}}), \quad \gamma_8 > 0. \quad (17)$$

In fact, for large n estimates (14) – (17) are even stronger. The non-resonant set $\mathcal{G}_n$ is proven to be extensive in $\mathbb{R}^2$:

$$|\mathcal{G}_n \cap \mathbf{B}_{\mathbf{R}}| \underset{R \rightarrow \infty}{=} |\mathbf{B}_{\mathbf{R}}| (1 + O(R^{-\gamma_3})). \quad (18)$$

Estimates (14) – (18) are uniform in n .

The set $\mathcal{D}_n(\lambda)$ is defined as the level (isoenergetic) set for the non-resonant eigenvalue $\lambda^{(n)}(\vec{\mathcal{Z}})$:

$$\mathcal{D}_n(\lambda) = \{ \vec{\mathcal{Z}} \in \mathcal{G}_n : \lambda^{(n)}(\vec{\mathcal{Z}}) = \lambda \}.$$

This set is proven to be a slightly distorted circle with a finite number of holes (see Fig. 1, 2). The set $\mathcal{D}_n(\lambda)$ can be described by the formula:

$$\mathcal{D}_n(\lambda) = \{ \vec{\mathcal{Z}} : \vec{\mathcal{Z}} = \mathcal{Z}^{(n)}(\lambda, \vec{\nu}) \vec{\nu}, \quad \vec{\nu} \in \mathcal{B}_n(\lambda) \}, \quad (19)$$

where $\mathcal{B}_n(\lambda)$ is a subset of the unit circle S_1 . The set $\mathcal{B}_n(\lambda)$ can be interpreted as the set of possible directions of propagation for almost plane waves $\Psi_n(\vec{\mathcal{Z}}, \vec{x})$, see (11), (15). It has an asymptotically full measure on S_1 as $\lambda \rightarrow \infty$:

$$L(\mathcal{B}_n(\lambda)) \underset{\lambda \rightarrow \infty}{=} 2\pi + O(\lambda^{-\gamma_4/2l}). \quad (20)$$

Each set $\mathcal{B}_n(\lambda)$ has only a finite number of holes, however their number is growing with n . More and more holes of a smaller and smaller size are added at each step. The value $\varkappa^{(n)}(\lambda, \vec{\nu}) - \lambda^{1/2l}$ gives the deviation of $\mathcal{D}_n(\lambda)$ from the perfect circle of the radius $\lambda^{1/2l}$ in the direction $\vec{\nu}$. It is proven that the deviation is asymptotically small:

$$\varkappa^{(n)}(\lambda, \vec{\nu}) = \lambda^{1/2l} + O(\lambda^{-\gamma_5}), \quad \frac{\partial \varkappa^{(n)}(\lambda, \vec{\nu})}{\partial \varphi} = O(\lambda^{-\gamma_{11}}), \quad \gamma_5, \gamma_{11} > 0, \quad (21)$$

φ being an angle variable, $\vec{\nu} = (\cos \varphi, \sin \varphi)$. Estimates (20), (21) are uniform in n .

On each step more and more points are excluded from the non-resonant sets $\mathcal{G}_n$, thus $\{\mathcal{G}_n\}_{n=1}^\infty$ is a decreasing sequence of sets. The set $\mathcal{G}_\infty$ is defined as the limit set: $\mathcal{G}_\infty = \bigcap_{n=1}^\infty \mathcal{G}_n$. It has an infinite number of holes, but nevertheless satisfies the relation (6). For every $\vec{\varkappa} \in \mathcal{G}_\infty$ and every n , there is a generalized eigenfunction of $H^{(n)}$ of the type (11), (15). It is proven that the sequence of $\Psi_n(\vec{\varkappa}, \vec{x})$ has a limit in $L_\infty(\mathbb{R}^2)$ when $\vec{\varkappa} \in \mathcal{G}_\infty$. The function $\Psi_\infty(\vec{\varkappa}, \vec{x}) = \lim_{n \rightarrow \infty} \Psi_n(\vec{\varkappa}, \vec{x})$ is a generalized eigenfunction of H . It can be written in the form (3) – (4). Naturally, the corresponding eigenvalue $\lambda_\infty(\vec{\varkappa})$ is the limit of $\lambda^{(n)}(\vec{\varkappa})$ as $n \rightarrow \infty$.

It is shown that $\{\mathcal{B}_n(\lambda)\}_{n=1}^\infty$ is a decreasing sequence of sets, on each step more and more directions being excluded. We consider the limit $\mathcal{B}_\infty(\lambda)$ of $\mathcal{B}_n(\lambda)$:

$$\mathcal{B}_\infty(\lambda) = \bigcap_{n=1}^\infty \mathcal{B}_n(\lambda). \quad (22)$$

This set has a Cantor type structure on the unit circle. It is proven that $\mathcal{B}_\infty(\lambda)$ has an asymptotically full measure on the unit circle (see (8)). We prove that the sequence $\varkappa^{(n)}(\lambda, \vec{\nu})$, $n = 1, 2, \dots$, describing the isoenergetic curves $\mathcal{D}_n(\lambda)$, quickly converges as $n \rightarrow \infty$. We show that $\mathcal{D}_\infty(\lambda)$ can be described as the limit of $\mathcal{D}_n(\lambda)$ in the sense (7), where $\varkappa_\infty(\lambda, \vec{\nu}) = \lim_{n \rightarrow \infty} \varkappa^{(n)}(\lambda, \vec{\nu})$ for every $\vec{\nu} \in \mathcal{B}_\infty(\lambda)$. It is shown that the derivatives of the functions $\varkappa^{(n)}(\lambda, \vec{\nu})$ (with respect to the angle variable on the unit circle) have a limit as $n \rightarrow \infty$ for every $\vec{\nu} \in \mathcal{B}_\infty(\lambda)$. We denote this limit by $\frac{\partial \varkappa_\infty(\lambda, \vec{\nu})}{\partial \varphi}$. Using (21), we prove that

$$\frac{\partial \varkappa_\infty(\lambda, \vec{\nu})}{\partial \varphi} = O(\lambda^{-\gamma_{11}}). \quad (23)$$

Thus, the limit curve $\mathcal{D}_\infty(\lambda)$ has a tangent vector in spite of its Cantor type structure, the tangent vector being the limit of corresponding tangent vectors for $\mathcal{D}_n(\lambda)$ as $n \rightarrow \infty$. The curve $\mathcal{D}_\infty(\lambda)$ looks as a slightly distorted circle with infinite number of holes for every sufficiently large λ , $\lambda > \lambda_*(V)$. It immediately follows that $[\lambda_*, \infty)$ is in the spectrum of H (Bethe-Sommerfeld conjecture).

The main technical difficulty to overcome is the construction of non-resonant sets $\mathcal{B}_n(\lambda)$ for every fixed sufficiently large λ , $\lambda > \lambda_0(V)$, where $\lambda_0(V)$ is the same for all n . The set $\mathcal{B}_n(\lambda)$ is obtained by deleting a “resonant” part from $\mathcal{B}_{n-1}(\lambda)$. Definition of $\mathcal{B}_{n-1}(\lambda) \setminus \mathcal{B}_n(\lambda)$ includes eigenvalues of $H^{(n-1)}(\vec{\varkappa})$. To describe $\mathcal{B}_{n-1}(\lambda) \setminus \mathcal{B}_n(\lambda)$ one has to consider




Figure 3: Set Φ_2

not only non-resonant eigenvalues of the type (10), (14), but also resonant eigenvalues, for which no suitable formulas are known. Absence of formulas causes difficulties in estimating the size of $\mathcal{B}_{n-1}(\lambda) \setminus \mathcal{B}_n(\lambda)$. To treat this problem we start with introducing an angle variable $\varphi \in [0, 2\pi)$, $\vec{\nu} = (\cos \varphi, \sin \varphi) \in S_1$ and consider sets $\mathcal{B}_n(\lambda)$ in terms of this variable. Next, we show that the resonant set $\mathcal{B}_{n-1}(\lambda) \setminus \mathcal{B}_n(\lambda)$ can be described as the set of zeros of functions of the type

$$\det \left(H^{(s)}(\vec{\mathcal{Z}}_{n-1}(\varphi) + 2\pi(\mathbf{n} + \alpha \mathbf{m})) - \lambda - \varepsilon \right), \quad s = 1, \dots, n-1, \quad (\mathbf{n}, \mathbf{m}) \in \Omega_n \setminus (\mathbf{0}, \mathbf{0}),$$

where $\vec{\mathcal{Z}}_{n-1}(\varphi)$ is a vector-function describing $\mathcal{D}_{n-1}(\lambda)$: $\vec{\mathcal{Z}}_{n-1}(\varphi) = \mathcal{Z}_{n-1}(\lambda, \vec{\nu})\vec{\nu}$. To obtain $\mathcal{B}_{n-1}(\lambda) \setminus \mathcal{B}_n(\lambda)$ we take all values of ε in a small interval and $(\mathbf{n}, \mathbf{m})$ in some subset of Ω_n . Further, we extend our considerations to a complex neighborhood Φ_0 of $[0, 2\pi)$. We show that the determinants are analytic functions of φ and, by this, reduce the problem of estimating the size of the resonant set to a problem in complex analysis. We use theorems for analytic functions to count zeros of the determinants and to investigate how far the zeros move when ε changes. It enables us to estimate the size of the zero set of the determinants, and, hence, the size of the non-resonant set $\Phi_n \subset \Phi_0$, which is defined as a non-zero set for the determinants. Proving that the non-resonant set Φ_n is sufficiently large, we obtain estimates (18) for $\mathcal{G}_n$ and (20) for $\mathcal{B}_n$, the set $\mathcal{B}_n$ corresponding to the real part of Φ_n .

To obtain Φ_n we delete from Φ_0 more and more discs (holes) of smaller and smaller radii at each step. Thus, the non-resonant set $\Phi_n \subset \Phi_0$ has a structure of Swiss Cheese (Fig. 3, 4). Deleting a resonance set from Φ_0 at each step of the recurrent procedure we call a “Swiss Cheese Method”. The essential difference of our method from constructions of non-resonant sets in similar situations before (see e.g. [30]–[32], [34]) is that we construct a non-resonant set not only in the whole space of a parameter ($\vec{\mathcal{Z}} \in \mathbb{R}^2$ here), but also on isoenergetic curves $\mathcal{D}_n(\lambda)$ in the space of the parameter, when λ is sufficiently large. Estimates for the size of non-resonant sets on a curve require more subtle techni-




Figure 4: Set Φ_3

cal considerations than those sufficient for description of a non-resonant set in the whole space of the parameter. But as a reward, such estimates enable us to show that every isoenergetic set for $\lambda > \lambda_0$ is not empty and thus, to prove Bethe-Sommerfeld conjecture.

Note that generalization of the results from the case $l > 1$, l being an integer, to the case of rational l satisfying the same inequality is relatively simple; it requires just slightly more careful technical considerations. The restriction $l > 1$ is also technical, though it is more difficult to lift. The condition $l > 1$ is needed only for the second step of the recurrent procedure. The authors plan to consider the case $l = 1$ in a forthcoming paper. The requirement $\mu < \infty$ is essential, since we use it to estimate the minimal values of $|\mathbf{n} + \alpha \mathbf{m}|$ when $(\mathbf{n}, \mathbf{m}) \in \Omega_n \setminus (\mathbf{0}, \mathbf{0})$. Such estimates are necessary for controlling small denominators in the perturbation series at each step.

The plan of the paper is the following. Preliminary considerations are in Section 2. Sections 3 – 7 describe steps of the recurrent procedure. Note, that Steps I,II are designed to start the procedure. Step III is already typical, however uses some “non-typical” estimates from Steps I,II. Step IV is completely typical: all other steps of the recurrent procedure differ from Step IV only by the change of indices. The proofs of convergence of the iteration procedure and of the results 1 – 3, listed at the beginning of the introduction, are in Section 8. The result 4 about absolutely continuous spectrum is proven in Section 9. Section 10 (Appendices) contains technical lemmas.

Acknowledgement The authors are very grateful to Prof. Parnowski for useful discussions and to Prof. Young-Ran Lee for allowing us to use figures 1-4 from [KL].

2 Preliminary Remarks

We consider two-dimensional quasi-periodic polyharmonic operator

$$(-\Delta)^l + V(\vec{x}), \quad l > 1 \tag{24}$$

which is perturbation of the free operator $H_0 := (-\Delta)^l$. Here V is the potential of the form (2). Without loss of generality we assume $0 < \alpha < 1$. We assume also that irrationality measure μ of α is finite: $\mu < \infty$, or in other words, that α is not a Liouville number. Note, that for irrational number α we always have $\mu \geq 2$. It follows from the definition of the irrationality measure that 1) *For any $\epsilon > 0$ there exists a constant C_ϵ such that for any irreducible rational number $\frac{\tilde{M}}{\tilde{N}}$ we have*

$$\left| \alpha - \frac{\tilde{M}}{\tilde{N}} \right| \geq \frac{C_\epsilon}{\tilde{N}^{\mu+\epsilon}}. \quad (25)$$

2) *For any $\epsilon > 0$ there exists a sequence $\frac{M}{N}$ of irreducible rational numbers such that*

$$\left| \alpha - \frac{M}{N} \right| \leq \frac{1}{N^{\mu-\epsilon}}. \quad (26)$$

For every pair of integer vectors $\mathbf{s}_1, \mathbf{s}_2 \in \mathbb{Z}^2$ we consider $\vec{p}_s := 2\pi(\mathbf{s}_1 + \alpha\mathbf{s}_2)$. We introduce the norm

$$|||\vec{p}_s||| := |\mathbf{s}_1| + |\mathbf{s}_2|.$$

We will also use the notation $p_s := |\vec{p}_s|$ and $\vec{p}_s = p_s(\cos \varphi_s, \sin \varphi_s)$.

Lemma 2.1. *For every $\vec{p}_s \neq 0$ we have*

$$p_s \leq 2\pi |||\vec{p}_s|||, \quad (27)$$

$$p_s \geq 2\pi C_\epsilon |||\vec{p}_s|||^{-(\mu-1+\epsilon)}. \quad (28)$$

Proof. The estimate (27) is obvious. To prove (28) we notice that if $\mathbf{s}_2 = 0$ then $p_s = 2\pi|\mathbf{s}_1| \geq 2\pi$. Let $\mathbf{s}_1 = (s_{11}, s_{12})$, $\mathbf{s}_2 = (s_{21}, s_{22})$. If, for example, $s_{21} \neq 0$ then from (25) and definition of $|||\vec{p}_s|||$ we obtain

$$\begin{aligned} p_s &\geq 2\pi |s_{11} + \alpha s_{21}| = 2\pi |s_{21}| \left| \alpha + \frac{s_{11}}{s_{21}} \right| \geq \\ &2\pi C_\epsilon |s_{21}|^{-\mu-\epsilon+1} \geq 2\pi C_\epsilon |||\vec{p}_s|||^{-(\mu-1+\epsilon)}. \end{aligned} \quad (29)$$

□

We introduce vector $\vec{\kappa}(\varphi) := (\kappa_1, \kappa_2) = \kappa \vec{v} := \kappa(\cos \varphi, \sin \varphi)$. Similar agreement will be used for other vectors. Let $H(\vec{\kappa}) = H(\kappa, \varphi)$ be the "fiber" operator acting in $L_2(\mathbb{Z}^4)$ with its matrix elements given by

$$(H(\vec{\kappa}))_{\mathbf{s}, \mathbf{s}+\mathbf{q}} = |\vec{\kappa} + \vec{p}_s|^{2l} \delta_{\mathbf{s}, \mathbf{s}+\mathbf{q}} + V_{\vec{p}_q}.$$

Here $V_{\vec{p}_q} := V_{\mathbf{q}_1, \mathbf{q}_2}$. (see (2))

$$V_{\vec{p}_q} = 0, \quad \text{when } |||\vec{p}_q||| > Q, \quad (Q < \infty). \quad (30)$$

To simplify the notation in what follows we will write V_q instead of $V_{\vec{p}_q}$ when it does not lead to confusion.

3 Step I

3.1 Operator $H^{(1)}$

Let δ be some small parameter, $0 < \delta < (100\mu)^{-1}$. We put

$$\Omega(\delta) := \{\mathbf{m} \in \mathbb{Z}^2 : \|\vec{p}_{\mathbf{m}}\| \leq k^\delta\}, \quad \tilde{\Omega}(\delta) := \{\mathbf{m} \in \mathbb{Z}^2 : \|\vec{p}_{\mathbf{m}}\| \leq 4k^\delta\}.$$

By $P(\delta)$ we denote orthogonal (diagonal) projection in $l^2(\mathbb{Z}^2)$ on the set of elements supported in $\Omega(\delta)$. We call it the characteristic projector of $\Omega(\delta)$. The dimension of the projector is equal to the number of elements in $\Omega(\delta)$ and, obviously, does not exceed $(8k^\delta)^4$. We have

$$(P(\delta)H_0(\vec{\mathcal{Z}})P(\delta))_{\mathbf{m},\mathbf{n}} = |\vec{\mathcal{Z}} + \vec{p}_{\mathbf{m}}|^{2l} \delta_{\mathbf{m},\mathbf{n}} \chi_{\Omega(\delta)}(\mathbf{m}),$$

where as usual $\chi_{\Omega(\delta)}(\mathbf{m})$ is the characteristic function of the set $\Omega(\delta)$. We are going to consider $H^{(1)}(\vec{\mathcal{Z}}) = P(\delta)H(\vec{\mathcal{Z}})P(\delta)$ as a perturbation of the operator $P(\delta)H_0(\vec{\mathcal{Z}})P(\delta)$.

3.2 Perturbation Formulas

Now we construct a “non-resonant” set of φ , for which the operator $H^{(1)}(\vec{\mathcal{Z}}(\varphi))$ can be constructively considered as a perturbation of $H_0^{(1)}(\vec{\mathcal{Z}}(\varphi))$ corresponding to $V = 0$. In what follows τ is an auxiliary parameter $\frac{1}{32} \leq \tau \leq 32$.

Lemma 3.1 (Geometric). *For every $k > 800$ there is a subset $\omega^{(1)}(k, \delta, \tau)$ of the interval $[0, 2\pi)$ such that:*

1. *For every $\varphi \in \omega^{(1)}(k, \delta, \tau)$ and $\mathbf{m} \in \tilde{\Omega}(\delta) \setminus \{0\}$, the following inequality holds:*

$$\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|^{2l} - k^{2l} \right| > \tau k^{2l-1-40\mu\delta}, \quad \vec{k} := k(\cos \varphi, \sin \varphi). \quad (31)$$

2. *For every φ in the real $\frac{\tau}{16}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$ and $\varkappa \in \mathbb{R} : |\varkappa - k| < \frac{\tau}{16}k^{-40\mu\delta}$, a slightly weaker inequality holds for $\vec{\mathcal{Z}}(\varphi) = \varkappa(\cos \varphi, \sin \varphi)$ and $\mathbf{m} \in \tilde{\Omega}(\delta) \setminus \{0\}$:*

$$\left| |\vec{\mathcal{Z}}(\varphi) + \vec{p}_{\mathbf{m}}|^{2l} - k^{2l} \right| > \frac{\tau^l}{2} k^{2l-1-40\mu\delta}. \quad (32)$$

3. *The set $\omega^{(1)}(k, \delta, \tau)$ has an asymptotically full measure in $[0, 2\pi)$ as $k \rightarrow \infty$. Namely,*

$$|\omega^{(1)}(k, \delta, \tau)| = 2\pi + O(k^{-37\mu\delta}), \quad k \rightarrow \infty. \quad (33)$$

Corollary 3.2. *If φ is in the real $\frac{\tau}{16}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$ and z is on the circle*

$$C_1 = \{z : |z - k^{2l}| = \frac{\tau^l}{4}k^{2l-1-40\mu\delta}\}, \quad (34)$$

then the following inequality holds for all $\mathbf{m} \in \tilde{\Omega}(\delta)$:

$$\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|^{2l} - z \right| \geq \frac{\tau^l}{4}k^{2l-1-40\mu\delta}, \quad z \in C_1. \quad (35)$$

The lemma is proved in Section 3.3 (Corollaries 3.7 and 3.10.) The corollary from the lemma is proven at the end of Section 3.3. Note that in Section 3.3 we construct non-resonance set of φ in the set of complex numbers. Such complex non-resonance set we need for construction of further steps of approximation.

Let $r = 1, 2, \dots$ and

$$g_r^{(1)}(\vec{\varkappa}) := \frac{(-1)^r}{2\pi i r} \text{Tr} \oint_{C_1} ((P(\delta)(H_0(\vec{\varkappa}) - zI)P(\delta))^{-1}VP(\delta))^r dz, \quad (36)$$

$$G_r^{(1)}(\vec{\varkappa}) := \frac{(-1)^{r+1}}{2\pi i} \oint_{C_1} ((P(\delta)(H_0(\vec{\varkappa}) - zI)P(\delta))^{-1}VP(\delta))^r (P(\delta)(H_0(\vec{\varkappa}) - zI)P(\delta))^{-1} dz. \quad (37)$$

Note that $g_1^{(1)}(\vec{\varkappa}) = 0$ since $V = 0$. Coefficient $g_2^{(1)}(\vec{\varkappa})$ admits representation:

$$\begin{aligned} g_2^{(1)}(\vec{\varkappa}) &= \sum_{\mathbf{q} \in \Omega(\delta) \setminus \{0\}} |V_{\mathbf{q}}|^2 (|\vec{\varkappa}|^{2l} - |\vec{\varkappa} + \vec{p}_{\mathbf{q}}|^{2l})^{-1} \\ &= -\frac{1}{2} \sum_{\mathbf{q} \in \Omega(\delta) \setminus \{0\}} \frac{|V_{\mathbf{q}}|^2 (|\vec{\varkappa} + \vec{p}_{\mathbf{q}}|^{2l} + |\vec{\varkappa} - \vec{p}_{\mathbf{q}}|^{2l} - 2|\vec{\varkappa}|^{2l})}{(|\vec{\varkappa}|^{2l} - |\vec{\varkappa} + \vec{p}_{\mathbf{q}}|^{2l})(|\vec{\varkappa}|^{2l} - |\vec{\varkappa} - \vec{p}_{\mathbf{q}}|^{2l})}, \end{aligned} \quad (38)$$

From now on $\|A\|_1$ means the norm of an operator A in the trace class.

Theorem 3.3. *Suppose φ is in the real $\frac{\tau}{16}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$ and $\varkappa \in \mathbb{R}$, $|\varkappa - k| \leq \frac{\tau}{16}k^{-40\mu\delta}$, $\vec{\varkappa} = \varkappa(\cos \varphi, \sin \varphi)$. Then, for sufficiently large $k > k_0(V, \delta, \tau)$ there exists a single eigenvalue of $H^{(1)}(\vec{\varkappa})$ in the interval $\varepsilon_1(k, \delta, \tau) = (k^{2l} - \frac{\tau^l}{2}k^{2l-1-40\mu\delta}, k^{2l} + \frac{\tau^l}{2}k^{2l-1-40\mu\delta})$. It is given by the absolutely converging series:*

$$\lambda^{(1)}(\vec{\varkappa}) = \varkappa^{2l} + \sum_{r=2}^{\infty} g_r^{(1)}(\vec{\varkappa}). \quad (39)$$

For coefficients $g_r^{(1)}(\vec{\varkappa})$ the following estimates hold:

$$|g_r^{(1)}(\vec{\varkappa})| \leq (Ck)^{-(r-1)(2l-1-40\mu\delta)+4\delta}. \quad (40)$$

Moreover,

$$|g_2^{(1)}(\vec{\mathcal{Z}})| \leq Ck^{-2l+(80\mu+6)\delta}. \quad (41)$$

The corresponding spectral projection is given by the series:

$$\mathcal{E}^{(1)}(\vec{\mathcal{Z}}) = \mathcal{E}_0(\vec{\mathcal{Z}}) + \sum_{r=1}^{\infty} G_r^{(1)}(\vec{\mathcal{Z}}), \quad (42)$$

$\mathcal{E}_0(\vec{\mathcal{Z}})$ being the unperturbed spectral projection. The operators $G_r^{(1)}(\vec{\mathcal{Z}})$ satisfy the estimates:

$$\|G_r^{(1)}(\vec{\mathcal{Z}})\|_1 < (Ck)^{-r(2l-1-44\mu\delta)}. \quad (43)$$

Matrix elements of $G_r^{(1)}(\vec{\mathcal{Z}})$ satisfy the following relations:

$$G_r^{(1)}(\vec{\mathcal{Z}})_{\mathbf{s}\mathbf{s}'} = 0, \quad \text{if } rQ < \|\vec{p}_{\mathbf{s}}\| + \|\vec{p}_{\mathbf{s}'}\|. \quad (44)$$

Corollary 3.4. For the perturbed eigenvalue and its spectral projection the following estimates hold:

$$\lambda^{(1)}(\vec{\mathcal{Z}}) = \mathcal{Z}^{2l} + O(k^{-2l+(80\mu+6)\delta}), \quad (45)$$

$$\|\mathcal{E}^{(1)}(\vec{\mathcal{Z}}) - \mathcal{E}_0(\vec{\mathcal{Z}})\|_1 < ck^{-2l+1+44\mu\delta}. \quad (46)$$

Matrix elements of spectral projection $\mathcal{E}^{(1)}(\vec{\mathcal{Z}})$ also satisfy the estimate:

$$|\mathcal{E}^{(1)}(\vec{\mathcal{Z}})_{\mathbf{s}\mathbf{s}'}| < (Ck)^{-d^{(1)}(\mathbf{s},\mathbf{s}')}, \quad d^{(1)}(\mathbf{s},\mathbf{s}') = Q^{-1}(\|\vec{p}_{\mathbf{s}}\| + \|\vec{p}_{\mathbf{s}'}\|)(2l-1-44\mu\delta). \quad (47)$$

The last estimate easily follows from the formula (44) and estimate (43).

Proof. The proof is based on expansion of the resolvent in perturbation series on the circle C_1 . Indeed, let us consider the series

$$(H^{(1)} - z)^{-1} = \sum_{r=0}^{\infty} (H_0^{(1)} - z)^{-1} \left(-P(\delta)VP(\delta) (H_0^{(1)} - z)^{-1} \right)^r \quad (48)$$

where $H_0^{(1)} = P(\delta)H_0$ and $z \in C_1$. It easily follows from (35) that

$$\left\| (H_0^{(1)}(\vec{\mathcal{Z}}) - z)^{-1} \right\| < \frac{8}{\tau l} k^{-2l+1+40\mu\delta}. \quad (49)$$

Hence,

$$\left\| (H^{(1)}(\vec{\mathcal{Z}}) - z)^{-1} \right\| < \frac{16}{\tau l} k^{-2l+1+40\mu\delta} \quad (50)$$

for sufficiently large k . Substituting the series into the formula $\mathcal{E}^{(1)}(\vec{\mathcal{Z}}) = -\frac{1}{2\pi i} \oint_{C_1} (H^{(1)}(\vec{\mathcal{Z}}) - z)^{-1} dz$ and integrating term-wise, we arrive at (42). Estimates (43) easily follow from (49) and the obvious inequality $\|P(\delta)\|_1 \leq (2k^\delta)^4$. It follows $\mathcal{E}^{(1)} = \mathcal{E}_0 + O(k^{-2l+1+44\mu\delta})$.

This means that there is a single eigenvalue of $H^{(1)}(\vec{\varkappa})$ inside C_1 . In a similar way (using (36), (38) and $V = 0$) we obtain the formula for the eigenvalue and (40), (41), for details see [K]. To prove (44) we consider the operator $A = VP(\delta) \left(H_0^{(1)} - z\right)^{-1}$ and represent it as $A = A_0 + A_1 + A_2$, where $A_0 = (P(\delta) - \mathcal{E}_0(\vec{\varkappa})) A (P(\delta) - \mathcal{E}_0(\vec{\varkappa}))$, $A_1 = (P(\delta) - \mathcal{E}_0(\vec{\varkappa})) A \mathcal{E}_0(\vec{\varkappa})$, $A_2 = \mathcal{E}_0(\vec{\varkappa}) A (P(\delta) - \mathcal{E}_0(\vec{\varkappa}))$. It is easy to see that $\mathcal{E}_0(\vec{\varkappa}) A \mathcal{E}_0(\vec{\varkappa}) = 0$ because of $V = 0$. Note that

$$\oint_{C_1} \left(H_0^{(1)} - z\right)^{-1} A_0^r dz = 0,$$

since the integrand is a holomorphic function inside C_1 . Therefore,

$$G_r^{(1)}(\vec{\varkappa}) = \frac{(-1)^{r+1}}{2\pi i} \sum_{j_1, \dots, j_r=0,1,2, \ j_1^2 + \dots + j_r^2 \neq 0} \oint_{C_1} \left(H_0^{(1)} - z\right)^{-1} A_{j_1} \dots A_{j_r} dz.$$

At least one of indices in each term is equal to 1, 2. We take into account that $(A_2)_{\mathbf{s}\mathbf{s}'} = (A_1)_{\mathbf{s}'\mathbf{s}} = 0$ if $\mathbf{s} \neq 0$ and $A_{\mathbf{s}\mathbf{s}'} = 0$ if $\|\vec{p}_{\mathbf{s}-\mathbf{s}'}\| > Q$. It follows that $G_r^{(1)}(\vec{\varkappa})_{\mathbf{s}\mathbf{s}'}$ can differ from zero only if $rQ \geq \|\vec{p}_{\mathbf{s}}\| + \|\vec{p}_{\mathbf{s}'}\|$.

□

It will be shown (Corollary 3.8) that coefficients $g_r^{(1)}(\vec{\varkappa})$ and operators $G_r^{(1)}(\vec{\varkappa})$ can be analytically extended into the complex $\frac{\tau}{16}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$ as functions of φ and to the complex $\frac{\tau}{8}k^{-(40\mu+1)\delta}$ -neighborhood of k as functions of $\varkappa$, estimates (40), (41), (43) being preserved. Now, we use formulae (36), (39) to extend $\lambda^{(1)}(\vec{\varkappa}) = \lambda^{(1)}(\varkappa, \varphi)$ as an analytic function. Obviously, series (39) is differentiable. Using Cauchy integral we get the following lemma.

Lemma 3.5. *Under conditions of Theorem 3.3 the following estimates hold when φ is in $\omega^{(1)}(k, \delta, \tau)$ or its complex $\frac{\tau}{32}k^{-(40\mu+1)\delta}$ -neighborhood and $\varkappa$ is in the complex $\frac{\tau}{16}k^{-40\mu\delta}$ -neighborhood of $\varkappa = k$:*

$$\lambda^{(1)}(\vec{\varkappa}) = \varkappa^{2l} + O\left(k^{-2l+(80\mu+6)\delta}\right), \quad (51)$$

$$\frac{\partial \lambda^{(1)}}{\partial \varkappa} = 2l\varkappa^{2l-1} + O\left(k^{-2l+(120\mu+6)\delta}\right), \quad \frac{\partial \lambda^{(1)}}{\partial \varphi} = O\left(k^{-2l+(120\mu+7)\delta}\right), \quad (52)$$

$$\begin{aligned} \frac{\partial^2 \lambda^{(1)}}{\partial \varkappa^2} &= 2l(2l-1)\varkappa^{2l-2} + O\left(k^{-2l+(160\mu+6)\delta}\right), \\ \frac{\partial^2 \lambda^{(1)}}{\partial \varkappa \partial \varphi} &= O\left(k^{-2l+(160\mu+7)\delta}\right), \quad \frac{\partial^2 \lambda^{(1)}}{\partial \varphi^2} = O\left(k^{-2l+(160\mu+8)\delta}\right). \end{aligned} \quad (53)$$

3.3 Geometric Considerations

In this section we prove Lemma 3.1 and its corollary. However, we will prove a version of this lemma for a complex set of φ . We need this complex version for further steps. Lemma 3.1 is a simple corollary of the result proven in this section. We will use the notation $|\mathbf{a}|_{\mathbb{R}}^2 := (\mathbf{a}, \mathbf{a})_{\mathbb{R}}$ where $(\mathbf{a}, \mathbf{b})_{\mathbb{R}} := a_1 b_1 + a_2 b_2$ when $\mathbf{a}, \mathbf{b} \in \mathbb{C}^2$. It is easy to see that $|\vec{\mathcal{Z}}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2$ is an analytic extension in $\varkappa$ and φ of

$$|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{m}}|^2 = \varkappa^2 + p_{\mathbf{m}}^2 + 2\varkappa p_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}})$$

defined for real $\varkappa, \varphi$. Note that $|\cdot|$ is the canonical norm in $\mathbb{C}$ or $\mathbb{R}^2$. For every fixed $k \geq 1$ and $\frac{1}{32} \leq \tau \leq 32$, we describe the resonance set $\mathcal{O}^{(1)} = \mathcal{O}^{(1)}(k, \tau)$ of $\varphi \in \mathbb{C}$. We put

$$\mathcal{O}^{(1)}(k, \tau) := \cup_{\mathbf{m} \in \tilde{\Omega}(\delta) \setminus \{0\}} \mathcal{O}_{\mathbf{m}}(k, \tau), \quad (54)$$

where

$$\begin{aligned} \mathcal{O}_{\mathbf{m}}(k, \tau) &:= \{\varphi \in \mathbb{C} : \left| |\vec{k} + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| \leq \tau k^{1-40\mu\delta} \} = \\ &\{\varphi \in \mathbb{C} : \left| p_{\mathbf{m}}^2 + 2k p_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}}) \right| \leq \tau k^{1-40\mu\delta} \}. \end{aligned} \quad (55)$$

In most cases parameter τ will be equal to 1. But sometimes we will use different choice of τ . It easily follows from the definition (55) and the estimate (27) that for any $\varkappa \in \mathbb{C}$ such that $|\varkappa - k| \leq 1$ and any $\varphi \in \mathcal{O}_{\mathbf{m}}(k, \tau)$ we have

$$\left| |p_{\mathbf{m}}^2 + 2\varkappa p_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}})| - |p_{\mathbf{m}}^2 + 2k p_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}})| \right| \leq \frac{\tau}{4} k^{1-40\mu\delta}, \quad (56)$$

provided $2(1 + 40\mu)\delta \leq 1$ and $k \geq 800$ which will be assumed in what follows.

Let $\mathcal{W}_0 := \{\varphi \in \mathbb{C} : |\Im \varphi| < 1\}$. We introduce a complex non-resonant set:

$$\mathcal{W}^{(1)}(k, \tau) := \mathcal{W}_0 \setminus \mathcal{O}^{(1)}(k, \tau). \quad (57)$$

Clearly, it is open. We also note that the set $\mathcal{O}^{(1)} \cap [0, 2\pi]$ is symmetric, i.e. $\mathcal{O}^{(1)} \cap [0, 2\pi] + \pi \pmod{2\pi} = \mathcal{O}^{(1)} \cap [0, 2\pi]$, since $\varphi_{-\mathbf{m}} = \varphi_{\mathbf{m}} + \pi$. We define $\omega^{(1)}(k, \delta, \tau)$ as a real part of $\mathcal{W}^{(1)}(k, \delta, \tau)$:

$$\omega^{(1)}(k, \delta, \tau) = \mathcal{W}^{(1)}(k, \tau) \cap [0, 2\pi]. \quad (58)$$

Lemma 3.6. *Let φ be in $\mathcal{W}^{(1)}(k, \tau)$, then*

$$\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| \geq \tau k^{1-40\mu\delta} \text{ for all } \mathbf{m} \in \tilde{\Omega}(\delta) \setminus \{0\}. \quad (59)$$

If φ is in the complex $k^{-(40\mu+1)\delta}$ -neighborhood of $\mathcal{W}^{(1)}(k, \tau)$ and $\varkappa \in \mathbb{C} : |\varkappa - k| < \frac{\tau}{8} k^{-40\mu\delta}$. Then, for $\vec{\mathcal{Z}} = \varkappa(\cos \varphi, \sin \varphi)$ the following estimate holds:

$$\left| |\vec{\mathcal{Z}}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| \geq \frac{\tau}{2} k^{1-40\mu\delta} \text{ for all } \mathbf{m} \in \tilde{\Omega}(\delta) \setminus \{0\}. \quad (60)$$

The lemma easily follows from (55) and (56).

Corollary 3.7. *Parts 1 and 2 of Lemma 3.1 hold.*

Corollary 3.8. *Coefficients $g_r^{(1)}(\vec{\kappa})$ and operators $G_r^{(1)}(\vec{\kappa})$ can be analytically extended into the complex $\frac{\tau}{16}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$ as functions of φ and to the complex $\frac{\tau}{16}k^{-(40\mu+1)\delta}$ -neighborhood of k as functions of κ , estimates (40), (41), (43) being preserved.*

Lemma 3.9. *The measure of the resonance set $\mathcal{O}^{(1)} \cap [0, 2\pi]$ satisfies the estimate:*

$$\text{meas}(\mathcal{O}^{(1)} \cap [0, 2\pi]) \leq Ck^{-37\delta\mu}. \quad (61)$$

Corollary 3.10. *Part 3 of Lemma 3.1 holds.*

Proof.

Let $\mathbf{m} \neq 0$ and $\varphi_{\mathbf{m}}^{\pm}$ be two (mod 2π) solutions of the equation

$$p_{\mathbf{m}}^2 + 2kp_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}}) = 0.$$

Obviously, $\varphi_{\mathbf{m}}^{\pm} - \varphi_{\mathbf{m}} = \pm \frac{\pi}{2} + O(k^{-1+\delta})$. Put

$$\Phi_{\mathbf{m}}^{\pm} := \{\varphi \in \mathbb{C} : |\varphi - \varphi_{\mathbf{m}}^{\pm}| \leq \tau k^{-39\delta\mu}\}.$$

Then, taking into account (28), it is not difficult to see that $\mathcal{O}_{\mathbf{m}} \subset \cup_{\pm, j \in \mathbb{Z}} (\Phi_{\mathbf{m}}^{\pm} + 2\pi j)$. Thus,

$$\text{meas}(\mathcal{O}^{(1)} \cap [0, 2\pi]) \leq 4\tau k^{-39\delta\mu} (8k^{\delta})^4 \leq Ck^{-37\delta\mu}. \quad (62)$$

□

Proof of Corollary 3.2. Let $C_1 := \{z \in \mathbb{C} : |z - k^{2l}| = \frac{\tau}{4}k^{2l-1-40\mu\delta}\}$ be the contour around eigenvalue k^{2l} of the unperturbed operator $H_0(\vec{k})$. Then it follows from (59) that for any $\varphi \in \mathcal{W}^{(1)}(k, \tau)$, $\mathbf{m} \in \tilde{\Omega}(\delta) \setminus \{0\}$, and $z : |z - k^{2l}| \leq \frac{\tau}{4}k^{2l-1-40\mu\delta}$ we have

$$\begin{aligned} ||\vec{k} + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} - z| &\geq ||\vec{k} + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} - k^{2l}| - \frac{\tau l}{4}k^{2l-1-40\mu\delta} \geq \\ &\tau l(1 - O(k^{\delta-1}))k^{2l-1-40\mu\delta} - \frac{\tau l}{4}k^{2l-1-40\mu\delta} \geq \frac{\tau l}{4}k^{2l-1-40\mu\delta}, \end{aligned} \quad (63)$$

for sufficiently large k . For $\mathbf{m} = 0$ the estimate follows from the definition of C_1 .

3.4 Isoenergetic Surface for Operator $H^{(1)}$

Lemma 3.11. *1. For every sufficiently large λ , $\lambda := k^{2l}$, and φ in the real $\frac{\tau}{32}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$, there is a unique $\kappa^{(1)}(\lambda, \varphi)$ in the interval $I_1 := [k - \frac{\tau}{32}k^{-40\mu\delta}, k + \frac{\tau}{32}k^{-40\mu\delta}]$, such that*

$$\lambda^{(1)}(\vec{\kappa}^{(1)}(\lambda, \varphi)) = \lambda, \quad \vec{\kappa}^{(1)}(\lambda, \varphi) := \kappa^{(1)}(\lambda, \varphi)\vec{v}(\varphi). \quad (64)$$

2. Furthermore, there exists an analytic in φ continuation of $\varkappa^{(1)}(\lambda, \varphi)$ to the complex $\frac{\tau}{32}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$ such that $\lambda^{(1)}(\vec{\varkappa}^{(1)}(\lambda, \varphi)) = \lambda$. Function $\varkappa^{(1)}(\lambda, \varphi)$ can be represented as $\varkappa^{(1)}(\lambda, \varphi) = k + h^{(1)}(\lambda, \varphi)$, where

$$|h^{(1)}| = O(k^{-4l+1+(80\mu+6)\delta}), \quad (65)$$

$$\frac{\partial h^{(1)}}{\partial \varphi} = O(k^{-4l+1+(120\mu+7)\delta}), \quad \frac{\partial^2 h^{(1)}}{\partial \varphi^2} = O(k^{-4l+1+(160\mu+8)\delta}), \quad (66)$$

$$\frac{\partial \varkappa^{(1)}}{\partial \lambda} = \frac{1}{2lk^{2l-1}} (1 + O(k^{-4l+1+(120\mu+6)\delta})). \quad (67)$$

Proof.

1. Let us prove existence of $\varkappa^{(1)}(\lambda, \varphi)$. By Theorem 3.3, there exists an eigenvalue $\lambda^{(1)}(\vec{\varkappa})$, given by (39), for all $\varkappa$ in the interval I_1 . Let $\mathcal{L}^{(1)}(\varphi) := \{\lambda^{(1)}(\vec{\varkappa}) : \varkappa \in I_1\}$. Using the definition of I_1 , (45), and continuity of $\lambda^{(1)}(\vec{\varkappa})$ is continuous in $\varkappa$, we easily obtain $\mathcal{L}^{(1)}(\varphi) \supset [k^{2l} - t, k^{2l} + t]$, $t = c_1 k^{2l-1-40\mu\delta}$, $0 < c_1 \neq c_1(k)$. Hence, there exists a $\varkappa^{(1)}$ such that $\lambda^{(1)}(\vec{\varkappa}^{(1)}) = k^{2l}$, $\varkappa^{(1)} \in I_1$.

Now we show that there is only one $\varkappa^{(1)}$ in the interval I_1 satisfying (64). Indeed, by (52), $\frac{\partial \lambda^{(1)}(\vec{\varkappa})}{\partial \varkappa} \geq 2lk^{2l-1}(1 + o(1))$. This implies that $\lambda^{(1)}(\vec{\varkappa})$ is monotone with respect to $\varkappa$ in I_1 . Thus, there is only one $\varkappa \in I_1$ satisfying (64).

2. We consider $\lambda^{(1)}(\vec{\varkappa}(\varphi))$ as a function of complex variable $\varkappa$ in the disc $|\varkappa - k| < \frac{\tau}{32}k^{-40\mu\delta}$. Taking into account (51) and applying Rouché's theorem, we obtain that for any φ in $\frac{\tau}{32}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, \delta, \tau)$ there exists unique value of $\varkappa^{(1)}(\varphi)$ such that $|\varkappa^{(1)}(\varphi) - k| < \frac{\tau}{32}k^{-40\mu\delta}$ and $\lambda^{(1)}(\vec{\varkappa}^{(1)}(\varphi)) = \lambda := k^{2l}$. Actually,

$$|\varkappa^{(1)}(\varphi) - k| < k^{-4l+1+(80\mu+6)\delta}. \quad (68)$$

Then it follows from (52) and implicit function theorem that $\varkappa^{(1)}(\varphi)$ is locally analytic. Combined with uniqueness this implies global analyticity.

The estimate (65) follows from (68). Applying standard arguments with the Cauchy formula we obtain (66). Using (52) we get (67).

□

Let us consider the set of points in $\mathbb{R}^2$ given by the formula: $\vec{\varkappa} = \vec{\varkappa}^{(1)}(\varphi)$, $\varphi \in \omega^{(1)}(k, \delta, \tau)$. By Lemma 3.11 this set of points is a slightly disturbed circle with holes, see Fig. 1. All the points of this curve satisfy the equation $\lambda^{(1)}(\vec{\varkappa}^{(1)}(\lambda, \varphi)) = k^{2l}$. We call it isoenergetic surface of the operator $H^{(1)}$ and denote by $\mathcal{D}_1(\lambda)$, see figure 1. The “radius” $\varkappa^{(1)}(\lambda, \varphi)$ of $\mathcal{D}_1(\lambda)$ monotonously increases with λ , see (67).

3.5 Preparation for Step II. Construction of the Second Non-resonant Set

3.5.1 Model Operator for Step II

Here we will describe an operator PHP , see (75), which will be used for constructing perturbation series in the second step. The operator PHP has a block structure, the size of blocks being of order k^δ .

Let r_1 be some fixed number $2 < r_1$. An upper bound on r_1 we will introduce in Step II. We defined $\mathcal{O}_{\mathbf{m}}$ by formula (55) for all $\mathbf{m}$: $0 < \|\vec{p}_{\mathbf{m}}\| \leq 4k^\delta$. Now we define $\mathcal{O}_{\mathbf{m}}$ by the formula

$$\begin{aligned} \mathcal{O}_{\mathbf{m}}(k, \tau) := \{ \varphi \in \mathbb{C} : \quad & \left| |\vec{k} + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| \leq \tau k^{-40\mu\delta} \} = \\ & \{ \varphi \in \mathbb{C} : \quad |p_{\mathbf{m}}^2 + 2kp_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}})| \leq \tau k^{-40\mu\delta} \}. \end{aligned} \quad (69)$$

for $\mathbf{m}$: $4k^\delta < \|\vec{p}_{\mathbf{m}}\| \leq k^{r_1}$. Note that the right-hand part in the inequality here is smaller than the corresponding one in (55). Obviously, $\mathcal{O}_{\mathbf{m}}$ contains the whole interval $[0, 2\pi)$ for sufficiently small $p_{\mathbf{m}}$. As in Step I let $\varphi_{\mathbf{m}}^\pm$ be two (mod 2π) solutions of the equation

$$p_{\mathbf{m}}^2 + 2kp_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}}) = 0. \quad (70)$$

Lemma 3.12. *The set $\mathcal{O}_{\mathbf{m}}(k, \tau)$ has the following properties:*

1. If $p_{\mathbf{m}} > 4k$, then $\mathcal{W}_0 \cap \mathcal{O}_{\mathbf{m}}(k, \tau) = \emptyset$.
2. If $k^{-1-39\mu\delta} \leq p_{\mathbf{m}} \leq 4k$ and $|4k^2 - p_{\mathbf{m}}^2| > 4\tau k^{-40\mu\delta}$, then $\mathcal{O}_{\mathbf{m}} \subset \cup_{\pm, j \in \mathbb{Z}} (\Phi_{\mathbf{m}}^\pm + 2\pi j)$, where

$$\Phi_{\mathbf{m}}^\pm := \left\{ \varphi \in \mathbb{C} : \quad |\varphi - \varphi_{\mathbf{m}}^\pm| \leq \frac{\tau k^{-1-40\mu\delta}}{p_{\mathbf{m}} \sqrt{1 - p_{\mathbf{m}}^2 (2k)^{-2}}} \right\},$$

and $\Phi_{\mathbf{m}}^+ \cap \Phi_{\mathbf{m}}^- = \emptyset$.

3. If $|4k^2 - p_{\mathbf{m}}^2| \leq 4\tau k^{-40\mu\delta}$, then $\mathcal{O}_{\mathbf{m}} \subset \cup_{\pm, j \in \mathbb{Z}} (\Phi_{\mathbf{m}}^\pm + 2\pi j)$, where

$$\Phi_{\mathbf{m}}^\pm := \{ \varphi \in \mathbb{C} : \quad |\varphi - \varphi_{\mathbf{m}}^\pm| \leq 32\tau k^{-1-20\mu\delta} \}.$$

In the proof we use the Taylor series with respect to φ for $|\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2$ near its zeros, see Appendix 1.

Let $\varphi_0 \in [0, 2\pi) \setminus \mathcal{O}^{(1)}(k, 8)$, where $\mathcal{O}^{(1)}(k, 8)$ is given by (54). We define $\mathcal{M}(\varphi_0) \subset \mathbb{Z}^2$ as follows:

$$\mathcal{M}(\varphi_0) := \{ \mathbf{m} : \quad 0 < \|\vec{p}_{\mathbf{m}}\| \leq k^{r_1} \text{ and } \varphi_0 \in \mathcal{O}_{\mathbf{m}}(k, 1) \}. \quad (71)$$

We will also need a larger set

$$\mathcal{M}'(\varphi_0) := \{\mathbf{m} : 0 < \|\vec{p}_{\mathbf{m}}\| \leq 2k^{r_1} \text{ and } \varphi_0 \in \mathcal{O}_{\mathbf{m}}(k, 1)\}.$$

In fact, $\mathcal{M}(\varphi_0)$, $\mathcal{M}'(\varphi_0)$ do not include $\mathbf{m} : \|\vec{p}_{\mathbf{m}}\| < 4k^\delta$, since $\varphi_0 \in [0, 2\pi) \setminus \mathcal{O}^{(1)}(k, 8)$.

We split $\mathcal{M}(\varphi_0)$ into two components $\mathcal{M} := \mathcal{M}_1 \cup \mathcal{M}_2$. By definition, $\mathbf{m} \in \mathcal{M}_1$ if

$$\min_{\mathbf{m}' \in \mathcal{M}'(\varphi_0), \mathbf{m}' \neq \mathbf{m}} \|\vec{p}_{\mathbf{m}-\mathbf{m}'}\| > k^\delta.$$

Let $\mathcal{M}_2 = \mathcal{M} \setminus \mathcal{M}_1$. Next, let $\tilde{\mathcal{M}}_{\mathbf{m}}$ be $(k^\delta/3)$ -neighborhood of $\mathbf{m}$ in $\|\cdot\|$ norm:

$$\tilde{\mathcal{M}}_{\mathbf{m}} := \{\mathbf{n} : \|\vec{p}_{\mathbf{n}-\mathbf{m}}\| < k^\delta/3 \text{ for a given } \mathbf{m} \in \mathcal{M}(\varphi_0)\},$$

Obviously,

$$\tilde{\mathcal{M}}_{\mathbf{m}}(\varphi_0) \cap \tilde{\mathcal{M}}_{\mathbf{m}'}(\varphi_0) = \emptyset, \quad \text{for any } \mathbf{m} \in \mathcal{M}_1 \text{ and } \mathbf{m}' \in \mathcal{M}', \quad \mathbf{m}' \neq \mathbf{m}.$$

Let $\tilde{\mathcal{M}}_1(\varphi_0)$ be $(k^\delta/3)$ -neighborhood of $\mathcal{M}_1$ in $\|\cdot\|$ norm:

$$\tilde{\mathcal{M}}_1(\varphi_0) := \cup_{\mathbf{m} \in \mathcal{M}_1(\varphi_0)} \tilde{\mathcal{M}}_{\mathbf{m}}(\varphi_0) = \{\mathbf{n} : \|\vec{p}_{\mathbf{n}-\mathbf{m}}\| < k^\delta/3 \text{ for some } \mathbf{m} \in \mathcal{M}_1(\varphi_0)\}.$$

Let us introduce an equivalence relation in $\mathcal{M}'$. We say $\mathbf{m}_0 \sim \mathbf{m}'_0$ if there is a sequence $\mathbf{m}_j \in \mathcal{M}'$, $j = 1, \dots, J$, such that $\min_{k < j} \|\vec{p}_{\mathbf{m}_j - \mathbf{m}_k}\| \leq k^\delta$ for all $j = 1, \dots, J$ and $\mathbf{m}_J = \mathbf{m}'_0$. We denote the equivalence class containing $\mathbf{m} \in \mathcal{M}_2$ by $\mathcal{M}_2^{(\mathbf{m})}$. By definition of $\mathcal{M}_2$ such equivalence class contains at least one more element. In the next lemma we prove that an equivalence class contains no more than 4 elements. Namely in this lemma the restriction $l > 1$ plays a crucial role.

Lemma 3.13. *Let $\mathbf{m}_0 \in \mathcal{M}_2$ and $\mathbf{m}_j \in \mathcal{M}'$, $j = 1, \dots, J$, are such that all $\mathbf{m}_j$, $j = 0, \dots, J$, are different and $\min_{k < j} \|\vec{p}_{\mathbf{m}_j - \mathbf{m}_k}\| \leq k^\delta$ for all $j = 1, \dots, J$. Then, $1 \leq J \leq 3$.*

The proof is in Appendix 2.

Obviously, for any pair $\mathbf{m}, \mathbf{m}' \in \mathcal{M}_2$ either $\mathcal{M}_2^{(\mathbf{m})} = \mathcal{M}_2^{(\mathbf{m}')} or $\mathcal{M}_2^{(\mathbf{m})} \cap \mathcal{M}_2^{(\mathbf{m}')} = \emptyset$. We can enumerate different equivalence classes $\mathcal{M}_2^{(\mathbf{m})}$ by an index j and denote them by $\mathcal{M}_2^j$, $j = 1, \dots, J_0$. By construction, $\mathcal{M}_2 \subset \cup_{j=1}^{J_0} \mathcal{M}_2^j \subset \mathcal{M}'$.$

Let $\tilde{\mathcal{M}}_2^j$ be $(k^\delta/3)$ -neighborhood of $\mathcal{M}_2^j$ in $\|\cdot\|$ norm:

$$\tilde{\mathcal{M}}_2^j(\varphi_0) := \{\mathbf{n} : \|\vec{p}_{\mathbf{n}-\mathbf{m}}\| < k^\delta/3 \text{ for an } \mathbf{m} \in \mathcal{M}_2^j(\varphi_0)\}.$$

Obviously,

$$\begin{aligned} \tilde{\mathcal{M}}_2^j &= \cup_{\mathbf{m} \in \mathcal{M}_2^j} \tilde{\mathcal{M}}_{\mathbf{m}}, \\ \tilde{\mathcal{M}}_2^j \cap \tilde{\mathcal{M}}_2^{j'} &= \emptyset, \text{ when } j \neq j', \end{aligned}$$

$$\tilde{\mathcal{M}}_2^j \cap \tilde{\mathcal{M}}_{\mathbf{m}} = \emptyset, \text{ when } \mathbf{m} \in \mathcal{M}_1.$$

Let

$$\begin{aligned}\tilde{\mathcal{M}}_2 &= \cup_{j=1}^{J_0} \tilde{\mathcal{M}}_2^j, \\ \tilde{\mathcal{M}} &= \tilde{\mathcal{M}}_1 \cup \tilde{\mathcal{M}}_2.\end{aligned}$$

Moreover, $||| \cdot |||$ distance between these sets is greater than $\frac{1}{3}k^\delta$. It is easy to see that $\tilde{\mathcal{M}} \subset \mathcal{M}'$. Hence, the number of elements in $\tilde{\mathcal{M}}$ does not exceed ck^{4r_1} .

We consider the diagonal projection P corresponding to $\tilde{\mathcal{M}}(\varphi_0)$:

$$P(\varphi_0)_{\mathbf{m}\mathbf{m}} = \begin{cases} 1, & \text{when } \mathbf{m} \in \tilde{\mathcal{M}}(\varphi_0), \\ 0, & \text{otherwise.} \end{cases}$$

We consider $PH(\vec{\mathcal{A}}^{(1)}(\varphi))P : PL_2(\mathbb{Z}^2) \rightarrow PL_2(\mathbb{Z}^2)$ for $\varphi \in \mathbb{C}$, $|\varphi - \varphi_0| < k^{-2-\delta(40\mu+1)}$. Since $\varphi_0 \in [0, 2\pi) \setminus \mathcal{O}^{(1)}(k, 8)$, perturbation series (39), (42) converge in the disc.

By construction, the set $\tilde{\mathcal{M}}(\varphi_0)$ is split into several nonintersecting components:

$$\tilde{\mathcal{M}}(\varphi_0) = \left(\cup_{\mathbf{m} \in \mathcal{M}_1} \tilde{\mathcal{M}}_{\mathbf{m}} \right) \cup \left(\cup_j \tilde{\mathcal{M}}_2^j \right). \quad (72)$$

Obviously,

$$P = \sum_{\mathbf{m} \in \mathcal{M}_1} P_{\mathbf{m}} + \sum_j P_2^j, \quad (73)$$

where $P_{\mathbf{m}}, P_2^j$ are diagonal projectors corresponding to the sets $\tilde{\mathcal{M}}_{\mathbf{m}}$ and $\tilde{\mathcal{M}}_2^j$, the projectors being orthogonal. Considering (30) and taking into account that $Q < k^\delta/3$ for sufficiently large k , we readily show:

$$P_{\mathbf{m}}VP_{\mathbf{m}'} = P_{\mathbf{m}}VP_2^j = P_2^jVP_{\mathbf{m}} = P_2^jVP_2^{j'} = 0, \text{ when } \mathbf{m}, \mathbf{m}' \in \mathcal{M}_1, \mathbf{m} \neq \mathbf{m}', j \neq j'. \quad (74)$$

Therefore,

$$PHP = \sum_{\mathbf{m} \in \mathcal{M}_1} P_{\mathbf{m}}HP_{\mathbf{m}} + \sum_j P_2^jHP_2^j. \quad (75)$$

Since (31) holds for any $\mathbf{m} \in \tilde{\Omega}(\delta) \setminus \{0\}$, we have $\tilde{\mathcal{M}}(\varphi_0) \cap \tilde{\Omega}(\delta) = \emptyset$. This means that the $||| \cdot |||$ -distance between $\tilde{\mathcal{M}}(\varphi_0)$ and $\Omega(\delta)$ is no less than $3k^\delta$. Hence,

$$P_{\mathbf{m}}VP(\delta) = P(\delta)VP_{\mathbf{m}} = P(\delta)VP_2^j = P_2^jVP(\delta) = 0. \quad (76)$$

3.5.2 Estimates for the Resolvent of the Model Operator

In the next lemma we use the restriction $l > 1$ for the first time. In fact, we need this restriction only in the second step of the procedure.

Lemma 3.14. *Let $\varphi_0 \in \omega^{(1)}(k, 8)$.*

1. If $\mathbf{m} \in \mathcal{M}_1(\varphi_0) : p_{\mathbf{m}} > 4k^\delta, |2k - p_{\mathbf{m}}| \geq 1$, then, the operator

$$(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_{\mathbf{m}})^{-1}$$

has no more than one pole in the disk $|\varphi - \varphi_0| < 2k^{-2-\delta(40\mu+1)}$. The following estimate holds:

$$\left\| (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_{\mathbf{m}})^{-1} \right\| < ck^{-2l+1}\varepsilon_0^{-1}, \quad \varepsilon_0 = \min\{\varepsilon, k^{-2-(40\mu+1)\delta}\}, \quad (77)$$

when φ is in the smaller disk $|\varphi - \varphi_0| < k^{-2-\delta(40\mu+1)}$, ε being the distance from φ to the nearest pole of the operator.

2. If $\mathbf{m} \in \mathcal{M} : |2k - p_{\mathbf{m}}| < 1$, then, in fact $\mathbf{m} \in \mathcal{M}_1$ and the operator

$$(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_{\mathbf{m}})^{-1}$$

has no more than two poles in the disk $|\varphi - \varphi_0| < 2k^{-2-\delta(40\mu+1)}$. The following estimate holds:

$$\left\| (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_{\mathbf{m}})^{-1} \right\| < ck^{-2l}\varepsilon_0^{-2}, \quad \varepsilon_0 = \min\{\varepsilon, k^{-2-\delta(40\mu+1)}\}, \quad (78)$$

when φ is in the smaller disk $|\varphi - \varphi_0| < k^{-2-\delta(40\mu+1)}$, ε being the distance from φ to the nearest pole of the operator.

3. If $\mathbf{m} \in \mathcal{M} : p_{\mathbf{m}} < 4k^\delta$, then, in fact $\mathbf{m} \in \mathcal{M}_1$ and the operator

$$(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_{\mathbf{m}})^{-1}$$

has no more than one pole in the disk $|\varphi - \varphi_0| < 2k^{-2-\delta(40\mu+1)}$. The following estimate holds:

$$\left\| (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_{\mathbf{m}})^{-1} \right\| < 8k^{-2l+1}p_{\mathbf{m}}^{-1}\varepsilon_0^{-1}, \quad \varepsilon_0 = \min\{\varepsilon, k^{-2l+1+\delta}\}, \quad (79)$$

when φ is in the smaller disk $|\varphi - \varphi_0| < k^{-2-\delta(40\mu+1)}$, ε being the distance from φ to the nearest pole of the operator.

4. The operator $(P_2^j(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_2^j)^{-1}$ has no more than four poles in the disk $|\varphi - \varphi_0| < 2k^{-2-\delta(40\mu+1)}$. The following estimate holds:

$$\left\| (P_2^j(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_2^j)^{-1} \right\| < ck^{-2l-2-120\mu\delta}\varepsilon_0^{-4}, \quad \varepsilon_0 = \min\{\varepsilon, k^{-1-40\mu\delta}\}, \quad (80)$$

when φ is in the smaller disk $|\varphi - \varphi_0| < k^{-2-\delta(40\mu+1)}$, ε being the distance from φ to the nearest pole of the operator.

Corollary 3.15. *Let $\varphi_0 \in \omega^{(1)}(k, 8)$. Then, the operator $(P(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P)^{-1}$ has no more than $64k^{4r_1}$ poles in the disk $|\varphi - \varphi_0| < 2k^{-2-\delta(40\mu+1)}$. The following estimate holds:*

$$\left\| (P(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P)^{-1} \right\| < ck^{\mu r_1} \varepsilon_0^{-1} + ck^{-2l} \varepsilon_0^{-2} + ck^{-2l-2} \varepsilon_0^{-4}, \quad \varepsilon_0 = \min\{\varepsilon, k^{-2l+1+\delta}\}, \quad (81)$$

when φ is in the smaller disk $|\varphi - \varphi_0| < k^{-2-\delta(40\mu+1)}$, ε being the distance from φ to the nearest pole of the operator.

Indeed, the number of blocks in PHP (see (75)) does not exceed $16k^{4r_1}$ (the number of elements in $\Omega(r_1)$). The resolvent of each block has no more than four poles. Therefore, the resolvent of PHP has no more than $64k^{4r_1}$ poles. Using (77)-(80) and, using that $p_{\mathbf{m}} > k^{-\mu r_1}$ in (79), we obtain the corollary.

Corollary 3.16. *If $\varepsilon = k^{-r'_1}$, $r'_1 \geq \mu r_1$, then*

$$\left\| (P(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P)^{-1} \right\| < ck^{4r'_1}, \quad (82)$$

$$\left\| (P(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P)^{-1} \right\|_1 < ck^{4r'_1+4r_1}. \quad (83)$$

The first formula follows from (81). The second formula follows from the fact that the dimension of P does not exceed k^{4r_1} .

Proof.

1. Let $|2k - p_{\mathbf{m}}| \geq 1$, $p_{\mathbf{m}} > 4k^\delta$. Clearly only the case $p_{\mathbf{m}} < 4k$ is significant, since otherwise $\mathcal{O}_{\mathbf{m}}(k, 1)$ cannot intersect the disc $|\varphi - \varphi_0| < k^{-2-40(\mu+1)\delta}$ by Lemma 3.12. It is easy to see that the set $\mathcal{O}_{\mathbf{m}}(k, 1)$ consists of two separate discs $\mathcal{O}_{\mathbf{m}}^\pm(k, 1)$, the distance between them being greater than $ck^{-1/2}$. Let us assume for definiteness $\varphi_0 \in \mathcal{O}_{\mathbf{m}}^+(k, 1)$. This means the disc $|\varphi - \varphi_0| < k^{-2-\delta(40\mu+1)}$ does not intersect $\mathcal{O}_{\mathbf{m}}^-(k, 1)$. Let us first show that the operator

$$(P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P_{\mathbf{m}})^{-1} \quad (84)$$

has exactly one pole inside $\mathcal{O}_{\mathbf{m}}^+(k, 1)$, which is, in fact, inside $\mathcal{O}_{\mathbf{m}}^+(k, 1/4)$. Note that $\vec{\mathcal{Z}}^{(1)}(\varphi)$ is defined in $\mathcal{O}_{\mathbf{m}}^+(k, 1)$, since the size of $\mathcal{O}_{\mathbf{m}}^+(k, 1)$ is much less than that of any circle in $\mathcal{O}^{(1)}$. It satisfies the estimate $\vec{\mathcal{Z}}^{(1)}(\varphi) = \vec{k}(\varphi) + o(k^{-2})$ in $\mathcal{O}_{\mathbf{m}}^+$. If $\varphi_0 \in \mathcal{O}_{\mathbf{m}}^+(k, 1) \setminus \mathcal{O}_{\mathbf{m}}^+(k, 1/4)$, then the estimates $|\vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}+\mathbf{q}}|^2 - k^2| > \frac{1}{4}k^{-40\mu\delta}$, hold for $0 \leq \|\vec{p}_{\mathbf{q}}\| < k^\delta$ (see definition of $\mathcal{M}_1(\varphi_0)$) and can be extended to the $(k^{-2-(40\mu+1)\delta})$ -neighborhood of φ_0 ($\frac{1}{4}$ becomes $\frac{1}{8}$). Thus,

$$\left\| (P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P_{\mathbf{m}})^{-1} \right\| < ck^{-2l+2+40\mu\delta} \quad (85)$$

when $|\varphi - \varphi_0| < k^{-2-(40\mu+1)\delta}$, $\varphi_0 \in \mathcal{O}_{\mathbf{m}}^+(k, 1) \setminus \mathcal{O}_{\mathbf{m}}^+(k, 1/4)$.

Clearly the resolvent (85) does not have poles in the set $|\varphi - \varphi_0| < k^{-2-(40\mu+1)\delta}$. The estimate (77) with $\varepsilon_0 = k^{-2-(40\mu+1)\delta}$ follows from (85) and Hilbert identity.

Now, suppose that $\varphi_0 \in \mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. The function $|\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} - k^{2l}$ has a single zero inside $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Using Rouché's theorem, we obtain that $|\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} - k^{2l}$ also has a single zero inside $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Note that the following inequality holds in $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$ for $0 < \|\vec{p}_{\mathbf{q}}\| < k^\delta$:

$$||\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}+\mathbf{q}}|_{\mathbb{R}}^{2l} - k^{2l}| > \frac{1}{4}k^{2l-2-40\mu\delta}.$$

Indeed, if $||\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}+\mathbf{q}}|_{\mathbb{R}}^{2l} - k^{2l}| \leq \frac{1}{4}k^{2l-2-40\mu\delta}$ for some $\mathbf{q} \neq (0, 0)$ and $\varphi \in \mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$, then

$$|2(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}, \vec{p}_{\mathbf{q}})_{\mathbb{R}} + p_{\mathbf{q}}^2| < \frac{1}{2}k^{-40\mu\delta}. \quad (86)$$

Considering that the size of $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$ is $\frac{k^{-1-40\mu\delta}}{p_{\mathbf{m}}\sqrt{1-p_{\mathbf{m}}^2(2k)^{-2}}}(1+o(1))$ and that $p_{\mathbf{m}} > 4k^\delta > 4p_{\mathbf{q}}/2\pi$, we obtain the inequality analogous to (86) for φ_0 with $\frac{3}{4}$ instead of $\frac{1}{2}$. This contradicts to the assumption $\varphi_0 \in \mathcal{M}_1$. Thus, the following inequality holds for all $\mathbf{q} : \|\vec{p}_{\mathbf{q}}\| < k^\delta$ including $\mathbf{q} = (0, 0)$:

$$||\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}+\mathbf{q}}|_{\mathbb{R}}^{2l} - k^{2l}| > \frac{1}{4}k^{2l-2-40\mu\delta},$$

when φ is on the boundary of $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Hence, the resolvent

$$(P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P_{\mathbf{m}})^{-1}$$

of the free operator $P_{\mathbf{m}}H_0$ has exactly one pole inside $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$ and

$$\left\| (P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P_{\mathbf{m}})^{-1} \right\| \leq 4k^{-2l+2+40\mu\delta}, \quad (87)$$

when φ is on the boundary on the disc $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Considering that the dimension of $P_{\mathbf{m}}$ does not exceed $16k^{4\delta}$ we obtain:

$$\left\| (P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P_{\mathbf{m}})^{-1} \right\|_1 \leq 64k^{-2l+2+\delta(40\mu+4)}. \quad (88)$$

It remains to prove the analogous result for the perturbed operator H . We introduce the determinant

$$D(\varphi) = \det \left(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I)^{-1}P_{\mathbf{m}} \right).$$

Obviously, $D(\varphi) = \det(I + A)$, where $I, A : P_{\mathbf{m}}L_2(\mathbb{Z}^2) \rightarrow P_{\mathbf{m}}L_2(\mathbb{Z}^2)$

$$A(\varphi) = P_{\mathbf{m}}V \left(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right)^{-1} P_{\mathbf{m}}.$$

Taking into account that

$$D(\varphi) = \frac{\det \left(P_{\mathbf{m}} \left(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right) P_{\mathbf{m}} \right)}{\det \left(P_{\mathbf{m}} \left(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right) P_{\mathbf{m}} \right)},$$

we see that $D(\varphi)$ is a meromorphic function inside $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Next, we employ a well-known inequality for the determinants, see [36]:

$$|\det(I + A) - \det(I + B)| \leq \|A - B\|_1 \exp(\|A\|_1 + \|B\|_1 + 1), \quad A, B \in \mathbf{S}_1. \quad (89)$$

Putting $A = A(\varphi)$, $B = 0$, we obtain

$$|\det(I + A) - 1| \leq \|A\|_1 \exp(\|A\|_1 + 1).$$

It is easy to see that

$$\|A\|_1 \leq \|V\| \|P_{\mathbf{m}} \left(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right)^{-1} P_{\mathbf{m}}\|_1.$$

Considering the estimate (88) for the resolvent of the free operator, we obtain $\|A_1(\varphi)\|_1 < 1/200$ on the boundary of $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$ for sufficiently large k . By Rouché's theorem, $D(\varphi)$ has only one zero in $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Thus, $\det P_{\mathbf{m}} \left(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right) P_{\mathbf{m}}$ has exactly one zero in $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Using this, we immediately obtain that operator $\left(P_{\mathbf{m}} \left(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right) P_{\mathbf{m}} \right)^{-1}$ has one pole inside $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Considering the estimate for the free resolvent and using Hilbert identity, we immediately obtain,

$$\left\| \left(P_{\mathbf{m}} \left(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right) P_{\mathbf{m}} \right)^{-1} \right\| \leq 8k^{-2l+2+40\mu\delta} \quad (90)$$

for all φ on the boundary of $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. Taking into account that the size of $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$ does not exceed $k^{-1-40\mu\delta}$, we obtain:

$$\left\| \left(P_{\mathbf{m}} \left(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I \right) P_{\mathbf{m}} \right)^{-1} \right\| \leq 8k^{-2l+2+40\mu\delta} (k^{-1-40\mu\delta} / \varepsilon) \quad (91)$$

when $\varphi \in \mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$ on the distance ε , from the pole. If $|\varphi - \varphi_0| < k^{-2-(40\mu+1)\delta}$, but $\varphi \notin \mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$, then φ is on the distance less than $k^{-2-(40\mu+1)\delta}$ from the boundary of $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$, since φ_0 is inside $\mathcal{O}_{\mathbf{m}}^+(k, \frac{1}{4})$. The estimate (90) holds on the boundary and stable with respect to such a small perturbation of φ . Thus, estimate (77) is proven.

2. Let $\mathbf{m} \in \mathcal{M}$, $|2k - p_{\mathbf{m}}| < 1$. Then, $\mathcal{O}_{\mathbf{m}}^+$ and $\mathcal{O}_{\mathbf{m}}^-$ can overlap. The case $\varphi_0 \in \mathcal{O}_{\mathbf{m}}(k, 1) \setminus \mathcal{O}_{\mathbf{m}}(k, 1/4)$ we consider in the same way as for $|2k - p_{\mathbf{m}}| \geq 1$. Suppose $\varphi_0 \in \mathcal{O}_{\mathbf{m}}(k, 1/4)$. Combining $\left| |\vec{k}(\varphi_0) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| < \frac{1}{4}k^{1-40\mu\delta}$ with $|2k - p_{\mathbf{m}}| < 1$, we obtain that the vectors $2\vec{k}(\varphi)$ and $-\vec{p}_{\mathbf{m}}$ are close:

$$|2\vec{k}(\varphi_0) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 < 5k.$$

Therefore, $(\vec{k}(\varphi_0) + \vec{p}_{\mathbf{m}}, \vec{p}_{\mathbf{q}})_{\mathbb{R}} = -(\vec{k}(\varphi_0), \vec{p}_{\mathbf{q}})_{\mathbb{R}} + O(k^{1/2+\delta})$ for all $0 < \|\vec{p}_{\mathbf{q}}\| < k^\delta$. Considering that the size of $\mathcal{O}_{\mathbf{m}}$ does not exceed $ck^{-1-20\mu\delta}$ (Lemma 3.12) and the distance between $\mathcal{O}_{\mathbf{m}}^+$ and $\mathcal{O}_{\mathbf{m}}^-$ is $O(k^{-1/2})$, we obtain the analogous estimate for all φ in $\mathcal{O}_{\mathbf{m}}$:

$$(\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}, \vec{p}_{\mathbf{q}})_{\mathbb{R}} = -(\vec{k}(\varphi), \vec{p}_{\mathbf{q}})_{\mathbb{R}} + O(k^{1/2+\delta}) \quad \text{for all } 0 < \|\vec{p}_{\mathbf{q}}\| < k^\delta.$$

It immediately follows:

$$|\vec{k}(\varphi) + \vec{p}_{\mathbf{m}+\mathbf{q}}|_{\mathbb{R}}^2 - |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 = |\vec{k}(\varphi) - \vec{p}_{\mathbf{q}}|_{\mathbb{R}}^2 - |\vec{k}(\varphi)|_{\mathbb{R}}^2 + O(k^{1/2+\delta})$$

for all $0 < \|\vec{p}_{\mathbf{q}}\| < k^\delta$. The size of $\mathcal{O}_{\mathbf{m}}$ is much smaller than that of $\mathcal{O}_{-\mathbf{q}}$ and $\mathcal{O}_{\mathbf{m}}$ is not completely in $\mathcal{O}_{-\mathbf{q}}$. Hence,

$$\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}+\mathbf{q}}|_{\mathbb{R}}^2 - |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 \right| > \frac{1}{2}k^{1-40\mu\delta}$$

for all $\varphi \in \mathcal{O}_{\mathbf{m}}$. Considering that $\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| \leq \frac{1}{2}k^{-40\mu\delta}$ in $\mathcal{O}_{\mathbf{m}}$, we obtain

$$\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}+\mathbf{q}}|_{\mathbb{R}}^2 - k^2 \right| > \frac{1}{2}k^{1-40\mu\delta}, \text{ when } \varphi \in \mathcal{O}_{\mathbf{m}}.$$

In particular, $\mathbf{m} \in \mathcal{M}_1(\varphi_0)$. Considering that $\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| = \frac{1}{2}k^{-40\mu\delta}$ on the boundary of $\mathcal{O}_{\mathbf{m}}$, we obtain (87) and (88). Considering as before, we show that the resolvent (84) has at most two poles inside $\mathcal{O}_{\mathbf{m}}$. It follows that

$$\left\| (P_{\mathbf{m}} (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}I) P_{\mathbf{m}})^{-1} \right\| \leq k^{-2l+2+40\mu\delta} (ck^{-1-20\mu\delta}/\varepsilon)^2$$

when φ is on the distance ε from the pole.

3. Let $\mathbf{m} \in \mathcal{M}$, $0 < p_{\mathbf{m}} \leq 4k^\delta$, $\mathbf{m} \notin \Omega(\delta)$. The case $\varphi_0 \in \mathcal{O}_{\mathbf{m}}(k, 1) \setminus \mathcal{O}_{\mathbf{m}}(k, \frac{1}{4})$ is considered the same way as in the previous steps, see (85). From now on we assume $\varphi_0 \in \mathcal{O}_{\mathbf{m}}(k, \frac{1}{4})$. There is an eigenvalue $\lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}})$ of $P_{\mathbf{m}}H(\vec{\mathcal{Z}}^{(1)}(\varphi))P_{\mathbf{m}}$ given by the perturbation series. Indeed,

$$\left| |\vec{k}(\varphi_0) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| < \frac{1}{4}k^{-40\mu\delta},$$

since $\varphi_0 \in \mathcal{O}_{\mathbf{m}}(k, \frac{1}{4})$. Considering that $\varphi_0 \notin \mathcal{O}^{(1)}(k, 8)$, we easily obtain that $\left|(\vec{k}(\varphi_0), \vec{p}_{\mathbf{q}})_{\mathbb{R}}\right| \gtrsim k^{1-40\mu\delta}$ for all $\mathbf{q} \in \Omega(\delta) \setminus \{0\}$. Taking into account that and $p_{\mathbf{m}} \leq 4k^\delta$ we arrive at the estimate:

$$\left|\left|\vec{k}(\varphi_0) + \vec{p}_{\mathbf{m}+\mathbf{q}}\right|_{\mathbb{R}}^2 - k^2\right| \gtrsim k^{1-40\mu\delta}$$

for all $\mathbf{q} \in \Omega(\delta) \setminus \{0\}$ and any $\varphi_0 \in \omega^{(1)}(k, 8) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{4})$. It follows $\mathbf{m} \in \mathcal{M}_1$. By Lemma 3.11, $\vec{\mathcal{Z}}^{(1)}(\varphi)$ is defined in $\frac{1}{4}k^{-(40\mu+1)\delta}$ -neighborhood of $\omega^{(1)}(k, 8)$ which we denote by $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4})$. It is easy to show that the estimates similar to the last two hold for $\vec{\mathcal{Z}}^{(1)}(\varphi)$, $\varphi \in \tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$. Therefore,

$$\left|\left|\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}\right|_{\mathbb{R}}^{2l} - k^{2l}\right| \lesssim \frac{1}{2}k^{2l-2-40\mu\delta}, \quad (92)$$

$$\left|\left|\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}+\mathbf{q}}\right|_{\mathbb{R}}^{2l} - k^{2l}\right| \gtrsim k^{2l-1-40\mu\delta} \quad (93)$$

for all $\mathbf{q} \in \Omega(\delta) \setminus \{0\}$. It follows from the last two estimates that the perturbation series for $\lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}})$ and $\lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi))$ converge. Both are holomorphic functions of φ in $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$. Using Rouché's theorem, it is not difficult to show (for details see Appendix 3, Lemma 10.1) that the equation

$$\lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}) = k^{2l} + \varepsilon_0, \quad |\varepsilon_0| \leq p_{\mathbf{m}}k^\delta, \quad (94)$$

has no more than two solutions $\varphi^\pm(\varepsilon_0)$ in the $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$. They satisfy the estimates:

$$|\varphi^\pm(\varepsilon_0) - \varphi_{\mathbf{m}}^\pm| < 4k^{-2l+1+2\delta}. \quad (95)$$

Considering that $\varphi_{\mathbf{m}}^\pm = \varphi_{\mathbf{m}} \pm \pi/2 + O(k^{-1+\delta})$, we see that the distance between two solutions is approximately equal to π . For any $\varphi \in \tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$ satisfying the estimate $|\varphi - \varphi_{\mathbf{m}}^\pm| < k^{-\delta}$,

$$\frac{\partial}{\partial \varphi} \lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}) = \pm 2lp_{\mathbf{m}}k^{2l-1}(1 + o(1)), \quad (96)$$

for details see Appendix 3, Lemma 10.2. Therefore (for details see Appendix 3, Lemma 10.3),

$$|\lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}) - k^{2l}| \geq k^{2l-1}p_{\mathbf{m}}\varepsilon \quad (97)$$

if $\varphi \in \tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$ is outside $\tilde{\mathcal{O}}_{\mathbf{m},\varepsilon}^+ \cup \tilde{\mathcal{O}}_{\mathbf{m},\varepsilon}^-$, here and below $\tilde{\mathcal{O}}_{\mathbf{m},\varepsilon}^\pm$ are the open discs of the radius ε , $0 < \varepsilon < k^{-2l+1+\delta}$, centered at $\varphi^\pm(0)$. It is shown in Appendix 3, Lemma 10.4 that

$$\left\|(\lambda^{(1)}(\vec{y}(\varphi)) - k^{2l})(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l})P_{\mathbf{m}})^{-1}\right\| \leq 8, \quad \vec{y}(\varphi) := \vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}, \quad (98)$$

for any φ in $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$.

If $|\varphi - \varphi_0| < 2k^{-2-\delta(40\mu+1)}$ and $\varphi_0 \in \omega(k, \delta, 8) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{4})$, then $\varphi \in \tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$ and, hence, (97), (98) hold. Now (79) easily follows from (97) and (98).

4. Let, now, $\mathbf{m}, \mathbf{m}'$ be two elements from the same set $\mathcal{M}_2^j$. It means that there exist elements $\vec{p}_{\mathbf{q}_i} \in \Omega(\delta)$, $i = 1, \dots, I$, such that $\mathbf{m}' = \mathbf{m} + \sum_{i=1}^I \mathbf{q}_i$ and $\mathbf{m} + \sum_{i=1}^s \mathbf{q}_i \in \mathcal{M}_2$ for any $1 \leq s \leq I$. We have proved in Lemma 3.13 that $I \leq 3$. Next, we consider $\tilde{\mathcal{O}}^j = \cup_{\mathbf{m} \in \mathcal{M}_2^j} \mathcal{O}_{\mathbf{m}}$. Each connected component of $\tilde{\mathcal{O}}^j$ contains no more than four discs. We have proven above that all $\mathbf{m} : p_{\mathbf{m}} < 4k^\delta$ or $|2k - p_{\mathbf{m}}| < 1$ belong to $\mathcal{M}_1$. Using this fact and applying Lemma 3.12 we see that the size of each component does not exceed $o(k^{-1-40\mu\delta})$. Considering as above, we show that

$$\left\| (P_{\mathbf{m}} (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l} I) P_{\mathbf{m}})^{-1} \right\| \leq 2k^{-2l+2+40\mu\delta} \quad (99)$$

for all φ on the boundary of $\tilde{\mathcal{O}}^j$. Each component contains no more than four poles of the resolvent. It follows that

$$\left\| (P_{\mathbf{m}} (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l} I) P_{\mathbf{m}})^{-1} \right\| \leq ck^{-2l+2+40\mu\delta} (k^{-1-40\mu\delta} / \varepsilon_0)^4$$

with $\varepsilon_0 = \min\{\varepsilon, k^{-1-40\mu\delta}\}$, when φ is on the distance ε from the poles.

□

3.5.3 Resonant and Nonresonant Sets for Step II

We divide $[0, 2\pi)$ into $[2\pi k^{2+\delta(40\mu+1)}] + 1$ intervals $\Delta_j^{(1)}$ with the length not bigger than $k^{-2-\delta(40\mu+1)}$. If a particular interval belongs to $\mathcal{O}^{(1)}(k, 8)$ we ignore it; otherwise, let $\varphi_0(j) \notin \mathcal{O}^{(1)}(k, 8)$ be a point inside the $\Delta_j^{(1)}$. Let

$$\mathcal{W}_j^{(1)} = \{\varphi \in \mathcal{W}^{(1)} : |\varphi - \varphi_j^{(0)}| < 2k^{-2-\delta(40\mu+1)}\}.$$

Clearly, neighboring sets $\mathcal{W}_j^{(1)}$ overlap (because of the multiplier 2 in the inequality), they cover the $2k^{-2-\delta(40\mu+1)}$ -neighborhood of $\omega(k, 8)$. We denote this neighborhood by $\hat{\mathcal{W}}^{(1)}(k, 2)$. For each φ in the neighborhood there is a j such that $|\varphi - \varphi_j^{(0)}| < k^{-2-\delta(40\mu+1)}$. We consider the poles of the operator $(P(\varphi_j^{(0)}) (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l}) P(\varphi_j^{(0)}))^{-1}$ in a $\mathcal{W}_j^{(1)}$ and denote them by φ_{jm} , $m = 1, \dots, M_j$. By Corollary 3.15, $M_j < 64k^{4r_1}$. Next, let $\mathcal{O}_{jm}^{(2)}$ be the disc of the radius $k^{-r'_1}$ around φ_{jm} , $r'_1 > \mu r_1$.

Definition 3.17. *The set*

$$\mathcal{O}^{(2)} = \cup_{jm} \mathcal{O}_{jm}^{(2)} \quad (100)$$

we call the second resonant set. The set

$$\mathcal{W}^{(2)} = \hat{\mathcal{W}}^{(1)}(k, 2) \setminus \mathcal{O}^{(2)} \quad (101)$$

is called the second nonresonant set. The set

$$\omega^{(2)} = \mathcal{W}^{(2)} \cap [0, 2\pi) \quad (102)$$

is called the second real nonresonant set.

Lemma 3.18. *Let $r'_1 > \mu r_1$, $\varphi \in \mathcal{W}^{(2)}$ and $\varkappa \in \mathbb{C} : |\varkappa - \varkappa^{(1)}(\varphi)| < k^{-4r'_1-2l+1-\delta}$. Then,*

$$\left\| (P(H(\vec{\varkappa}(\varphi)) - k^{2l}I)P)^{-1} \right\| < ck^{4r'_1}, \quad (103)$$

$$\left\| (P(H(\vec{\varkappa}(\varphi)) - k^{2l}I)P)^{-1} \right\|_1 < ck^{4r'_1+4r_1}, \quad (104)$$

where P is the projection (73) corresponding to the interval $\Delta_j^{(1)}$ containing $\Re\varphi$.

Proof. For $\vec{\varkappa} = \vec{\varkappa}^{(1)}(\varphi)$ the lemma follows immediately from the definition of $\mathcal{W}^{(2)}$ and Corollary 3.16. Considering the Hilbert identity, it is easy to see that estimates (82) and (83) are stable with respect to perturbation of $\varkappa^{(1)}$ of order $k^{-4r'_1-2l+1-\delta}$. This stability ensure (103) and (104).

□

By total size of the set $\mathcal{O}^{(2)}$ we mean the sum of the sizes of its connected components.

Lemma 3.19. *Let $r'_1 \geq (\mu + 4)r_1$. Then, the size of each connected component of $\mathcal{O}^{(2)}$ is less than $128k^{4r_1-r'_1}$. The total size of $\mathcal{O}^{(2)}$ is less than $128k^{2+\delta(40\mu+1)+4r_1-r'_1}$, where $2 + \delta(40\mu + 1) + 4r_1 - r'_1 < 0$.*

Corollary 3.20. *If a connected component of $\mathcal{O}^{(2)}$ intersects $[0, 2\pi)$ or its $\frac{1}{2}k^{-2-\delta(40\mu+1)}$ -neighborhood, then it is strictly inside $\tilde{\mathcal{W}}^{(1)}$.*

Proof. Indeed, each set $\mathcal{W}_j^{(1)}$ contains no more than $64k^{4r_1}$ discs $\mathcal{O}_{jm}$. Therefore, the size of $\mathcal{O}^{(2)} \cap \mathcal{W}_j^{(1)}$ is less than $128k^{-r'_1+4r_1}$. Considering that $128k^{-r'_1+4r_1}$ is much smaller than the length of $\Delta_j^{(1)}$, we obtain that there is no connected components which go across the whole set $\mathcal{W}_j^{(1)}$ and the size of each connected component of $\mathcal{O}^{(2)}$ is less than $128k^{4r_1-r'_1}$. Considering that $j < k^{2+\delta(40\mu+1)}$, we obtain the required estimate for the total size of $\mathcal{O}^{(2)}$.

□

We will also need the estimates for the resolvent in the neighborhood of $\mathbf{m} = 0$. From the definition of $\vec{\varkappa}^{(1)}(\varphi)$ we obtain the following lemma.

Lemma 3.21. *Let $\varphi \in \mathcal{W}_j^{(1)}$ and C_2 be the circle $|z - k^{2l}| = (2ck^{4r'_1})^{-1}$ (where we use the same constant as in Lemma 3.18). Then*

$$\left\| (P(\delta)(H(\vec{\varkappa}^{(1)}(\varphi)) - z)P(\delta))^{-1} \right\| \leq 8ck^{4r'_1}.$$

Proof. This estimate is sufficiently obvious and can be obtained in many different ways. Here though we will use the construction which we often will keep in mind when stating similar estimates in what follows. We apply for the z variable the "squeezing" arguments which we used in the proof of Lemma 3.14 for the variable φ . Namely, by (50) (or rather its identical analogues for complex $\vec{\mathcal{Z}}^{(1)}(\varphi)$),

$$\left\| (P(\delta)(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P(\delta))^{-1} \right\| \leq \frac{16}{\tau l} k^{-2l+1+40\mu\delta},$$

when $|z - k^{2l}| = \frac{\tau l}{4} k^{2l-1-40\mu\delta}$. Let us show that analytic function $\det(P(\delta)(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P(\delta))$ has the single simple zero $z = k^{2l}$ inside the circle C_1 . Indeed, consider

$$D(\varphi) = \det \left(P(\delta) (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - zI) (P(\delta) (H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - zI) P(\delta))^{-1} \right).$$

Obviously, $D(\varphi) = \det(I + A)$, where $I, A : P(\delta)L_2(\mathbb{Z}^2) \rightarrow P(\delta)L_2(\mathbb{Z}^2)$,

$$A(\varphi) = P(\delta)V (H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - zI)^{-1} P(\delta).$$

Obviously, $D(\varphi)$ is a meromorphic function inside C_1 . Next, we employ (89) putting $A = A(\varphi)$, $B = 0$. We obtain $|\det(I + A) - 1| \leq \|A\|_1 \exp(\|A\|_1 + 1)$. By (49), $\|A(\varphi)\|_1 < ck^{-2l+1+\delta(40\mu+4)}$ when $z \in C_1$. By Rouché's theorem, $D(\varphi)$ has only one zero in C_1 . Thus, $\det P(\delta) (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - zI) P(\delta)$ has one zero in C_1 . Using this, we immediately obtain that operator $(P(\delta) (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - zI) P(\delta))^{-1}$ has exactly one pole inside C_1 , the pole being at the point $z = k^{2l}$. Using the maximum principle, we obtain the required estimate on the circle C_2 . $\square$

We also notice that the statement of the Lemma 3.18 still holds (with $2c$ instead of c) if we use $z \in C_2$ instead of k^{2l} . Thus, if we put

$$P_j := P(\varphi_j(0)), \quad \tilde{P}_j := P_j + P(\delta), \quad (105)$$

then (notice that $P_j V P(\delta) = 0$ by (76))

$$\left\| \left(\tilde{P}_j (H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z) \tilde{P}_j \right)^{-1} \right\| \leq 8ck^{4r'_1}, \quad |z - k^{2l}| = (2ck^{4r'_1})^{-1}, \quad \varphi \in \mathcal{W}_j^{(1)}. \quad (106)$$

Here we also used Lemma 3.21. At last, considering from the beginning the discs $\mathcal{O}_{jm}^{(2)}$ with radius $\frac{1}{2}k^{-r'_1}$ instead of $k^{-r'_1}$ one can easily see that similar estimates (with probably larger constants) hold in $k^{-r'_1-\delta}$ -neighborhood of $\mathcal{W}^{(2)}$.

4 Step II

4.1 Operator $H^{(2)}$. Perturbation Formulas

Let $P(r_1)$ be an orthogonal projector onto $\Omega(r_1) := \{\mathbf{m} : \|\vec{p}_{\mathbf{m}}\| \leq k^{r_1}\}$ and $H^{(2)} = P(r_1)HP(r_1)$. From now on we assume

$$r'_1 = 40\mu r_1 + 2l, \quad 2 < r_1 < k^{\delta/8}. \quad (107)$$

We consider $H^{(2)}(\vec{\varkappa}^{(1)}(\varphi))$ as a perturbation of

$$\tilde{H}^{(1)} = \tilde{P}_j H(\vec{\varkappa}^{(1)}(\varphi)) \tilde{P}_j + \left(P(r_1) - \tilde{P}_j \right) H_0(\vec{\varkappa}^{(1)}(\varphi)) \left(P(r_1) - \tilde{P}_j \right), \quad (108)$$

where $\tilde{P}_j = P_j + P(\delta)$ and P_j is the projection P corresponding to the interval $\Delta_j^{(1)}$ containing φ . By (74), (76), the first term on the right-hand side of (108) has a block structure:

$$\tilde{P}_j H(\vec{\varkappa}^{(1)}(\varphi)) \tilde{P}_j = P(\delta) H P(\delta) + P H P = P(\delta) H P(\delta) + \sum_{\mathbf{m} \in \mathcal{M}_1} P_{\mathbf{m}} H P_{\mathbf{m}} + \sum_j P_2^j H P_2^j.$$

The second term in (108) is, obviously, diagonal. Thus, $\tilde{H}^{(1)}$ has a block-diagonal structure. Let W be the perturbation of $\tilde{H}^{(1)}$, i.e, $W = H^{(2)} - \tilde{H}^{(1)}$. It is easy to see that:

$$W = P(r_1) V P(r_1) - \tilde{P}_j V \tilde{P}_j. \quad (109)$$

By analogy with (36), (37),

$$g_r^{(2)}(\vec{\varkappa}) := \frac{(-1)^r}{2\pi i r} \text{Tr} \oint_{C_2} \left(W(\tilde{H}^{(1)}(\vec{\varkappa}) - zI)^{-1} \right)^r dz, \quad (110)$$

$$G_r^{(2)}(\vec{\varkappa}) := \frac{(-1)^{r+1}}{2\pi i} \oint_{C_2} (\tilde{H}^{(1)}(\vec{\varkappa}) - zI)^{-1} \left(W(\tilde{H}^{(1)}(\vec{\varkappa}) - zI)^{-1} \right)^r dz. \quad (111)$$

Theorem 4.1. *Suppose φ is in the real $k^{-r'_1-\delta}$ -neighborhood of $\omega^{(2)}(k, \delta, \tau)$ and $\varkappa \in \mathbb{R}$, $|\varkappa - \varkappa^{(1)}(\varphi)| \leq k^{-4r'_1-2l+1-\delta}$, $\vec{\varkappa} = \varkappa(\cos \varphi, \sin \varphi)$. Then, for sufficiently large $k > k_1(V, \delta, \tau)$ there exists a single eigenvalue of $H^{(2)}(\vec{\varkappa})$ in the interval $\varepsilon_2(k, \delta, \tau) = (k^{2l} - (2ck^{4r'_1})^{-1}, k^{2l} + (2ck^{4r'_1})^{-1})$. It is given by the absolutely converging series:*

$$\lambda^{(2)}(\vec{\varkappa}) = \lambda^{(1)}(\vec{\varkappa}) + \sum_{r=2}^{\infty} g_r^{(2)}(\vec{\varkappa}). \quad (112)$$

For coefficients $g_r^{(2)}(\vec{\varkappa})$ the following estimates hold:

$$|g_r^{(2)}(\vec{\varkappa})| < ck^{-2k^\delta Q^{-1}} k^{-(2l-2-40\mu\delta)(r-1)+2+8\delta}. \quad (113)$$

The corresponding spectral projection is given by the series:

$$\mathcal{E}^{(2)}(\vec{\varkappa}) = \mathcal{E}^{(1)}(\vec{\varkappa}) + \sum_{r=1}^{\infty} G_r^{(2)}(\vec{\varkappa}), \quad (114)$$

$\mathcal{E}^{(1)}(\vec{\varkappa})$ being the spectral projection of $H^{(1)}(\vec{\varkappa})$. The operators $G_r^{(2)}(\vec{\varkappa})$ satisfy the estimates:

$$\|G_r^{(2)}(\vec{\varkappa})\|_1 < ck^{-k^\delta Q^{-1}} k^{-(2l-2-40\mu\delta)r+1+4\delta}. \quad (115)$$

$$G_r^{(2)}(\vec{\varkappa})_{\text{ss}'} = 0, \quad \text{if } 10rk^\delta < \|\vec{p}_{\text{s}}\| + \|\vec{p}_{\text{s}'}\| \quad (116)$$

Corollary 4.2. *For the perturbed eigenvalue and its spectral projection the following estimates hold:*

$$\lambda^{(2)}(\vec{\mathcal{Z}}) = \lambda^{(1)}(\vec{\mathcal{Z}}) + O\left(k^{-2k^\delta Q^{-1}} k^{-2l+4+48\mu\delta}\right), \quad (117)$$

$$\|\mathcal{E}^{(2)}(\vec{\mathcal{Z}}) - \mathcal{E}^{(1)}(\vec{\mathcal{Z}})\|_1 < ck^{-k^\delta Q^{-1}} k^{-(2l-3-40\mu\delta-4\delta)}. \quad (118)$$

$$|\mathcal{E}^{(2)}(\vec{\mathcal{Z}})_{\mathbf{s}\mathbf{s}'}| < k^{-d^{(2)}(\mathbf{s}, \mathbf{s}')}, \quad \text{when } \|\vec{p}_{\mathbf{s}}\| > k^\delta \text{ or } \|\vec{p}_{\mathbf{s}'}\| > k^\delta, \quad (119)$$

$$d^{(2)}(\mathbf{s}, \mathbf{s}') = \frac{1}{10}(\|\vec{p}_{\mathbf{s}}\| + \|\vec{p}_{\mathbf{s}'}\|)k^{-\delta}(2l-2-40\mu\delta) + k^\delta Q^{-1} - 1 - 4\delta.$$

Formulas (117) and (118) easily follow from (112), (114) and (113) and (115). The estimate (119) follows from (114), (115) and (116). Indeed, using these estimates, we obtain $|\mathcal{E}^{(2)}(\vec{\mathcal{Z}}) - \mathcal{E}^{(1)}(\vec{\mathcal{Z}})|_{\mathbf{s}\mathbf{s}'} < k^{-d^{(2)}(\mathbf{s}, \mathbf{s}')}.$ Considering that $\mathcal{E}^{(1)}(\vec{\mathcal{Z}})_{\mathbf{s}\mathbf{s}'} = 0$ when $\|\vec{p}_{\mathbf{s}}\| > k^\delta$ or $\|\vec{p}_{\mathbf{s}'}\| > k^\delta$, we arrive at (119).

Proof. Let $\varphi \in \mathcal{W}_j^{(1)}$ for some j . Put $P'_j := P(r_1) - \tilde{P}_j$ (we will omit the index j in what follows). By (108), (109),

$$\tilde{H}^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi)) := \tilde{P}H(\vec{\mathcal{Z}}^{(1)}(\varphi))\tilde{P} + P'H_0(\vec{\mathcal{Z}}^{(1)}(\varphi))P', \quad W := P'VP' + P'V\tilde{P} + \tilde{P}VP'.$$

We will often omit $\vec{\mathcal{Z}}^{(1)}(\varphi)$ in the arguments when it cannot lead to confusion. By (106), we have

$$\|(\tilde{H}^{(1)} - zI)^{-1}\| < 8ck^{4r'_1}. \quad (120)$$

Let us consider the perturbation series

$$(H^{(2)} - z)^{-1} = \sum_{r=0}^{\infty} (\tilde{H}^{(1)} - z)^{-1} A^r, \quad (121)$$

$$A = -W(\tilde{H}^{(1)} - z)^{-1}.$$

To check the convergence it is enough to show that

$$\|A\| < ck^{-2l+2+40\mu\delta}. \quad (122)$$

Estimates (120), (122) yield:

$$\|(H^{(2)}(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)^{-1}\| < 16ck^{4r'_1}. \quad (123)$$

To prove (122) it suffice to establish the following estimates:

$$\begin{aligned} \|P'VP'(\tilde{H}^{(1)} - z)^{-1}\| &< ck^{-2l+2+40\mu\delta}, \\ \|P'V\tilde{P}(\tilde{H}^{(1)} - z)^{-1}\| &< ck^{-2l+2+40\mu\delta}, \\ \|\tilde{P}VP'(\tilde{H}^{(1)} - z)^{-1}\| &< ck^{-2l+2+40\mu\delta}. \end{aligned} \quad (124)$$

The first and the third inequalities in (124) are simple. They follow from the definition of P' and $\tilde{P}$ and identities

$$(\tilde{H}^{(1)} - z)^{-1}P' = P'(\tilde{H}^{(1)} - z)^{-1} = P'(H_0(\tilde{\mathcal{Z}}^{(1)}(\varphi)) - z)^{-1}P'.$$

Indeed, by definition, $(P')_{\mathbf{mm}} = 1$ if and only if $\varphi_j(0) \notin \mathcal{O}_{\mathbf{m}}(k, 8)$ and $\|\vec{p}_{\mathbf{m}}\| > k^\delta$. Therefore, $\|\tilde{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} - k^{2l}\| > k^{2l-2-40\mu\delta}$ when $(P')_{\mathbf{mm}} = 1$ and the estimate is stable with respect to perturbation of order $k^{-1-40\mu\delta}$. The first and third inequalities in (124) easily follow. Let us prove the second estimate. We represent $\tilde{P}(\tilde{H}^{(1)} - z)^{-1}$ as follows:

$$\begin{aligned} \tilde{P}(\tilde{H}^{(1)} - z)^{-1} = \\ \sum_{r=0}^{r_0} \left((H_0 - z)^{-1} \tilde{P} V \tilde{P} \right)^r (H_0 - z)^{-1} \tilde{P} + \left((H_0 - z)^{-1} \tilde{P} V \tilde{P} \right)^{r_0+1} (\tilde{H}^{(1)} - z)^{-1} \tilde{P}, \end{aligned}$$

where r_0 to be fixed later. Then,

$$\begin{aligned} \|P'V\tilde{P}(\tilde{H}^{(1)} - z)^{-1}\| &\leq \sum_{r=0}^{r_0} \|B_r\| + \left\| P'V \left((H_0 - z)^{-1} \tilde{P} V \tilde{P} \right)^{r_0+1} \right\| \|(\tilde{H}^{(1)} - z)^{-1} \tilde{P}\|, \\ B_r &:= P'V \left((H_0 - z)^{-1} \tilde{P} V \tilde{P} \right)^r (H_0 - z)^{-1} \tilde{P}. \end{aligned} \tag{125}$$

Note that $B_r = P'B_r\tilde{P}$ and matrix elements $(B_r)_{\mathbf{js}}$ are equal to zero if $\|\vec{p}_{\mathbf{j}} - \vec{p}_{\mathbf{s}}\| > Q(r+1)$ (see (30)). Thus, the only non-trivial elements $(B_r)_{\mathbf{js}}$ are such that

$$\mathbf{j} \in \Omega(r_1) \setminus \left(\tilde{\mathcal{M}}(\varphi_0) \cup \Omega(\delta) \right), \quad \mathbf{s} \in \tilde{\mathcal{M}}(\varphi_0) \cup \Omega(\delta), \quad \|\vec{p}_{\mathbf{j}} - \vec{p}_{\mathbf{s}}\| \leq Q(r+1).$$

Let $r : Q(r+1) \leq k^\delta/6$. It follows that $(B_r)_{\mathbf{js}} = 0$ if $\mathbf{s} \in \mathcal{M}$ or $\mathbf{s} = 0$, since such $\mathbf{s}$ have a distance greater than $\frac{1}{3}k^\delta$ from $\mathbf{j}$. If $\mathbf{s} \notin \mathcal{M}$ or $\mathbf{s} \neq 0$, then $\|\tilde{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{s}}|_{\mathbb{R}}^{2l} - z\| > k^{2l-2-40\mu\delta}$. Therefore, for $r : Q(r+1) \leq k^\delta/6$ we have:

$$\|B_r\| \leq (Ck^{-2l+2+40\mu\delta})^{r+1}, \quad \left\| P'V \left((H_0 - z)^{-1} \tilde{P} V \tilde{P} \right)^{r+1} \right\| \leq (Ck^{-2l+2+40\mu\delta})^{r+1},$$

with some absolute constant C . Now, we fix $r_0 := k^\delta/(6Q) - 1$. Then the condition $Q(r+1) \leq k^\delta/6$ is satisfied for all $r \leq r_0$ and

$$\|P'V\tilde{P}(\tilde{H}^{(1)} - z)^{-1}\| \leq \sum_{r=0}^{r_0} (Ck^{-2l+2+40\mu\delta})^{r+1} + (Ck^{-2l+2+40\mu\delta})^{r_0+1} 4ck^{4r'_1}.$$

Assuming that k is large enough (in particular, $\frac{(2l-2-40\mu\delta)k^\delta}{6Q} > 5r'_1$) we obtain the second inequality in (124).

To prove (115) we consider the operator $AP(\delta) = WP(\delta) \left(\tilde{H}^{(1)} - z \right)^{-1}$ and represent it as $AP(\delta) = A_0 + A_1 + A_2$, where $A_0 = (P(r_1) - \mathcal{E}^{(1)}(\vec{z})) A (P(r_1) - \mathcal{E}^{(1)}(\vec{z}))$, $A_1 = (P(r_1) - \mathcal{E}^{(1)}(\vec{z})) A \mathcal{E}^{(1)}(\vec{z})$, $A_2 = \mathcal{E}^{(1)}(\vec{z}) A (P(r_1) - \mathcal{E}^{(1)}(\vec{z}))$. Note that we have $\mathcal{E}^{(1)}(\vec{z}) W \mathcal{E}^{(1)}(\vec{z}) = 0$, because of (109). We see that

$$\oint_{C_2} \left(\tilde{H}^{(1)} - z \right)^{-1} A_0^r dz = 0,$$

since the integrand is a holomorphic function inside C_2 . Therefore,

$$G_r^{(2)}(\vec{z}) = \frac{(-1)^r}{2\pi i} \sum_{j_1, \dots, j_r=0,1,2, \ j_1^2+\dots+j_r^2 \neq 0} I_{j_1 \dots j_r}, \quad I_{j_1 \dots j_r} := \oint_{C_2} \left(\tilde{H}^{(1)} - z \right)^{-1} A_{j_1} \dots A_{j_r} dz. \quad (126)$$

At least one of indices in each term is equal to 1 or 2. Let us show that

$$\|A_2\|_1 < ck^{-k^\delta Q^{-1}} k^{-(2l-2-40\mu\delta)+1+4\delta}. \quad (127)$$

First, we notice that $\mathcal{E}^{(1)} W (P(r_1) - \mathcal{E}^{(1)}) = \mathcal{E}^{(1)} W P'$ by (109) and (76). It suffices to show that

$$\|\mathcal{E}^{(1)} W P'\|_1 < k^{-k^\delta Q^{-1}+1+4\delta}, \quad (128)$$

since $\|P' \left(\tilde{H}^{(1)} - z \right)^{-1}\| = \|P' (H_0 - z)^{-1}\| < k^{-(2l-2-40\mu\delta)}$ for $z \in C_2$. Indeed,

$$(\mathcal{E}^{(1)} W P')_{\mathbf{s}\mathbf{s}'} = \sum_{\mathbf{s}'': \|\vec{p}_{\mathbf{s}''}\| \leq k^\delta, \ \|\vec{p}_{\mathbf{s}''} - \mathbf{s}'\| \leq Q} \mathcal{E}_{\mathbf{s}\mathbf{s}''}^{(1)} W_{\mathbf{s}'' - \mathbf{s}'}$$

when $\|\vec{p}_{\mathbf{s}'}\| > k^\delta$ and it is equal to zero otherwise. Hence,

$$|(\mathcal{E}^{(1)} W P')_{\mathbf{s}\mathbf{s}'}| \leq \|W\| \sum_{\mathbf{s}'': k^\delta - Q \leq \|\vec{p}_{\mathbf{s}''}\| \leq k^\delta} \mathcal{E}_{\mathbf{s}\mathbf{s}''}^{(1)}$$

if $\|\vec{p}_{\mathbf{s}'}\| < k^\delta + Q$ and zero otherwise. Using (47), we obtain

$$|(\mathcal{E}^{(1)} W P')_{\mathbf{s}\mathbf{s}'}| < ck^{4\delta} \max_{\|\vec{p}_{\mathbf{s}''}\| > k^\delta - Q} k^{-d^{(1)}(\mathbf{s}, \mathbf{s}'')}. \quad (129)$$

It easily follows:

$$|(\mathcal{E}^{(1)} W P')_{\mathbf{s}\mathbf{s}'}| < ck^{4\delta} k^{-(2l-1-40\mu\delta)(k^\delta Q^{-1}-1+\|\vec{p}_{\mathbf{s}}\|Q^{-1})}$$

when $\|\vec{p}_{\mathbf{s}'}\| < k^\delta + Q$, and zero otherwise. It follows $\|\mathcal{E}^{(1)} W P'\| < ck^{-k^\delta Q^{-1}+1+4\delta}$. Considering that $\mathcal{E}^{(1)}$ is a one-dimensional projection, we obtain the same estimate for $\mathbf{S}_1$ -norm, namely, (128). Thus, we have proved (127). Let us estimate $I_{j_1 \dots j_r}$. Suppose

one of the indices is equal to 2. Substituting (127) into (126) and taking into account $\|\mathcal{E}^{(1)}(\tilde{H}^{(1)} - z)^{-1}\| < b_2^{-1}$, where b_2 is the radius of C_2 , we obtain:

$$\|I_{j_1 \dots j_r}\| < ck^{-k^\delta Q^{-1}} k^{-(2l-2-40\mu\delta)r+1+4\delta}.$$

Note that the operator A_1 is always followed by A_2 unless A_1 occupies the very last position in the product. Thus, it remains to consider the case $A_{j_1} \dots A_{j_r} = A_0^{r-1} A_1$. It is easy to see that

$$\left(\tilde{H}^{(1)} - z\right)^{-1} A_0^{r-1} A_1 = \left(\left(\tilde{H}^{(1)} - \bar{z}\right)^{-1} A_2(\bar{z}) A_0^{r-1}(\bar{z})\right)^*.$$

This implies the estimate for this case too. Therefore,

$$\|G_r^{(2)}(\vec{\varkappa})\| < ck^{-k^\delta Q^{-1}} k^{-(2l-2-40\mu\delta)r+1+4\delta}.$$

The same estimate can be written for the $\mathbf{S}_1$ norm of this operator, since $\mathcal{E}^{(1)}$ is one-dimensional.

Let us obtain the estimate for $g_r(\vec{\varkappa})$. Obviously,

$$g_r^{(2)}(\vec{\varkappa}) = \frac{(-1)^r}{2\pi i r} \sum_{j_1, \dots, j_r=0,1,2, \ j_1^2 + \dots + j_r^2 \neq 0} \text{Tr} \oint_{C_2} A_{j_1} \dots A_{j_r} dz. \quad (130)$$

Note that each term contains both A_1 and A_2 , since we compute the trace of the integral. Using (128), we obtain: $\|A_1\|_1 < cb_2^{-1} k^{-k^\delta Q^{-1}+1+4\delta}$. Combining this estimate with (127) and (122), we obtain (113) for $r \geq 2$. Finally, applying (110) in the case $r = 1$, we see that $g_1^{(2)}(\vec{\varkappa}) = 0$, since $\mathcal{E}^{(1)} W \mathcal{E}^{(1)} = 0$.

To prove (116) it's enough to notice that the biggest block of $\tilde{H}^{(1)}$ has the size not greater than $2k^\delta$.

□

It is easy to see that coefficients $g_r^{(2)}(\vec{\varkappa})$ and operators $G_r^{(2)}(\vec{\varkappa})$ can be analytically extended into the complex $k^{-r'_1-\delta}$ -neighborhood of $\omega^{(2)}$ (in fact, into $k^{-r'_1-\delta}$ -neighborhood of $\mathcal{W}^{(2)}$) as functions of φ and to the complex $(k^{-4r'_1-2l+1-\delta})$ -neighborhood of $\varkappa = \varkappa^{(1)}(\varphi)$ as functions of $\varkappa$, estimates (113), (117) being preserved. Now, we use formulae (110), (112) to extend $\lambda^{(2)}(\vec{\varkappa}) = \lambda^{(2)}(\varkappa, \varphi)$ as an analytic function. Obviously, series (112) is differentiable. Using Cauchy integral and Lemma 3.5 we get the following lemma.

Lemma 4.3. *Under conditions of Theorem 4.1 the following estimates hold when $\varphi \in \omega^{(2)}(k, \delta)$ or its complex $k^{-r'_1-\delta}$ -neighborhood and $\varkappa \in \mathbb{C} : |\varkappa - \varkappa^{(1)}(\varphi)| < k^{-4r'_1-2l+1-\delta}$:*

$$\lambda^{(2)}(\vec{\varkappa}) = \lambda^{(1)}(\vec{\varkappa}) + O\left(k^{-2k^\delta Q^{-1}} k^{-2l+4+48\mu\delta}\right), \quad (131)$$

$$\frac{\partial \lambda^{(2)}}{\partial \varkappa} = \frac{\partial \lambda^{(1)}}{\partial \varkappa} + O\left(k^{-2k^\delta Q^{-1}} k^{4r'_1+3+(48\mu+1)\delta}\right), \quad (132)$$

$$\frac{\partial \lambda^{(2)}}{\partial \varphi} = \frac{\partial \lambda^{(1)}}{\partial \varphi} + O\left(k^{-2k^\delta Q^{-1}} k^{r'_1-2l+4+(48\mu+1)\delta}\right), \quad (133)$$

$$\frac{\partial^2 \lambda^{(2)}}{\partial \varkappa^2} = \frac{\partial^2 \lambda^{(1)}}{\partial \varkappa^2} + O\left(k^{-2k^\delta Q^{-1}} k^{8r'_1+2l+2+(48\mu+2)\delta}\right), \quad (134)$$

$$\frac{\partial^2 \lambda^{(2)}}{\partial \varkappa \partial \varphi} = \frac{\partial^2 \lambda^{(1)}}{\partial \varkappa \partial \varphi} + O\left(k^{-2k^\delta Q^{-1}} k^{5r'_1+3+(48\mu+2)\delta}\right), \quad (135)$$

$$\frac{\partial^2 \lambda^{(2)}}{\partial \varphi^2} = \frac{\partial^2 \lambda^{(1)}}{\partial \varphi^2} + O\left(k^{-2k^\delta Q^{-1}} k^{2r'_1-2l+4+(48\mu+2)\delta}\right). \quad (136)$$

4.2 Isoenergetic Surface for Operator $H^{(2)}$

Lemma 4.4. 1. For every sufficiently large λ , $\lambda := k^{2l}$, and φ in the real $\frac{1}{2}k^{-r'_1-\delta}$ -neighborhood of $\omega^{(2)}(k, \delta, \tau)$, there is a unique $\varkappa^{(2)}(\lambda, \varphi)$ in the interval $I_1 := [\varkappa^{(1)}(\lambda, \varphi) - \frac{1}{2}k^{-4r'_1-2l+1-\delta}, \varkappa^{(1)}(\lambda, \varphi) + \frac{1}{2}k^{-4r'_1-2l+1\delta}]$, such that

$$\lambda^{(2)}(\vec{\varkappa}^{(2)}(\lambda, \varphi)) = \lambda, \quad \vec{\varkappa}^{(2)}(\lambda, \varphi) := \varkappa^{(2)}(\lambda, \varphi) \vec{v}(\varphi). \quad (137)$$

2. Furthermore, there exists an analytic in φ continuation of $\varkappa^{(2)}(\lambda, \varphi)$ to the complex $\frac{1}{2}k^{-r'_1-\delta}$ -neighborhood of $\omega^{(2)}(k, \delta, \tau)$ such that $\lambda^{(2)}(\vec{\varkappa}^{(2)}(\lambda, \varphi)) = \lambda$. Function $\varkappa^{(2)}(\lambda, \varphi)$ can be represented as $\varkappa^{(2)}(\lambda, \varphi) = \varkappa^{(1)}(\lambda, \varphi) + h^{(2)}(\lambda, \varphi)$, where

$$|h^{(2)}(\varphi)| = O\left(k^{-2k^\delta Q^{-1}} k^{-4l+5+48\mu\delta}\right), \quad (138)$$

$$\frac{\partial h^{(2)}}{\partial \varphi} = O\left(k^{-2k^\delta Q^{-1}} k^{r'_1-4l+5+49\mu\delta}\right), \quad \frac{\partial^2 h^{(2)}}{\partial \varphi^2} = O\left(k^{-2k^\delta Q^{-1}} k^{2r'_1-4l+5+50\mu\delta}\right). \quad (139)$$

Proof. The proof is completely analogous to that of Lemma 3.11, estimates (131)–(136) being used. $\square$

Let us consider the set of points in $\mathbb{R}^2$ given by the formula: $\vec{\varkappa} = \vec{\varkappa}^{(2)}(\varphi)$, $\varphi \in \omega^{(2)}(k, \delta, \tau)$. By Lemma 4.4 this set of points is a slight distortion of $\mathcal{D}_1$, see Fig. 2. All the points of this curve satisfy the equation $\lambda^{(2)}(\vec{\varkappa}^{(2)}(\varphi)) = k^{2l}$. We call it isoenergetic surface of the operator $H^{(2)}$ and denote by $\mathcal{D}_2$.

4.3 Preparation for Step III - Geometric Part. Properties of the Quasiperiodic Lattice

Let

$$\mathcal{S}(k, \varepsilon_0) = \left\{ \vec{\mathcal{R}} \in \mathbb{R}^2 : \left\| (H^{(1)}(\vec{\mathcal{R}}) - k^{2l})^{-1} \right\| > \varepsilon_0^{-1} \right\}. \quad (140)$$

In this section we prove that the number of the lattice points $\vec{\mathcal{R}}_0 + \vec{p}_{\mathbf{m}}$, $\|\vec{p}_{\mathbf{m}}\| < k^{r_1}$ in $\mathcal{S}(k, \varepsilon_0)$ does not exceed $Ck^{\frac{2r_1}{3}+1}$ when ε_0 is sufficiently small and $\vec{\mathcal{R}}_0$ is fixed. For this we split $\mathcal{S}$ into two subsets: “non-resonant” and “resonant”, the non-resonant set being just a vicinity of $\mathcal{D}_1(k^{2l})$. An estimate for the number of lattice points in the non-resonant set is proven in Lemma 4.8. An estimate for the number of lattice points in the resonant set is proven in Lemma 4.10. These estimates play an important role in the further construction.

4.3.1 General Lemmas

We consider $\vec{p}_{\mathbf{m}} = 2\pi(\mathbf{s}_1 + \alpha\mathbf{s}_2)$ with integer vectors $\mathbf{s}_j$ such that $|\mathbf{s}_j| \leq 4k^{r_1}$.

It is easy to see that there exists a pair $(q, p) \in \mathbb{Z}^2$ such that $0 < q \leq 4k^{r_1}$ and

$$|\alpha q + p| \leq 16k^{-r_1}. \quad (141)$$

We choose a pair (p, q) which gives the best approximation. In particular, p and q are mutually simple. Put $\epsilon_q := \alpha + \frac{p}{q}$. We have

$$k^{-2r_1\mu} \leq |\epsilon_q| \leq 16q^{-1}k^{-r_1}. \quad (142)$$

We write any $\mathbf{s}_2$ in the form

$$\mathbf{s}_2 = q\mathbf{s}'_2 + \mathbf{s}''_2 \quad (143)$$

with integer vectors $\mathbf{s}'_2$ and $\mathbf{s}''_2$, $0 \leq (\mathbf{s}''_2)_j < q$ for $j = 1, 2$. Hence, $|(\mathbf{s}'_2)_j| \leq 4k^{r_1}/q + 1$. It follows

$$(2\pi)^{-1}\vec{p}_{\mathbf{m}} = (\mathbf{s}_1 - p\mathbf{s}'_2) + \left(-\frac{p}{q}\mathbf{s}''_2 + \epsilon_q\mathbf{s}''_2\right) + \epsilon_q q\mathbf{s}'_2.$$

Denote $\mathbf{s} := \mathbf{s}_1 - p\mathbf{s}'_2$. Then $|\mathbf{s}| \leq 8k^{r_1}$. The number of different vectors $\tilde{\mathbf{s}} := -\frac{p}{q}\mathbf{s}''_2 + \epsilon_q\mathbf{s}''_2$ is not greater than $(2q)^2$. For each fixed pair $\tilde{\mathbf{s}}, \mathbf{s}$ we obtain a lattice parameterized by $\mathbf{s}'_2$. We call this lattice a cluster corresponding to given $\tilde{\mathbf{s}}, \mathbf{s}$. Each cluster, obviously, is a square lattice with the step $\epsilon_q q$. It contains no more than $(9k^{r_1}q^{-1})^2$ elements, since $|(\mathbf{s}'_2)_j| \leq 4k^{r_1}q^{-1} + 1$, $j = 1, 2$. The size of each cluster is less than $5|\epsilon_q|k^{r_1}$. If ϵ_q satisfies slightly stronger inequality, than (142) than clusters don't overlap, see the following lemma.

Lemma 4.5. *Suppose that ϵ_q satisfies the inequality*

$$|\epsilon_q| \leq \frac{1}{64}q^{-1}k^{-r_1}. \quad (144)$$

Then, the size of each cluster is less than $\frac{1}{8q}$. The distance between clusters is greater than $\frac{1}{2q}$.

Proof. Let us estimate the distance between points $\mathbf{s}'_2 = \mathbf{0}$ of two different clusters. Indeed, $\mathbf{s} - \frac{p}{q}\mathbf{s}''_2 \neq \mathbf{0}$, since $p\mathbf{s}''_2$, see (143), is not a multiple of q . Therefore, $\left| \left(\mathbf{s} - \frac{p}{q}\mathbf{s}''_2 \right)_j \right| \geq \frac{1}{q}$, $j = 1, 2$. Considering that $0 \leq (\mathbf{s}''_2)_j < q$, $j = 1, 2$, we obtain that the distance between two points where $\mathbf{s}'_2 = \mathbf{0}$ is greater than $\frac{1}{q} - |\epsilon_q|q$, that is greater than $\frac{15}{16q}$. The size of each cluster is obviously less than $|\epsilon_q|q(4k^{r_1}q^{-1} + 1) \leq \frac{1}{8q}$. Thus, two clusters cannot overlap, the distance between them being greater than $\frac{1}{2q}$. $\square$

We need two more properties of the lattice $\vec{p}_{\mathbf{m}}$, $\|\vec{p}_{\mathbf{m}}\| < 2k^{r_1}$.

Lemma 4.6. *The number of vectors $\vec{p}_{\mathbf{m}}$, satisfying the inequalities $\|\vec{p}_{\mathbf{m}}\| < 2k^{r_1}$, $p_{\mathbf{m}} < |\epsilon_q|qk^{r_1/3}$, does not exceed $k^{2r_1/3}$.*

Proof. Suppose vectors $\vec{p}_{\mathbf{m}}$ and $\vec{p}_{\mathbf{m}'}$ satisfy the conditions of the lemma. Then, $\|\vec{p}_{\mathbf{m}} - \vec{p}_{\mathbf{m}'}\| < 4k^{r_1}$. By definition of ϵ_q , $(2\pi)^{-1}|\vec{p}_{\mathbf{m}} - \vec{p}_{\mathbf{m}'}| \geq |\epsilon_q|q$. Thus, the distance between the points $\vec{p}_{\mathbf{m}}, \vec{p}_{\mathbf{m}'}$ is greater than $2\pi|\epsilon_q|q$ and each point can be surrounded by the disc of the radius $\pi|\epsilon_q|q$, the discs being disjoint. Dividing the area of the disc of the radius $2|\epsilon_q|qk^{r_1/3}$ (we increased radius to take into account points $\vec{p}_{\mathbf{m}}$ near the boundary of the disc $p_{\mathbf{m}} < |\epsilon_q|qk^{r_1/3}$) by the area of a disc of the radius $\pi|\epsilon_q|q$, we obtain that the number of vectors satisfying the inequality $p_{\mathbf{m}} < |\epsilon_q|qk^{r_1/3}$ does not exceed $k^{2r_1/3}$. $\square$

Lemma 4.7. *Suppose q in the inequality (141) satisfies the estimate $q > k^{2r_1/3}$. Then, the number of vectors $\vec{p}_{\mathbf{m}}$, $\|\vec{p}_{\mathbf{m}}\| < 2k^{r_1}$, satisfying the inequality $p_{\mathbf{m}} < k^{-2r_1/3}$ does not exceed $2^{12} \cdot k^{2r_1/3}$.*

Proof. First assume $|\epsilon_q| > \frac{1}{64}q^{-1}k^{-r_1}$. Then, dividing the area of the disc of the radius $2k^{-2r_1/3}$ by the area of a disc of the radius $\pi|\epsilon_q|q > \frac{1}{32}k^{-r_1}$, we obtain that the number of vectors satisfying the inequality $p_{\mathbf{m}} < k^{-2r_1/3}$ does not exceed $2^{12}k^{2r_1/3}$.

Second, we consider the case $|\epsilon_q| \leq \frac{1}{64}q^{-1}k^{-r_1}$. According to Lemma 4.5, the clusters do not overlap. The distance between clusters is greater than $\frac{1}{2q}$. Therefore, dividing the area of a disc with radius $\frac{3}{2}k^{-2r_1/3}$ by the area of a disc with radius $\frac{1}{4q}$, the last number being smaller than $\frac{1}{4}k^{-2r_1/3}$ by the conjecture of the lemma, we obtain that the number of clusters intersecting the disc of the radius $k^{-2r_1/3}$ is less than $(6k^{-2r_1/3}q)^2$. Each cluster contains less than $(9k^{r_1}q^{-1})^2$ points. Therefore, the total number of vectors $\vec{p}_{\mathbf{m}}$, $\|\vec{p}_{\mathbf{m}}\| < 2k^{r_1}$, satisfying the inequality $p_{\mathbf{m}} < k^{-2r_1/3}$ does not exceed $(6k^{-2r_1/3}q)^2 \cdot (9k^{r_1}q^{-1})^2 < 2^{12} \cdot k^{2r_1/3}$. $\square$

4.3.2 Lattice Points in the Nonresonant Set

Lemma 4.8. *Let $N(k, r_1, \vec{\mathbf{z}}_0, \varepsilon_0)$ be the number of points $\vec{\mathbf{z}}_0 + \vec{p}_{\mathbf{n}}$, $\|\vec{p}_{\mathbf{n}}\| < k^{r_1}$ in the ε_0 -neighborhood of $\mathcal{D}_1(k^{2l})$, where $\varepsilon_0 = k^{-5\mu r_1}$ and $\vec{\mathbf{z}}_0 \in \mathbb{R}^2$ being fixed. Then,*

$$N(k, r_1, \vec{\mathbf{z}}_0, \varepsilon_0) < 1000 \cdot k^{\frac{2r_1}{3}+1}.$$

Proof. Let us consider the segment $\vec{p}_{\mathbf{n}-\mathbf{n}'}$ between two points $\vec{\mathbf{z}}_0 + \vec{p}_{\mathbf{n}}$ and $\vec{\mathbf{z}}_0 + \vec{p}_{\mathbf{n}'}$ in the neighborhood. Obviously, $\|\vec{p}_{\mathbf{n}-\mathbf{n}'}\| < 2k^{r_1}$ and $p_{\mathbf{n}-\mathbf{n}'} > k^{-\mu r_1} \gg \varepsilon_0$. This means that the direction of the segment cannot be orthogonal to the curve (in fact they are almost parallel to the curve) and each end can be assigned its own angle coordinate $\varphi_{\mathbf{n}}, \varphi_{\mathbf{n}'}$, $\varphi_{\mathbf{n}} \neq \varphi_{\mathbf{n}'}$. We enumerate the points $\vec{\mathbf{z}}_0 + \vec{p}_{\mathbf{n}}$ in the order of increasing $\varphi_{\mathbf{n}}$ and connect neighboring points by segments. First we consider the segments with the length greater or equal to $\frac{1}{64}k^{-\frac{2r_1}{3}}$. Since the length of $\mathcal{D}_1(k^{2l})$ does not exceed $3\pi k$, the number of such segments does not exceed $650k^{\frac{2r_1}{3}+1}$.

It remains to estimate the number of segments with the length less than $\frac{1}{64}k^{-\frac{2r_1}{3}}$. First, we prove that no two segments $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}, \vec{p}_{\mathbf{n}_2-\mathbf{n}'_2}$ can be equal to each other. We use concavity of the curve $\mathcal{D}_1(k^{2l})$ and a small size ε_0 of its neighborhood. We show that for every $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$ with both ends in the neighborhood, there is a point on the curve where the tangent vector is parallel to $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$. Since the tangent vector changes monotonously with φ , no two vectors $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$ can have the same direction. Indeed, let us consider a segment $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$. Let (x, y) be local coordinates associated with $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$, the beginning of the segment being at the origin and the end having the coordinates $(\tau, 0)$, $\tau = p_{\mathbf{n}_1-\mathbf{n}'_1}$. The curve is described by the equation $y = y(x)$. It easily follows from Lemma 3.11 that $y'(x) = o(1)$ and the curvature κ of the curve $\mathcal{D}_1(k^{2l})$ is $\frac{1}{k}(1+o(1))$ at all points of the curve. Using the formula $\kappa(x) = |y''(x)| (1 + y'(x)^2)^{-3/2}$, we easily obtain $y''(x) = -\frac{1}{k}(1+o(1))$. Using a Taylor formula, we get $y(\tau) = y(0) + y'(0)\tau - \frac{1}{2k}(1+o(1))\tau^2$. Note that $|y(0)|, |y(\tau)| < 2\varepsilon_0$, since both ends of the segment are in the ε_0 -neighborhood of the curve. Considering also that $\tau > k^{-r_1\mu}$ and the estimate on ε_0 , we conclude: $\frac{\tau}{k} = 2y'(0)(1+o(1)) + O(k^{-4r_1\mu})$. Substituting this into the Taylor formula

$$y'(\tau) = y'(0) - \frac{\tau}{k}(1+o(1)), \quad (145)$$

we obtain: $y'(\tau) = -y'(0)(1+o(1)) + O(k^{-4r_1\mu})$. If $y'(\tau)$ and $y'(0)$ have the same sign or one of them is zero, the last relation yields $|y'(\tau)| + |y'(0)| = O(k^{-4r_1\mu})$. This contradicts to (145), since $\tau > k^{-r_1\mu}$. Therefore, $y'(\tau)$ and $y'(0)$ have different signs. Considering that $y'(x)$ is continuous, we obtain that there is a point x_0 in $(0, \tau)$ such that $y'(x_0) = 0$. This means that the isoenergetic curve at this point is parallel to $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$.

To finish the proof of the lemma we consider two cases. Suppose q in the inequality (141) satisfies the estimate $q > k^{2r_1/3}$. Then, by Lemma 4.7, the number of vectors $\vec{p}_{\mathbf{n}}$, $\|\vec{p}_{\mathbf{n}}\| < 2k^{r_1}$, satisfying the inequality $p_{\mathbf{n}} < \frac{1}{64}k^{-2r_1/3}$ does not exceed $2^{12} \cdot k^{2r_1/3}$. Since

each of them can be used only once, the total number of short segments does not exceed $2^{12} \cdot k^{2r_1/3}$.

Let $q \leq k^{2r_1/3}$. If $|\epsilon_q| > \frac{1}{64}q^{-1}k^{-r_1}$. Then, obviously, $\frac{1}{64}k^{-2r_1/3} < |\epsilon_q|qk^{r_1/3}$. Applying Lemma 4.6, we obtain that the number of segments with the length less than $\frac{1}{64}k^{-2r_1/3}$ is less than $k^{2r_1/3}$. Since each of them can be used only once, the total number of short segments does not exceed $k^{2r_1/3}$. It remains to consider the case $q \leq k^{2r_1/3}$, $|\epsilon_q| \leq \frac{1}{64}q^{-1}k^{-r_1}$. By Lemma 4.5, clusters are well separated. Considering that the distance between clusters is greater than $\frac{1}{2q}$ and the size of each cluster is less than $\frac{1}{8q}$, we obtain that no more than $8\pi qk$ clusters can intersect ε_0 -neighborhood of $\mathcal{D}_1(k^{2l})$. The part of the curve inside the clusters has the length L_{in} which is less than the double size of a cluster $10|\epsilon_q|k^{r_1}$ (the curve is concave) multiplied by the number of clusters $8\pi qk$, i.e., $L_{in} < 80\pi|\epsilon_q|qk^{r_1+1}$. Next, the segments with the length less than $\frac{1}{2}k^{-2r_1/3}$ cannot connect different clusters, since the distance between clusters is greater than $\frac{1}{2q} \geq \frac{1}{2}k^{-2r_1/3}$. Therefore, any segment of the length less than $\frac{1}{2}k^{-2r_1/3}$ is inside one cluster. If we consider the segments with the length greater than $|\epsilon_q|qk^{r_1/3}$, then the number of such segments is less than $L_{in}/|\epsilon_q|qk^{r_1/3}$, i.e., it is less than $80\pi k^{2r_1/3+1}$. By Lemma 4.6, the total number of segments of the length less than $|\epsilon_q|qk^{r_1/3}$ is less than $k^{2r_1/3}$. Each of them can be used only once. Thus, the total number of segments is less than $300k^{2r_1/3+1}$. $\square$

4.3.3 Lattice Points in the Resonant Set

Let $\mathcal{Q}_0 = \{\mathbf{0}\} \cup \mathcal{Q}$ where $\mathcal{Q} = \{\mathbf{q}_1, \dots, \mathbf{q}_J\}$, $J \geq 1$ and $\min_{\mathbf{q}, \mathbf{q}' \in \mathcal{Q}_0} \|\vec{p}_{\mathbf{q}-\mathbf{q}'}\| < k^\delta$. We assume that all elements of $\mathcal{Q}_0$ are different. We say that $\vec{\mathcal{Z}} \in \mathcal{R}_{\mathcal{Q}_0} \subset \mathbb{R}^2$ if all the following inequalities hold:

$$\begin{aligned} \left| \|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{q}}\|_{\mathbb{R}}^2 - k^2 \right| &< k^{-40\mu\delta} \quad \text{when } \mathbf{q} \in \mathcal{Q}_0, \\ \left| \|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{q}'}\|_{\mathbb{R}}^2 - k^2 \right| &\geq k^{-40\mu\delta} \quad \text{when } \mathbf{q}' \notin \mathcal{Q}_0 \text{ and } \min_{\mathbf{q} \in \mathcal{Q}_0} \|\vec{p}_{\mathbf{q}'-\mathbf{q}}\| < k^\delta. \end{aligned} \quad (146)$$

By Lemma 3.13 such $\vec{\mathcal{Z}}$ may exist only if $J \leq 3$. Let $P_{\mathcal{Q}_0}$ be a diagonal projection: $(P_{\mathcal{Q}_0})_{\mathbf{nn}} = 1$ if and only if $\min_{\mathbf{q} \in \mathcal{Q}_0} \|\vec{p}_{\mathbf{n}-\mathbf{q}}\| < k^\delta$. The dimension of $P_{\mathcal{Q}_0}$ clearly does not exceed $(2(J+1)k^\delta)^4$. Suppose $\vec{\mathcal{Z}} \in \mathcal{R}_{\mathcal{Q}_0}$. We consider operator $P_{\mathcal{Q}_0} (H(\vec{\mathcal{Z}}) - (k^{2l} + \varepsilon_0)I) P_{\mathcal{Q}_0}$, $|\varepsilon_0| < 1$, and its determinant $D(\vec{\mathcal{Z}}, k^{2l} + \varepsilon_0)$. Let $\mathcal{S}_{\mathcal{Q}_0}(k, \varepsilon_0) \subset \mathcal{R}_{\mathcal{Q}_0}$ be the set:

$$\mathcal{S}_{\mathcal{Q}_0}(k, \varepsilon_0) = \left\{ \vec{\mathcal{Z}} \in \mathcal{R}_{\mathcal{Q}_0} : D(\vec{\mathcal{Z}}, k^{2l} + \varepsilon'_0) = 0 \text{ for some } |\varepsilon'_0| < \varepsilon_0 \right\}. \quad (147)$$

Obviously, $\mathcal{S}_{\mathcal{Q}_0}(k, \varepsilon_0) = \mathcal{S}(k, \varepsilon_0) \cap \mathcal{R}_{\mathcal{Q}_0}$, see (140). Let (τ_1, τ_2) be new orthogonal coordinates with the origin at the point $\frac{1}{2}\vec{p}_{\mathbf{q}_1}$, τ_1 -axis being in the direction of $\vec{p}_{\mathbf{q}_1}$. It is easy to see that $\mathcal{R}_{\mathcal{Q}_0} \subset \{(\tau_1, \tau_2) : |\tau_1| < k^{-39\mu\delta}, \frac{1}{2}k < |\tau_2| < 2k\}$.

Lemma 4.9. *The set $D(\vec{\mathcal{Z}}, k^{2l}) = 0$ has the following properties in $\mathcal{R}_{\mathcal{Q}_0}$:*

1. *The equation $D(\vec{\mathcal{Z}}, k^{2l}) = 0$ describes at most 8 curves. They are described by the equations $\tau_2 = f_i(\tau_1)$, where $|f'_i(\tau_1)| < ck^{-1+\delta}$, $0 \leq i \leq \tilde{J}$, $\tilde{J} \leq 8$.*

2. The set $\mathcal{S}_{\mathcal{Q}_0}(k, \varepsilon_0)$ belongs to $\cup_i^{\tilde{J}} \mathcal{S}_i(k, \varepsilon_0)$, $\mathcal{S}_i(k, \varepsilon_0) = \{\vec{\mathcal{X}} : |\tau_2 - f_i(\tau_1)| < 2\varepsilon_0, |\tau_1| < k^{-39\mu\delta}\}$.
3. The curves $\tau_2 = f_i(\tau_1)$, $0 \leq i \leq \tilde{J}$, all together have no more than $2^{31}l^2 \cdot k^{8\delta}$ inflection points.
4. Let $\mathbf{l}$ is a segment of a straight line,

$$\mathbf{l} = \{\vec{\mathcal{X}} = (\tau_1, \beta_1\tau_1 + \beta_2), \tau_{1,0} < \tau_1 < \tau_{1,0} + \eta\}, \quad |\tau_{1,0}| < k^{-39\mu\delta}, \quad (148)$$

such that both its ends belong to $\mathcal{S}_i(k, \varepsilon_0)$, $2\varepsilon_0 < \eta^8 k^{-8l}$, $0 < \eta < 1$. Then, there is an inner part $\mathbf{l}'$ of the segment which is not in $\mathcal{S}_i(k, \varepsilon_0)$. Moreover, there is a point (τ_{1*}, τ_{2*}) in $\mathbf{l}'$ such that $f'_i(\tau_{1*}) = \beta_1$, i.e., the curve and the segment have the same direction when $\tau_1 = \tau_{1*}$.

Proof.

1. Let us consider eigenvalues $\hat{\lambda}_i(\vec{\mathcal{X}})$ of $P_{\mathcal{Q}_0}H(\vec{\mathcal{X}})P_{\mathcal{Q}_0}$. Obviously, there are no more than $J+1$ eigenvalues satisfying the inequality $|\hat{\lambda}_i(\vec{\mathcal{X}}) - k^{2l}| < \frac{1}{2}k^{2l-2-40\mu\delta}$ when $\vec{\mathcal{X}} \in \mathcal{R}_{\mathcal{Q}_0}$. Let E_0 be the diagonal projection: $(E_0)_{\mathbf{m}'\mathbf{m}'} = 1$ if and only if $\mathbf{m}' \in \mathcal{Q}_0$. Let

$$T = P_{\mathcal{Q}_0} \frac{\partial H(\vec{\mathcal{X}})}{\partial \tau_2} = P_{\mathcal{Q}_0} \frac{\partial H_0(\vec{\mathcal{X}})}{\partial \tau_2}.$$

Considering that other $\vec{p}_{\mathbf{q}_i}$, if any, have directions close to $\vec{p}_{\mathbf{q}_1}$, and T is a diagonal operator, we obtain:

$$TE_0 = 2l\tau_2 k^{2l-2} E_0(I + o(1)), \quad |\tau_2| \approx k, \quad (149)$$

where $o(\cdot)$ stands in the sense of the norm of bounded operators. Let $\hat{e}_i$ be a normalized eigenvector corresponding to $\hat{\lambda}_i$: $|\hat{\lambda}_i(\vec{\mathcal{X}}) - k^{2l}| < \frac{1}{2}k^{2l-2-40\mu\delta}$. By (146) and regular perturbation formulas, $\hat{e}_i = E_0\hat{e}_i + o(1)$. Hence,

$$\frac{\partial \hat{\lambda}_i}{\partial \tau_2} = (T\hat{e}_i, \hat{e}_i) = 2l\tau_2 k^{2l-2}(1 + o(1)). \quad (150)$$

By simple perturbation arguments, $\frac{\partial \hat{\lambda}_i}{\partial \tau_1} = O(k^{2l-2+\delta})$. Hence, the number of curves satisfying the equations $\hat{\lambda}_i(\vec{\mathcal{X}}) = k^{2l}$ in $\mathcal{R}_{\mathcal{Q}_0}$ is at most $2(J+1)$, each corresponding to a particular i , $1 \leq i \leq J+1$, and a sign of τ_2 . They can be described as $\tau_2 = f_i(\tau_1)$, $f_i(\tau_1)$ being piecewise differentiable and $|f'_i(\tau_1)| < ck^{-1+\delta}$.

2. It easily follows from (147) and (150) that $\mathcal{S}_{\mathcal{Q}_0} \subset \cup_{i=1}^{\tilde{J}} \mathcal{S}_i$, $1 \leq \tilde{J} \leq 8$.

3. Inflection points of the curves $\tau_2 = f_i(\tau_1)$ are described by the system of equations:

$$D = 0, \quad (151)$$

$$D_{\tau_2\tau_2}(D_{\tau_1})^2 + 2D_{\tau_1\tau_2}D_{\tau_1}D_{\tau_2} + D_{\tau_1\tau_1}(D_{\tau_2})^2 = 0 \quad (152)$$

where $D = D(\vec{\mathcal{Z}}, k^{2l})$. The left hand sides of (151) and (152) are polynomials of the degree K and $3K - 4$, $K \leq 2l(2(J+1)k^\delta)^4 \leq 2^{13}lk^{4\delta}$, with respect to τ_1, τ_2 . If they are mutually irreducible than, by Bezout theorem, the number of inflection points does not exceed $K(3K - 4) < 2^{28}l^2 \cdot k^{8\delta}$. Suppose the left hand sides of (151) and (152) are mutually reducible. Then, there is a solution $\tau_2 = f_i(\tau_1)$ with the zero curvature everywhere, i.e. a straight line $\tau_2 = a\tau_1 + b$. Considering that $D(\tau_1, a\tau_1 + b) = (\tau_1^{2l} + a^{2l}\tau_1^{2l})^{K/(2l)}(1 + o(1))$ as $\tau_1 \rightarrow \infty$, we conclude that a straight line cannot satisfy the equation (151). Thus, the total number of inflection points for the curves $\tau_2 = f_i(\tau_1)$ all together does not exceed $2^{31}l^2 \cdot k^{8\delta}$.

4. Let us consider a segment (148) of a straight line, such that both its ends are in $\mathcal{S}_i(k, \varepsilon_0)$ and $2\varepsilon_0 < \eta^8 k^{-8l}$. It follows that $\beta_1 = O(k^{-1+\delta})$, $|\beta_2| \approx k$, since $f'_i(\tau_1) = O(k^{-1+\delta})$. Next, we show that there is a part $\mathbf{I}'$ of $\mathbf{I}$ which is outside of $\mathcal{S}_i(k, \varepsilon_0)$. Note that $D(\vec{\mathcal{Z}}, k^{2l} + \varepsilon'_0) = 0$ if and only if $\hat{D}(\vec{\mathcal{Z}}, k^{2l} + \varepsilon'_0) = 0$ where $\hat{D}(\vec{\mathcal{Z}}, \lambda)$ is the determinant of the matrix

$$\hat{H}^{-1}P_{\mathcal{Q}_0}(H - \lambda I)P_{\mathcal{Q}_0}, \quad \hat{H} = (H_0 - \lambda I)(I - E_0) + E_0. \quad (153)$$

Note that diagonal terms of the matrix $\hat{H}^{-1}P_{\mathcal{Q}_0}(H - \lambda I)P_{\mathcal{Q}_0}$ are equal to 1 unless they correspond to E_0 and

$$\|\hat{H}^{-1}(I - E_0)\| < ck^{-(2l-2-40\mu\delta)} \quad \text{when } \vec{\mathcal{Z}} \in \mathcal{R}_{\mathcal{Q}_0}. \quad (154)$$

Let us extend $\hat{D}(\tau_1, \beta_1\tau_1 + \beta_2)$ as an analytic function of τ_1 into the complex disc $\mathbf{D} = \{\tau_1 \in \mathbb{C} : |\tau_1| < k^{-39\mu\delta}\}$. We also consider the following regions in $\mathbb{C}$:

$$\mathbf{D}_{\mathbf{q}} = \{\tau_1 \in \mathbb{C} : \|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{q}}|_{\mathbb{R}}^{2l} - k^{2l}\| < k^\delta, \quad \vec{\mathcal{Z}} = (\tau_1, \beta_1\tau_1 + \beta_2)\}, \quad \mathbf{q} \in \mathcal{Q}_0.$$

By $\hat{\mathbf{D}}_{\mathbf{q}}$ we will denote the set of vectors $\vec{\mathcal{Z}}$ corresponding to $\mathbf{D}_{\mathbf{q}}$. We are interested only in the connected component(s) of $\cup_{\mathbf{q} \in \mathcal{Q}_0} \hat{\mathbf{D}}_{\mathbf{q}}$ having the non-empty intersection with $\mathcal{R}_{\mathcal{Q}_0}$. It is easy to show that $\mathbf{I} \cap \mathcal{S}_{\mathcal{Q}_0} \subset \cup_{\mathbf{q} \in \mathcal{Q}_0} \hat{\mathbf{D}}_{\mathbf{q}}$. Therefore, we assume that $\mathbf{I} \subset \cup_{\mathbf{q} \in \mathcal{Q}_0} \hat{\mathbf{D}}_{\mathbf{q}}$ (otherwise the lemma is proved). Note that the estimate (154) is preserved for such regions, since each $\mathbf{D}_{\mathbf{q}}$ can be included in the balls of the radius $ck^{-l+1+\delta/2}$ centered at the points $\|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{q}}|_{\mathbb{R}}^{2l} - k^{2l} = 0$ and for any pair $\mathbf{q}, \mathbf{q}'$ $\frac{d}{d\tau_1} (\|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{q}}|_{\mathbb{R}}^2 - \|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{q}'}|_{\mathbb{R}}^2) = O(|\vec{p}_{\mathbf{q}} - \vec{p}_{\mathbf{q}'}|)$. Let

$$\hat{D}_0(\tau_1, \beta_1\tau_1 + \beta_2) = \prod_{\mathbf{q} \in \mathcal{Q}_0} \|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{q}}|_{\mathbb{R}}^{2l} - k^{2l}\|, \quad \vec{\mathcal{Z}} = (\tau_1, \beta_1\tau_1 + \beta_2).$$

Obviously $\hat{D}_0$ has at most $2(J+1)$ roots inside $\cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$. We denote the number of roots by $\hat{J}$, $\hat{J} \leq 2(J+1) \leq 8$. It is easy to see that $|\hat{D}_0| > k^{\delta(J+1)}$ on the boundary of $\cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$. Using (154) and (89), it is easy to show that $\hat{D} = \hat{D}_0 + O(k^{\delta J})$ on the boundary. Applying Rouchet's theorem, we see that $\hat{D}$ has $\hat{J}$ roots inside the union of the disks and satisfies the estimate $|\hat{D}| > \frac{1}{2}k^{(J+1)\delta}$ on the boundary $\cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$. Therefore, $\hat{D}$ can be represented in the form:

$$\hat{D}(\tau_1, \beta_1 \tau_1 + \beta_2) = \tilde{f}(\tau_1) \prod_{i=1}^{\hat{J}} (\tau_1 - \tau_1^{(i)}), \quad \tau_1 \in \cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}, \quad 0 \leq \hat{J} \leq 8. \quad (155)$$

Note that each $\mathbf{D}_{\mathbf{q}}$ can be included in the balls of the radius $ck^{-l+1+\delta/2}$ centered at the points $|\vec{z} + \vec{p}_{\mathbf{q}}|_{\mathbb{R}}^{2l} - k^{2l} = 0$. From the minimum principle ($\tilde{f} \neq 0$) it follows:

$$|\tilde{f}| \geq ck^{\gamma}, \quad \gamma = \delta(J+1) + (l-1-\delta/2)\hat{J} > 0, \quad (156)$$

when $\tau_1 \in \cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$. It easily follows that (155), (156) hold in $\mathcal{R}_{\mathcal{Q}_0}$. Let us consider a segment of the straight line $\vec{z} = [\tau_1, \beta_1 \tau_1 + \beta_2]$, $\tau_1 \in (\tau_{1,0}, \tau_{1,0} + \eta)$, $0 < \eta < 1$. By (155), there is a point $(\tau'_1, \beta_1 \tau'_1 + \beta_2)$ in this segment, where $|\hat{D}| > ck^{\gamma} \eta^{\hat{J}}$. Considering that $\hat{D}(\tau'_1, f_i(\tau'_1)) = 0$ by the definition of the curve and the obvious inequality $|\hat{D}_{\tau_2}(\tau_1, \tau_2)| < k^{2l(J+1)-1}$, we obtain $|f_i(\tau'_1) - (\beta_1 \tau'_1 + \beta_2)| > ck^{\gamma} \eta^{\hat{J}} / ck^{2l(J+1)-1} > k^{-2l(J+1)} \eta^{\hat{J}}$. If $\eta^8 k^{-8l} > 2\varepsilon_0$, then there are points in the segment which are outside $\mathcal{S}_i(k, \varepsilon_0)$. At one of these points the function $|f_i(\tau'_1) - (\beta_1 \tau'_1 + \beta_2)|$ attains its maximum value. At this point the curve and the line are parallel.

Remark. Note that the perturbation series for $(P_{\mathcal{Q}_0}(H - k^{2l}I)P_{\mathcal{Q}_0})^{-1}$ converges (with respect to $(P_{\mathcal{Q}_0}(H_0 - k^{2l}I)P_{\mathcal{Q}_0})^{-1}$) when τ_1 is on the boundary of $\cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$ and

$$\left\| (P_{\mathcal{Q}_0}(H - k^{2l}I)P_{\mathcal{Q}_0})^{-1} \right\| < ck^{\delta}. \quad (157)$$

By (155), the resolvent has no more than $\hat{J}$ poles inside $\cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$. Considering that each $\mathbf{D}_{\mathbf{q}}$ can be included in a ball of the radius $ck^{-l+1+\delta/2}$ and applying the maximum principle for the norm of a holomorphic operator, we obtain the following estimate inside $\cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$:

$$\left\| (P_{\mathcal{Q}_0}(H - k^{2l}I)P_{\mathcal{Q}_0})^{-1} \right\| < ck^{\delta} \left(\frac{ck^{-l+1+\delta/2}}{d} \right)^{\hat{J}}, \quad (158)$$

where d is the distance from a point $\tau_1 \in \cup_{\mathbf{q} \in \mathcal{Q}_0} \mathbf{D}_{\mathbf{q}}$ to a nearest pole of the resolvent. If $d \geq \eta/16$, then

$$\left\| (P_{\mathcal{Q}_0}(H - k^{2l}I)P_{\mathcal{Q}_0})^{-1} \right\| < ck^{\delta} \left(\frac{ck^{-l+1+\delta/2}}{\eta} \right)^{\hat{J}} < \varepsilon_0^{-1}. \quad (159)$$

It also shows that there is a point in $\mathbf{l}$ which is not in $\mathcal{S}(k, \varepsilon_0)$.

□

Let $N_{Q_0}(k, r_1, \vec{z}_0, \varepsilon_0)$ be the number of points $\vec{z}_0 + \vec{p}_n$, $\|\vec{p}_n\| < k^{r_1}$ in $S_{Q_0}(k, \varepsilon_0)$, $\vec{z}_0$ being fixed.

Lemma 4.10. *Let $\delta < r_1 < \infty$. If $\varepsilon_0 < k^{-16\mu r_1}$ then the number of points $N_{Q_0}(k, r_1, \vec{z}_0, \varepsilon_0)$ admits the estimate*

$$N_{Q_0}(k, r_1, \vec{z}_0, \varepsilon_0) \leq 2^{44} l^2 \cdot k^{2r_1/3+8\delta}. \quad (160)$$

Proof. The proof of the lemma is analogous to that of Lemma 4.8, when we replace properties of a distorted circle $\mathcal{D}_1(\lambda)$ by analogous properties of the set $D(\vec{z}) = 0$ proven in the previous lemma. Indeed, let us consider the segment $\vec{p}_{n-n'}$ between two points $\vec{z}_0 + \vec{p}_n$ and $\vec{z}_0 + \vec{p}_{n'}$ in the ε_0 -neighborhood of a concave component of a curve $\tau_2 = f_i(\tau_1)$. Obviously, $\|\vec{p}_{n-n'}\| < 2k^{r_1}$ and $p_{n-n'} > k^{-\mu r_1} \gg \varepsilon_0$. This means that the direction of the segment cannot be orthogonal to the curve and each end can be assigned its own coordinate $\tau_{1n}, \tau_{1n'}, \tau_{1n} \neq \tau_{1n'}$. We enumerate the points $\vec{z}_0 + \vec{p}_n$ in the order of increasing τ_{1n} and connect neighboring points by segments. Consider the segments with the length greater or equal to $\frac{1}{64}k^{-\frac{2r_1}{3}}$. Obviously, the length of the curve does not exceed 1. Hence, the number of such segments does not exceed $128k^{\frac{2r_1}{3}}$. It remains to estimate the number of segments with the length less than $\frac{1}{64}k^{-\frac{2r_1}{3}}$.

First, we prove that no two segments $\vec{p}_{n_1-n'_1}, \vec{p}_{n_2-n'_2}$ can be equal to each other in the same concave component of a curve $\tau_2 = f_i(\tau_1)$. Indeed, both end of $\vec{p}_{n_1-n'_1}$ are in $S_{Q_0}(k, \varepsilon_0)$. By Lemma 4.9, part 4, there is a point on the curve between two ends of the segment, where the curve is parallel to a segment (we notice that now we use the lemma for $\eta > k^{-\mu r_1}$). The same is true for $\vec{p}_{n_2-n'_2}$. Since we consider a concave component of a curve, it cannot be true.

To finish the proof of the lemma we consider two cases. Suppose q in the inequality (141) satisfies the estimate $q > k^{2r_1/3}$. Then, by Lemma 4.7, the number of vectors $\vec{p}_n$, $\|\vec{p}_n\| < 2k^{r_1}$, satisfying the inequality $p_n < \frac{1}{64}k^{-2r_1/3}$ does not exceed $2^{12} \cdot k^{2r_1/3}$. Since each of them can be used only once, the total number of short segments does not exceed $2^{12} \cdot k^{2r_1/3}$.

Let $q \leq k^{2r_1/3}$. If $|\epsilon_q| > \frac{1}{64}q^{-1}k^{-r_1}$. Then, obviously, $\frac{1}{64}k^{-2r_1/3} < |\epsilon_q|qk^{r_1/3}$. Applying Lemma 4.6, we obtain that the number of segments with the length less than $\frac{1}{64}k^{-2r_1/3}$ is less than $k^{2r_1/3}$. Since each of them can be used only once, the total number of short segments does not exceed $k^{2r_1/3}$. It remains to consider the case $q \leq k^{2r_1/3}$, $|\epsilon_q| \leq \frac{1}{64}q^{-1}k^{-r_1}$. By Lemma 4.5, clusters are well separated. Considering that the distance between clusters is greater than $\frac{1}{2q}$ and the size of each cluster is less than $\frac{1}{8q}$, we obtain that no more than $4q$ clusters can intersect ε_0 -neighborhood of a concave component of a curve $\tau_2 = f_i(\tau_1)$. The part of the curve inside the clusters has the length L_{in} which is less than the double size of a cluster $10|\epsilon_q|k^{r_1}$ multiplied by the number of clusters $4q$, i.e., $L_{in} < 40|\epsilon_q|qk^{r_1}$. Next, the segments with the length less than $\frac{1}{2}k^{-2r_1/3}$ cannot connect different clusters, since the distance between clusters is greater than $\frac{1}{2q} \geq \frac{1}{2}k^{-2r_1/3}$.

Therefore, any segment of the length less than $\frac{1}{2}k^{-2r_1/3}$ is inside one cluster. If we consider the segments with the length greater than $|\epsilon_q|qk^{r_1/3}$, then the number of such segments is less than $L_{in}/|\epsilon_q|qk^{r_1/3}$, i.e., it is less than $40k^{2r_1/3}$. By Lemma 4.6, the total number of segments of the length less than $|\epsilon_q|qk^{r_1/3}$ is less than $k^{2r_1/3}$. Each of them can be used only once. Thus, the total number of segments is less than $41k^{2r_1/3}$.

We proved that the number of segments in ε_0 -neighborhood of each concave component of a curve $\tau_2 = f_i(\tau_1)$ does not exceed $2^{13}k^{2r_1/3}$. By Lemma 4.9, part 3, the number of such components does not exceed $2^{31}l^2 \cdot k^{8\delta}$. The estimate (160) easily follows. $\square$

4.4 Preparation for Step III - Analytic Part

4.4.1 Model Operator for Step III

Let $r_2 > r_1 > 10^8$. Further we use the notation:

$$\Omega(r_2) = \{\mathbf{m} : \|\vec{p}_{\mathbf{m}}\| < k^{r_2}\}. \quad (161)$$

We repeat for r_2 the construction of the section 3.5.1 which was done for an arbitrary $r_1 > 2$. It is easy to see that the whole construction is monotonous with respect to r_1 . Namely,

$$\mathcal{M}(\varphi_0, r_1) \subseteq \mathcal{M}(\varphi_0, r_2), \quad \mathcal{M}'(\varphi_0, r_1) \subseteq \mathcal{M}'(\varphi_0, r_2), \quad \mathcal{M}_1(\varphi_0, r_1) \subseteq \mathcal{M}_1(\varphi_0, r_2),$$

$$\tilde{\mathcal{M}}_2(\varphi_0, r_1) \subseteq \tilde{\mathcal{M}}_2(\varphi_0, r_2), \quad \tilde{\mathcal{M}}_1(\varphi_0, r_1) \subseteq \tilde{\mathcal{M}}_1(\varphi_0, r_2), \quad \tilde{\mathcal{M}}_2(\varphi_0, r_1) \subseteq \tilde{\mathcal{M}}_2(\varphi_0, r_2).$$

Let $\varphi_0 \in \omega^{(2)}(k, \delta, 1)$. Put

$$\mathcal{M}^{(2)} := \mathcal{M}^{(2)}(\varphi_0, r_2) = \{\mathbf{m} \in \mathcal{M}(\varphi_0, r_2) : \varphi_0 \in \mathcal{O}_{\mathbf{m}}^{(2)}(10r'_1, 1)\}, \quad (162)$$

where $\mathcal{O}_{\mathbf{m}}^{(2)}(10r'_1, \tau)$ is the union of the disks of the radius $\tau k^{-10r'_1}$ with the centers at poles of the resolvent of k^δ -component containing $\vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}}$. More precisely, for each $\mathbf{m} \in \mathcal{M}(\varphi_0, r_2)$ we construct k^δ -box around it. We establish $3k^\delta$ equivalence relation between such boxes. Such components separated by k^δ from each other we call k^δ -components. Then $\mathcal{O}_{\mathbf{m}}^{(2)}(10r'_1, \tau)$ is the union of the disks of the radius $\tau k^{-10r'_1}$ with the centers at poles of the operator $(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P_{\mathbf{m}})^{-1}$, where $P_{\mathbf{m}}$ is the projection onto a particular k^δ -component containing $\vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}}$. We notice (see the proof of Lemma 3.13 with $3k^\delta$ and k^{r_2} instead of k^δ and k^{r_1}) that each k^δ -component contains not more than 4 elements $\mathbf{m} \in \mathcal{M}(\varphi_0, r_2)$. For such $\mathbf{m}$ corresponding sets $\mathcal{O}_{\mathbf{m}}^{(2)}(10r'_1, \tau)$ are identical. By construction of the non-resonant set $\omega^{(2)}(k, \delta, 1)$, we have $\mathcal{M}^{(2)} \cap \Omega(r_1) = \emptyset$.

Further we use the property of the set $\mathcal{M}^{(2)}$ formulated in the next lemma.

Lemma 4.11. *Let $\mathbf{m}_0 \in \Omega(r_2)$, $1/20 < \gamma' < 20$ and $\Pi_{\mathbf{m}_0}$ be the $k^{\gamma'r_1}$ -neighborhood (in $\|\cdot\|$ -norm) of $\mathbf{m}_0$. Then, the set $\Pi_{\mathbf{m}_0}$ contains less than $ck^{2\gamma'r_1/3+1}$ elements of $\mathcal{M}^{(2)}$.*

Proof. If $\mathbf{m} \in \mathcal{M}^{(2)}$, then there is a $\varphi_* : |\varphi_0 - \varphi_*| < k^{-10r'_1}$ such that

$$\det \left(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_*)) - k^{2l}I)P_{\mathbf{m}} \right) = 0. \quad (163)$$

Therefore, for some $\varepsilon'_0 : |\varepsilon'_0| < \varepsilon_0$, $\varepsilon_0 := ck^{2l-1-10r'_1}$,

$$\det \left(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - (k^{2l} + \varepsilon'_0)I)P_{\mathbf{m}} \right) = 0. \quad (164)$$

Indeed, if (164) holds for no ε' , then $\left\| \left(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P_{\mathbf{m}} \right)^{-1} \right\| < ck^{-2l+1+10r'_1}$, since φ_0 is real and, hence, $H(\vec{\mathcal{Z}}^{(1)}(\varphi_0))$ is selfadjoint. Using Hilbert identity, we obtain that $\left(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_*)) - k^{2l}I)P_{\mathbf{m}} \right)^{-1}$ is bounded. This contradicts to (163). Hence, (164) holds for some $\varepsilon'_0 : |\varepsilon'_0| < \varepsilon_0$.

Suppose $\mathbf{m} \in \mathcal{M}_1(\varphi_0, r_2)$. Then, (164) means that $|\lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}}) - k^{2l}| < \varepsilon_0$. Introducing the notation $\vec{\mathcal{Z}}_0 = \vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}_0}$, we rewrite the last inequality in the form: $|\lambda^{(1)}(\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{m}-\mathbf{m}_0}) - k^{2l}| < \varepsilon_0$, where $\|\vec{p}_{\mathbf{m}-\mathbf{m}_0}\| < k^{\gamma' r_1}$. It follows that $\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{m}-\mathbf{m}_0}$ is in the real $c\varepsilon_0 k^{-2l+1}$ -neighborhood of $\mathcal{D}_1(k^{2l})$. Applying Lemma 4.8, we obtain that the number of such points does not exceed $ck^{2\gamma' r_1/3+1}$. Let $\mathbf{m} \in \mathcal{M}_2(\varphi_0, r_2)$. Namely, let $\mathbf{m}$ belongs to a component $\mathcal{M}_2^j(\varphi_0, r_2)$. Then, $|\vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2| < k^{-40\mu\delta}$ and (164) holds, $P_{\mathbf{m}}$ being the projection on $\tilde{\mathcal{M}}_2^j(\varphi_0, r_2)$. Using again the notation $\vec{\mathcal{Z}}_0 = \vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}_0}$, and the definition of $\mathcal{M}_2^j(\varphi_0, r_2)$, we obtain: $|\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{m}-\mathbf{m}_0} + \vec{p}_{\mathbf{q}}|^2 - k^2| < k^{-40\mu\delta}$ for all $\mathbf{q} \in \mathcal{Q}_0$, where $\mathcal{Q}_0 := \mathcal{M}_2^j(\varphi_0, r_2) - \mathbf{m}$. In terms of Section 4.3.3, (164) means $\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{m}-\mathbf{m}_0} \in \mathcal{S}_{\mathcal{Q}_0}(k, \varepsilon_0)$, see (147). Applying Lemma 4.10 and using (164), we obtain that the number of such points does not exceed $ck^{2\gamma' r_1/3+8\delta}$ for a fixed $\mathcal{Q}_0$. Considering that the total number of sets $\mathcal{Q}_0$ does not exceed $ck^{12\delta}$, we obtain that the number of points $\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{m}-\mathbf{m}_0} \in \cup_{\mathcal{Q}_0} \mathcal{S}_{\mathcal{Q}_0}(k, \varepsilon_0)$ does not exceed $ck^{2\gamma' r_1/3+20\delta}$. Adding the estimates for the total number of resonant and non-resonant sets, we prove it is strictly less than $C_0 k^{2\gamma' r_1/3+1}$.

□

Let us split k^{r_2} -box into $k^{\gamma r_1}$ -boxes as described below. In the whole construction below we will have $\gamma = \frac{1}{5}$, but in some cases we will refer to the similar estimates with other values of γ . That's why in what follows we prefer to use implicit notation. The procedure consists of several steps. On each step we introduce a new scale of a box. Further structure will acquire additional scales at each step of approximation procedure. This is why we call the procedure Multiscale Construction in the Space of Momenta.

1. *Simple region.* Let $\Omega_s^{(2)}(r_2)$ be the collection of $\mathbf{m} \in \Omega(r_2)$ with small values of $p_{\mathbf{m}}$, namely, $\Omega_s^{(2)}(r_2) = \{\mathbf{m} \in \Omega(r_2) : 0 < p_{\mathbf{m}} \leq k^{-5r'_1}\}$. It is easy to see that $\Omega_s^{(2)}(r_2) \subset \mathcal{M}(\varphi_0, r_2)$, since $p_{\mathbf{m}}$ is small, see (71), (69). Next, if $\mathbf{m} \in \Omega_s^{(2)}(r_2)$, then there are no other elements of $\mathcal{M}(\varphi_0, r_2)$ in the k^{δ} -box around $\mathbf{m}$. Indeed,

let $\vec{\mathcal{Z}} = \vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}}$. It is a small perturbation of $\vec{\mathcal{Z}}^{(1)}(\varphi_0)$, hence it satisfies $||\vec{\mathcal{Z}} + \vec{p}_{\mathbf{n}}|^2 - |\vec{\mathcal{Z}}|^2| > \frac{\tau}{2}k^{-40\mu\delta}(1 + o(1))$ when $0 < ||\vec{p}_{\mathbf{n}}|| < k^\delta$, see (60). This means $\mathbf{m} + \mathbf{n} \notin \mathcal{M}(\varphi_0, r_2)$. Further, if $\mathbf{m} \in \Omega_s^{(2)}(r_2)$, then there are no other elements of $\Omega_s^{(2)}(r_2)$ in the surrounding box of the size k^{r_1} , see (28). Last, $\mathbf{m}$ itself can belong or do not belong to $\mathcal{M}^{(2)}$, there are no other elements of $\mathcal{M}^{(2)}$ in the k^{r_1} -box around such $\mathbf{m}$. Indeed, $\vec{\mathcal{Z}}^{(1)}(\varphi_0)$ satisfies the conditions of Lemma 3.18. This means that the k^δ -cluster around each $\mathbf{q}$: $0 < ||\vec{p}_{\mathbf{q}}|| < k^{r_1}$ is non-resonant. Moreover, the k^δ -box around each $\mathbf{m} + \mathbf{q}$: $0 < ||\vec{p}_{\mathbf{q}}|| < k^{r_1}$ is non-resonant too, since $p_{\mathbf{m}}$ is sufficiently small. This means $\mathbf{m} + \mathbf{q} \notin \mathcal{M}^{(2)}$.

For each $\mathbf{m} \in \Omega_s^{(2)}(r_2)$ we consider its $k^{r_1/2}$ -neighborhood. The union of such boxes we call the simple region and denote it by $\Pi_s(r_2)$. The corresponding projection is P_s . Note, that the distance from the simple region to the nearest point of $\mathcal{M}^{(2)}$ is greater than $\frac{1}{2}k^{r_1}$.

2. *Black region.* Next, we split $\Omega(r_2) \setminus (\Omega(r_1) \cup \Pi_s)$ into boxes of the size $k^{\gamma r_1}$. All elements $\mathbf{m} \in \mathcal{M}^{(2)}$ there satisfy $p_{\mathbf{m}} > k^{-5r_1'}$. We call a box black, if together with its neighbors it contains more than $k^{\gamma r_1/2 + \delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$, $\delta_0 = \gamma/100$ (in particular $\delta_0 r_1 > 100$). Let us consider all "black" boxes together with their $k^{\gamma r_1/2 + \delta_0 r_1}$ -neighborhoods. We call this the black region. Note that the size of the neighborhoods involved is much smaller than the size of the neighborhoods $k^{r_1/2}$ for the simple region, since $\gamma + \delta_0 < \frac{1}{2}$. The estimates for the size of the black region will be proven in Lemma 4.12. We denote the black region by Π_b . The corresponding projector is P_b . Obviously the distance between black and simple regions is greater than $\frac{1}{2}k^{r_1}$.
3. *Grey region.* By a white box we mean a $k^{\gamma r_1}$ -box, which together with its neighbors contains no more than $k^{\gamma r_1/2 + \delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$. Every white box we split into "small" boxes of the size $k^{\gamma r_1/2 + 2\delta_0 r_1}$. We call a small box "grey", if together with its neighbors it contains more than $k^{\gamma r_1/6 - \delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$. The grey region is the union of all grey small boxes together with their $k^{\gamma r_1/2 + 2\delta_0 r_1}$ -neighborhoods. Note that the size of the neighborhoods involved is much smaller than the size of the neighborhoods the simple and black regions. The estimates for the size of the grey region will be proven in Lemma 4.13. The notation for this region is Π_g . The corresponding projector is P_g . The part of the grey region, which is outside the black region, we denote by Π_g' and the corresponding projection by P_g' . Obviously, the distance between grey and simple regions is greater than $\frac{1}{2}k^{r_1}$.
4. *White region.* By a white small box we mean a small box, which together with its neighbors has no more than $k^{\gamma r_1/6 - \delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$. In each small white box we consider $k^{\gamma r_1/6}$ -boxes around each point of $\mathcal{M}^{(2)}$. The union of such $k^{\gamma r_1/6}$ -boxes we call the white region and denote it by Π_w . The corresponding projection is P_w .

Note that the size of the neighborhoods involved is much smaller than the size of the neighborhoods the simple, black and grey regions. The estimates for the size of the white region will be proven in Lemma 4.14. The part of the white region which is outside the black and grey regions, we denote Π'_w and the corresponding projection by P'_w . Obviously, the distance between grey and simple regions is greater than $\frac{1}{2}k^{r_1}$.

5. *Non-resonant region.* We also consider $\frac{1}{3}k^\delta$ -neighborhoods of all points in the set $\mathcal{M}(r_2, \varphi_0) \setminus \left(\mathcal{M}(r_1, \varphi_0) \cup \mathcal{M}^{(2)} \cup \Omega_s^{(2)}(r_2) \right)$. The union of this neighborhoods we call the non-resonance region Π_{nr} . The corresponding projection is P_{nr} . The part of the non-resonant region which is outside $\Pi_s \cup \Pi_b \cup \Pi_g \cup \Pi_w$, we denote Π'_{nr} and the corresponding projection by P'_{nr} .

Let

$$P_r := P_s + P_b + P'_g + P'_w, \quad P^{(2)} := P_r + P'_{nr} + P(r_1)$$

index r standing for "resonant".

First, we establish $3k^{\gamma r_1 + \delta_0 r_1}$ -equivalence relation between black boxes. Then the set Π_b can be represented as the union of components (clusters) separated by distance no less than $k^{\gamma r_1 + \delta_0 r_1}$. We denote such a component by Π_b^j .

Lemma 4.12. 1. *Each Π_b^j contains no more than $ck^{\gamma r_1/2 - \delta_0 r_1 + 3}$ black boxes.*

2. *The size of Π_b^j in $||| \cdot |||$ norm is less than $ck^{3\gamma r_1/2 + 3}$.*

3. *Each Π_b^j contains no more than $ck^{\gamma r_1 + 3}$ elements of $\mathcal{M}^{(2)}$. Moreover, any box of $||| \cdot |||$ -size $ck^{3\gamma r_1/2 + 3}$ containing Π_b^j has no more than $ck^{\gamma r_1 + 3}$ elements of $\mathcal{M}^{(2)}$ inside.*

Proof. Let n_b be the number of black boxes in Π_b^j , L_b be the size of Π_b^j and N_b the number of elements of $\mathcal{M}^{(2)}$ in Π_b^j . Obviously, $L_b < n_b 3k^{\gamma r_1 + \delta_0 r_1}$ and $N_b > cn_b k^{\gamma r_1/2 + \delta_0 r_1}$. By Lemma 4.11, $N_b < cL_b^{2/3}k$. Solving the last three inequalities for n_b , we get $n_b < ck^{\gamma r_1/2 - \delta_0 r_1 + 3}$. It follows $L_b < ck^{3\gamma r_1/2 + 3}$. Next, we consider a box of the size $k^{3\gamma r_1/2 + 3}$, containing Π_b^j . Using again Lemma 4.11, we obtain that the number of elements of $\mathcal{M}^{(2)}$ in this box is less than $cL_b^{2/3}k$. Therefore, $N_b < ck^{\gamma r_1 + 3}$. $\square$

Second, we establish $3k^{\gamma r_1/2 + 2\delta_0 r_1}$ -equivalence relation between small grey boxes. Then the set Π_g can be represented as the union of components separated by distance no less than $k^{\gamma r_1/2 + 2\delta_0 r_1}$. We denote each such component as Π_g^j .

Lemma 4.13. 1. *Each Π_g^j contains no more than $ck^{\gamma r_1/3 + 2\delta_0 r_1}$ grey boxes.*

2. *The size of Π_g^j in $||| \cdot |||$ norm is less than $ck^{5\gamma r_1/6 + 4\delta_0 r_1}$.*

3. *Each Π_g^j contains no more than $k^{\gamma r_1/2 + \delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$.*

Proof. Let us consider a part of Π_g^j belonging to one "big" white box. Let n_g be the number of grey boxes in Π_g^j . Let L_g be the size of Π_g^j and N_g be the number of elements of $\mathcal{M}^{(2)}$ in Π_g^j . Obviously, $N_g > cn_g k^{\gamma r_1/6 - \delta_0 r_1}$. By definition of a big white box $N_g < k^{\gamma r_1/2 + \delta_0 r_1}$. Therefore, $n_g < ck^{\gamma r_1/3 + 2\delta_0 r_1}$. Clearly, $L_g < n_g 3k^{\gamma r_1/2 + 2\delta_0 r_1} < ck^{5\gamma r_1/6 + 4\delta_0 r_1}$. Since $\delta_0 < \gamma/24$, we obtain that the size of each grey component is much less than the size $k^{\gamma r_1}$ of a big box. The lemma is proven under condition that Π_g^j is inside one of white boxes. Suppose Π_g^j intersects more than one white box. Considering that the size of Π_g^j in each big white box is much less than the size of this box, we conclude that Π_g^j fits into neighboring boxes and satisfies the estimates proven above. $\square$

Third, we consider points of $\mathcal{M}^{(2)}$ in small white boxes. We establish $3k^{\gamma r_1/6}$ -equivalence relation between them. Considering $k^{\gamma r_1/6}$ -neighborhoods of the points in $\mathcal{M}^{(2)}$, we see that this neighborhoods form clusters Π_w^j of Π_w separated by the distance no less than $k^{\gamma r_1/6}$. The number of $\mathcal{M}^{(2)}$ points in a white cluster we denote by N_w^j .

Lemma 4.14. 1. The size of Π_w^j in $\|\cdot\|$ norm is less than $ck^{\gamma r_1/3 - \delta_0 r_1}$.

2. Each Π_w^j contains no more than $k^{\gamma r_1/6 - \delta_0 r_1}$ points of $\mathcal{M}^{(2)}$.

Proof. Let us consider points of $\mathcal{M}^{(2)}$ in a small white box. By the definition of the white small box, the number of such points does not exceed $k^{\gamma r_1/6 - \delta_0 r_1}$. We consider the $k^{\gamma r_1/6}$ -neighborhoods of these points. They can form clusters. The total contribution from all points of $\mathcal{M}^{(2)}$ in the small white box and its neighbors, obviously, does not exceed $3k^{\gamma r_1/3 - \delta_0 r_1}$, which is much less than the size of a small white box. Therefore, each Π_w^j can't spread outside of the small white box and its neighbors. This proves both statements of the lemma. $\square$

At last, we also establish k^δ -equivalence relation between all points in $\mathcal{M}(r_2, \varphi_0) \setminus (\mathcal{M}(r_1, \varphi_0) \cup \mathcal{M}^{(2)} \cup \Omega_s^{(2)}(r_2))$. Then the set Π_{nr} can be represented as the union of components (clusters) separated by distance no less than $\frac{1}{3}k^\delta$. We denote such a component by Π_{nr}^j .

Next, we slightly change definitions of the black, grey and white areas to adjust their boundary to the structure of clusters. Namely, if k^δ -cluster containing points of $\mathcal{M}(\varphi_0, r_2) \setminus \mathcal{M}^{(2)}$ has a distance less than k^δ to a white, grey or black area, then we include it into the area with the lightest color. This "addition" does not change formulation of Lemmas 4.12, 4.13, 4.14, since the size of a k^δ -cluster is much smaller than the sizes of Π_b^j , Π_g^j , Π_w^j . If a white cluster has a distance less than $k^{\gamma r_1/6}$ to the grey or black area, we include it into that with the lighter color. This "addition" also does not change formulation of Lemmas 4.12, 4.13, since the size of a white cluster is much smaller than the sizes of Π_b^j , Π_g^j . If a grey cluster has a distance less than $k^{\gamma r_1/2 + 2\delta_0 r_1}$ to the black area, we include it into one of these areas. This "addition" does not change formulation of Lemma 4.12, since the size of a grey cluster is much smaller than the size of any Π_b^j . The new structure has the following properties: if the intersection of the k^δ -neighborhood of a k^δ -cluster with white, grey or

black area is not empty, then this cluster is completely in this area. If the intersection of the $k^{\gamma r_1/6}$ -neighborhood of a white cluster with grey or black area is not empty, then this cluster is completely in this area. If the intersection of the $k^{\gamma r_1/2+2\delta_0 r_1}$ -neighborhood of a grey cluster with the black area is not empty, then this cluster is completely in this area. $\delta_0 < \gamma/24$, $\gamma < 1/3$. This means that each component of the white, grey, black and non-resonance region is much smaller in $||| \cdot |||$ -size than $\Omega(r_1)$. Moreover, there are no points of $\mathcal{M}^{(2)}$ inside $\Omega(r_1)$. If the $k^{\gamma r_1/6}$ -neighborhood of a white cluster intersect $\Omega(r_1)$, we reduce $\Omega(r_1)$ by this neighborhood. This insignificant reduction does not change Step II. We make a similar reduction of $\Omega(r_1)$ if it is intersected by neighborhoods of grey or black clusters. Sometimes it will be convenient to numerate the projections $P(r_1)$, P_b , P'_g , P'_w , P'_{nr} , P_s by indices 0,1,2,3,4,5 as $P_0, P_1, P_2, P_3, P_4, P_5$. The corresponding sets are Π_i . Note that each Π_i consists of the components Π_{ij} , $j = 1, \dots, J(i)$ as described in the construction of the sets Π_i . The distance between closest components Π_{ij} , $i = 1, 2, 3, 4, 5$ with the same first index is greater than $k^{\gamma r_1+\delta_0 r_1}$, $k^{\gamma r_1/2+2\delta_0 r_1}$, $k^{\gamma r_1/6}$, $\frac{1}{3}k^\delta$, $k^{\gamma r_1}$, correspondingly. Then we can rewrite

$$P^{(2)} = \sum_{i=0}^5 P_i. \quad (165)$$

We introduce the boundaries $\partial\Omega(r_1)$, $\partial\Pi_b$, $\partial\Pi'_g$, $\partial\Pi'_w$, $\partial\Pi'_{nr}$, $\partial\Pi_s$ of the sets $\Omega(r_1)$, Π_b , Π'_g , Π'_w , Π'_{nr} , Π_s as follows: $\partial\Omega(r_1)$, $\partial\Pi_b$, $\partial\Pi'_g$, $\partial\Pi'_w$, $\partial\Pi'_{nr}$, $\partial\Pi_s$ are the sets of points in $\Omega(r_1)$, Π_b , Π'_g , Π'_w , Π'_{nr} , Π_s whose $||| \cdot |||$ -distance to the complements of $\Omega(r_1)$, Π_b , Π'_g , Π'_w , Π'_{nr} , Π_s , respectively, is less than $\frac{1}{3}k^\delta$. The corresponding projectors we denote as $P^\partial(r_1)$, P_b^∂ , P'^∂_g , P'^∂_w , P'^∂_{nr} , P_s^∂ or P_i^∂ , $i = 0, 1, 2, 3, 4, 5$.

Lemma 4.15. *Let $i, i' = 0, 1, 2, 3, 4, 5$, $i \neq i'$. The following relations hold:*

$$P_i P_{i'} = 0, \quad (166)$$

$$P_i V P_{i'} = 0, \quad (167)$$

$$(I - P^{(2)}) V P_i = (I - P^{(2)}) V P_i^\partial. \quad (168)$$

Corollary 4.16. *Operators $P^{(2)} V P^{(2)}$ and $P^{(2)} H P^{(2)}$ have a block structure. Namely,*

$$P^{(2)} V P^{(2)} = \sum_{i=0}^5 P_i V P_i, \quad P^{(2)} H P^{(2)} = \sum_{i=0}^5 P_i H P_i. \quad (169)$$

$$P_i V P_i = \sum_j P_{ij} V P_{ij}, \quad P_i H P_i = \sum_j P_{ij} H P_{ij}, \quad (170)$$

The lemma easily follows from the construction of the projectors and the fact that $V_{jj'} = 0$ when $|j - j'| \geq \frac{1}{3}k^\delta$.

Remark. Thus, we have constructed a multiscale structure inside $P^{(2)} H P^{(2)}$, blocks of different colors having distinctly different size. Merging blocks of a smaller size (a lighter color) with neighboring blocks of a bigger size (a darker color), we made the blocks to be separated by the $||| \cdot |||$ distance greater then k^δ .

Lemma 4.17. *Let $\varphi_0 \in \omega^{(2)}(k, \delta, \tau)$, $|\varphi - \varphi_0| < k^{-2l-40r'_1-\delta}$. Then,*

$$\left\| \left(P_{nr} (H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l} I) P_{nr} \right)^{-1} \right\| < ck^{40r'_1}. \quad (171)$$

Proof. The set Π_{nr} can be presented as $\cup_j \Pi_{nr}^j$, each Π_{nr}^j being a k^δ -cluster, and the distances between sets Π_{nr}^j is no less than $\frac{1}{3}k^\delta$. Therefore, $P_{nr} = \sum_j P_{nr}^j$, where P_{nr}^j are projections corresponding to Π_{nr}^j , and $P_{nr} H P_{nr} = \sum_j P_{nr}^j H P_{nr}^j$. Hence, it is enough to prove

$$\left\| \left(P_{nr}^j (H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l} I) P_{nr}^j \right)^{-1} \right\| < ck^{40r'_1}. \quad (172)$$

It follows from Lemma 3.14. Indeed, by construction, each Π_{nr}^j contains $\mathbf{m} \in \mathcal{M}(\varphi, r_2)$, $\mathcal{M}(\varphi, r_2) = \mathcal{M}_1(\varphi, r_2) \cup \mathcal{M}_2(\varphi, r_2)$. We can apply Lemma 3.14, since it is proven for any r_1 (no restrictions from above). We take $\varepsilon_0 = k^{-10r'_1}$ in Lemma 3.14, since the distance from φ_0 to the nearest pole of the operator $\left(P_{nr}^j (H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l} I) P_{nr}^j \right)^{-1}$ is greater than $k^{-10r'_1}$. By analogy with Corollary 3.16, we obtain (recall that now $p_{\mathbf{m}} > k^{-5r'_1}$)

$$\left\| \left(P_{nr}^j (H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l} I) P_{nr}^j \right)^{-1} \right\| < ck^{40r'_1}. \quad (173)$$

Taking into account that $\vec{\mathcal{Z}}^{(2)}(\varphi_0) - \vec{\mathcal{Z}}^{(1)}(\varphi_0) = o(k^{-2l-40\mu r'_1})$ and $\vec{\mathcal{Z}}^{(2)}(\varphi) - \vec{\mathcal{Z}}^{(2)}(\varphi_0) = o(k^{-2l+1-40r'_1})$, we arrive at (172). $\square$

Lemma 4.18. *Let $\varphi_0 \in \omega^{(2)}(k, \delta, \tau)$, and $|\varphi - \varphi_0| < k^{-44r'_1-2l-\delta}$, $i=1,2,3$. Then,*

1. *The number of poles of the resolvent $\left(P_i (H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l} I) P_i \right)^{-1}$ in the disc $|\varphi - \varphi_0| < k^{-44r'_1-2l-\delta}$ is no greater than $N_i^{(1)}$, where $N_1^{(1)} = k^{\gamma r_1+3}$, $N_2^{(1)} = k^{\gamma r_1/2+\delta_0 r_1}$, $N_3^{(1)} = k^{\gamma r_1/6-\delta_0 r_1}$.*
2. *Let ε be the distance to the nearest pole of the resolvent in $\mathcal{W}^{(2)}$ and let $\varepsilon_0 := \min\{\varepsilon, k^{-11r'_1}\}$. Then, the following estimates hold:*

$$\left\| \left(P_i (H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l} I) P_i \right)^{-1} \right\| < ck^{44r'_1} \left(\frac{k^{-11r'_1}}{\varepsilon_0} \right)^{N_i^{(1)}}, \quad (174)$$

$$\left\| \left(P_i (H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l} I) P_i \right)^{-1} \right\|_1 < ck^{44r'_1+8\gamma r_1} \left(\frac{k^{-11r'_1}}{\varepsilon_0} \right)^{N_i^{(1)}}. \quad (175)$$

Proof. Let Π be a component Π_b^j , Π_g^j or Π_w^j . By Lemmas 4.12, 4.13, 4.14 the number N of elements $\mathcal{M}^{(2)} \cap \Pi$ does not exceed $ck^{\gamma r_1+3}$. Let us recall that the set $\mathcal{M}^{(2)}$ is

defined by the formula (162), where $\mathcal{O}_{\mathbf{m}}^{(2)}$ is the union of the disks of the radius $k^{-10r'_1}$ with the centers at poles of the resolvent of k^δ -components containing $\vec{\mathcal{Z}}^{(1)}(\varphi_0) + \vec{p}_{\mathbf{m}}$. Let us consider $\mathcal{O}_{\Pi}^{(2)} = \cup_{\mathbf{m} \in \Pi \cap \mathcal{M}^{(2)}} \mathcal{O}_{\mathbf{m}}^{(2)}$ and an analogous set consisting of smaller discs: $\tilde{\mathcal{O}}_{\Pi}^{(2)} = \cup_{\mathbf{m} \in \Pi \cap \mathcal{M}^{(2)}} \tilde{\mathcal{O}}_{\mathbf{m}}^{(2)}$, where $\tilde{\mathcal{O}}_{\mathbf{m}}^{(2)}$ have the radius $k^{-11r'_1}$. Since $N < ck^{\gamma r_1 + 3}$, the total size of $\tilde{\mathcal{O}}_{\Pi}^{(2)}$ is less than $k^{-11r'_1 + \gamma r_1 + 3} = o(k^{-10r'_1})$.

First, assume $\varphi_0 \notin \tilde{\mathcal{O}}_{\Pi}^{(2)}$. Then, we can apply Lemma 3.14 and Corollary 3.16. Indeed, let us consider a k^δ -component in Π . We denote it by $\tilde{\mathcal{M}}(\mathbf{m})$ and the corresponding projection $P_{\mathbf{m}}$. By the definitions of $\mathcal{O}_{\mathbf{m}}^{(2)}$, $\tilde{\mathcal{O}}_{\mathbf{m}}^{(2)}$, the distance from φ_0 to the nearest pole of $(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}) - k^{2l}I)P_{\mathbf{m}})^{-1}$ is greater than $k^{-11r'_1}$. Applying Lemma 3.14 to these resolvents, we obtain (recall that now $p_{\mathbf{m}} > k^{-10\mu r_1}$):

$$\left\| (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P_{\mathbf{m}})^{-1} \right\| < ck^{44r'_1}, \quad (176)$$

$$\left\| (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P_{\mathbf{m}})^{-1} \right\|_1 < ck^{44r'_1 + 4\delta}. \quad (177)$$

By analogy with Corollary 3.16,

$$\begin{aligned} \left\| (P(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P)^{-1} \right\| &< ck^{44r'_1}, \\ \left\| (P(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P)^{-1} \right\|_1 &< ck^{44r'_1}L^4, \end{aligned}$$

where P is the projection for all k^δ components in Π , L is the size of Π . Arguing as in the proof of Theorem 4.1, we show that the perturbation series for the resolvent $(P_{\Pi}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P_{\Pi})^{-1}$ converges when we take $PH(\vec{\mathcal{Z}}^{(1)}(\varphi_0))P + (P_{\Pi} - P)H_0$ as the unperturbed operator. Therefore,

$$\left\| (P_{\Pi}(H(\vec{\mathcal{Z}}^{(1)}(\varphi_0)) - k^{2l}I)P_{\Pi})^{-1} \right\| < ck^{44r'_1},$$

no poles being inside of the disc. Taking into account that $|\varphi - \varphi_0| < k^{-44r'_1 - 2l - \delta}$ and $|\vec{\mathcal{Z}}^{(2)} - \vec{\mathcal{Z}}^{(1)}| = o(k^{-44r'_1 - 2l})$, we obtain

$$\left\| (P_{\Pi}(H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l}I)P_{\Pi})^{-1} \right\| < ck^{44r'_1}.$$

By Lemmas 4.12, 4.13, 4.14, $L < k^{2\gamma r_1}$,

$$\left\| (P_{\Pi}(H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l}I)P_{\Pi})^{-1} \right\|_1 < ck^{8\gamma r_1 + 44r'_1}.$$

Thus, the resolvent has no poles inside the disk around φ_0 and the estimates (174), (175) hold with $\varepsilon_0 := k^{-11r'_1}$. Second, if $\varphi \notin \tilde{\mathcal{O}}_{\Pi}^{(2)}$, then $\varphi_0 \notin \tilde{\mathcal{O}}_{\Pi}^{(2)}(11r'_1, \frac{1}{2})$. Therefore the estimate similar to the last two hold. Now estimates (174), (175) easily follow.

It remains to consider the case $\varphi_0, \varphi \in \tilde{\mathcal{O}}_{\Pi}^{(2)}$. Obviously, φ_0, φ belong to the same connected component of $\tilde{\mathcal{O}}_{\Pi}^{(2)}$ or to different components being at the distance less than $k^{-44r'_1-2l-\delta}$ from each other. We consider a $\varphi_* \in \partial\tilde{\mathcal{O}}_{\Pi}^{(2)}$, where $\partial\tilde{\mathcal{O}}_{\Pi}^{(2)}$ is the boundary of the component(s) containing φ_0, φ . Note that $\varphi_* \notin \mathcal{O}_{\mathbf{m}}^{(2)}(11r'_1, 1)$ for all $\mathbf{m} \in \Pi$. Indeed, for $\mathbf{m} \in \mathcal{M}^{(2)}$, it follows from the relations $\varphi_* \in \partial\tilde{\mathcal{O}}_{\Pi}^{(2)}$ and the definitions of $\mathcal{O}_{\Pi}^{(2)}, \tilde{\mathcal{O}}_{\Pi}^{(2)}$. If $\mathbf{m} \in \Pi \setminus \mathcal{M}^{(2)}$, then φ_0 is not in $\mathcal{O}_{\mathbf{m}}^{(2)}(10r'_1, 1)$ by the definition of $\mathcal{M}^{(2)}$. Since $\varphi_0, \varphi \in \tilde{\mathcal{O}}_{\Pi}^{(2)}$ and the length of $\tilde{\mathcal{O}}_{\Pi}^{(2)}$ is $o(k^{-10r'_1})$, we have $\varphi_* \notin \mathcal{O}_{\mathbf{m}}^{(2)}(10r'_1, \frac{1}{2})$. Now, considering as in the case $\varphi_0 \notin \tilde{\mathcal{O}}_{\Pi}^{(2)}$, we obtain that the perturbation series for the resolvent $(P_{\Pi}(H(\tilde{\mathcal{Z}}^{(1)}(\varphi_*)) - k^{2l}I)P_{\Pi})^{-1}$ converges when we take $PH(\tilde{\mathcal{Z}}^{(1)}(\varphi_*))P + (P_{\Pi} - P)H_0$ as the unperturbed operator. Therefore,

$$\left\| (P_{\Pi}(H(\tilde{\mathcal{Z}}^{(2)}(\varphi_*)) - k^{2l}I)P_{\Pi})^{-1} \right\| < ck^{44r'_1}.$$

The number of poles of the resolvent $(P_{\Pi}(H(\tilde{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l}I)P_{\Pi})^{-1}$ in $\tilde{\mathcal{O}}_{\Pi}^{(2)}$ is the same as the number of poles of the resolvent of unperturbed operator. Hence, it is N . Using the Maximum principle, we get (174) for the case $\varepsilon \leq k^{-11r'_1}$, where $N_i = N$ and depends on color of Π . Considering that the dimension of P_{Π} does not exceed $k^{8\gamma r_1}$, we obtain (175) $\square$

At last, let Π_s^j be a particular $k^{r_1/2}$ -box around $\mathbf{m}$ containing. Let P_s^j be corresponding projection.

Lemma 4.19. *Let $\varphi_0 \in \omega^{(2)}(k, \delta, \tau)$. Then, the operator $(P_s^j(H(\tilde{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l}I)P_s^j)^{-1}$ has no more than one pole in the disk $|\varphi - \varphi_0| < k^{-r'_1-\delta}$. Moreover,*

$$\left\| (P_s^j(H(\tilde{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l}I)P_s^j)^{-1} \right\| < \frac{8k^{-2l+1}}{p_{\mathbf{m}}\varepsilon_0}, \quad (178)$$

$$\left\| (P_s^j(H(\tilde{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l}I)P_s^j)^{-1} \right\|_1 < \frac{8k^{-2l+1+4r_1}}{p_{\mathbf{m}}\varepsilon_0}, \quad (179)$$

$\varepsilon_0 = \min\{\varepsilon, k^{-r'_1-\delta}\}$, where ε is the distance to the pole of the operator.

Proof. The proof is similar to that of Lemma 3.14 (part 3). Indeed, when $p_{\mathbf{m}} < k^{-5r'_1}$, the series for $\lambda^{(2)}(\tilde{\mathcal{Z}}^{(2)}(\varphi) + \vec{p}_{\mathbf{m}})$ converges in the complex $k^{-r'_1-\delta}$ neighborhood of $\omega^{(2)}(k, \delta, \tau)$ and $\lambda^{(2)}(\tilde{\mathcal{Z}}^{(2)}(\varphi) + \vec{p}_{\mathbf{m}}) = \lambda^{(1)}(\tilde{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}) + o(k^{-100r_1})$, see (131). By Lemma 10.1, the equation $\lambda^{(1)}(\tilde{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}) = k^{2l} + \varepsilon_0$, $|\varepsilon_0| \leq p_{\mathbf{m}}k^{\delta}$ has no more than two solutions in this neighborhood of $\omega^{(2)}(k, \delta, \tau)$. Using Rouché's theorem, we obtain the same fact for $\lambda^{(2)}(\tilde{\mathcal{Z}}^{(2)}(\varphi) + \vec{p}_{\mathbf{m}}) = \varepsilon_0$. It is easy to show that the analogs of Lemmas 10.2, 10.3 and 10.4 hold for $\lambda^{(2)}(\tilde{\mathcal{Z}}^{(2)}(\varphi) + \vec{p}_{\mathbf{m}})$. Thus, we obtain (178), (179). $\square$

4.4.2 Resonant and Nonresonant Sets for Step III

We divide $[0, 2\pi)$ into $[2\pi k^{44r'_1+2l+\delta}] + 1$ intervals $\Delta_m^{(2)}$ with the length not bigger than $k^{-44r'_1-2l-\delta}$. If a particular interval belongs to $\mathcal{O}^{(2)}$ we ignore it; otherwise, let $\varphi_0(m) \notin \mathcal{O}^{(2)}$ be a point inside the $\Delta_m^{(2)}$. Let

$$\mathcal{W}_m^{(2)} = \{\varphi \in \mathcal{W}^{(2)} : |\varphi - \varphi_0(m)| < 4k^{-44r'_1-2l-\delta}\}. \quad (180)$$

Clearly, neighboring sets $\mathcal{W}_m^{(2)}$ overlap (because of the multiplier 4 in the inequality), they cover $\hat{\mathcal{W}}^{(2)}$, which is the restriction of $\mathcal{W}^{(2)}$ to the $2k^{-44r'_1-2l-\delta}$ -neighborhood of $[0, 2\pi)$. For each $\varphi \in \hat{\mathcal{W}}^{(2)}$ there is an m such that $|\varphi - \varphi_0(m)| < 4k^{-44r'_1-2l-\delta}$. We consider the poles of the resolvent $(P^{(2)}(H(\vec{\mathcal{Z}}^{(2)}(\varphi)) - k^{2l})P^{(2)})^{-1}$ in $\hat{\mathcal{W}}_m^{(2)}$ and denote them by $\varphi_{mj}^{(2)}$, $j = 1, \dots, M_m$. By Corollary 4.16 the resolvent has a block structure. The number of blocks clearly cannot exceed the number of elements in $\Omega(r_2)$, i.e. k^{4r_2} . Using the estimates for the number of poles for each block, the estimate being provided by Lemma 4.18 Part 1, we can roughly estimate the number of poles of the resolvent by $k^{4r_2+r_1}$.

Next, let $r'_2 > 11r'_1$ and $\mathcal{O}_{mj}^{(3)}$ be the disc of the radius $k^{-r'_2}$ around $\varphi_{mj}^{(2)}$.

Definition 4.20. *The set*

$$\mathcal{O}^{(3)} = \cup_{mj} \mathcal{O}_{mj}^{(3)} \quad (181)$$

we call the third resonant set. The set

$$\mathcal{W}^{(3)} = \hat{\mathcal{W}}^{(2)} \setminus \mathcal{O}^{(3)} \quad (182)$$

is called the third non-resonant set. The set

$$\omega^{(3)} = \mathcal{W}^{(3)} \cap [0, 2\pi) \quad (183)$$

is called the third real non-resonant set.

Lemma 4.21. *Let $r'_2 > \mu r_2 > 44r'_1$, $\varphi \in \mathcal{W}^{(3)}$, $\varphi_0(m)$ corresponds to an interval $\Delta_m^{(2)}$ containing $\Re \varphi$. Let Π be one of the components $\Pi_s^j(\varphi_0(m))$, $\Pi_b^j(\varphi_0(m))$, $\Pi_g^j(\varphi_0(m))$, $\Pi_w^j(\varphi_0(m))$ and $P(\Pi)$ be the projection corresponding to Π . Let also $\varkappa \in \mathbb{C} : |\varkappa - \varkappa^{(2)}(\varphi)| < k^{-r'_2} k^{2\gamma r_1}$. Then,*

$$\left\| (P(\Pi) (H(\vec{\mathcal{Z}}(\varphi)) - k^{2l} I) P(\Pi))^{-1} \right\| < ck^{2\mu r_2 + r'_2 N^{(1)}}, \quad (184)$$

$$\left\| (P(\Pi) (H(\vec{\mathcal{Z}}(\varphi)) - k^{2l} I) P(\Pi))^{-1} \right\|_1 < ck^{(2\mu+1)r_2 + r'_2 N^{(1)}}, \quad (185)$$

$N^{(1)}$ corresponding to the color of Π : $N^{(1)} = 1$, $k^{\gamma r_1+3}$, $k^{\gamma r_1/2+\delta_0 r_1}$, $k^{\gamma r_1/6-\delta_0 r_1}$ for simple, black, grey and white clusters, correspondingly.

Proof. For $\vec{\mathfrak{z}} = \vec{\mathfrak{z}}^{(2)}(\varphi)$ the lemma follows immediately from the definition of $\mathcal{W}^{(3)}$ and Lemmas 4.18 and 4.19 ($p_{\mathbf{m}} > k^{-2\mu r_2}$). It is easy to see that estimates (184) and (185) are stable with respect to perturbation of $\mathfrak{z}^{(2)}$ of order $k^{-r'_2 k^{2\gamma r_1}}$. $\square$

By total size of the set $\mathcal{O}^{(3)}$ we mean the sum of the sizes of its connected components.

Lemma 4.22. *Let $r'_2 \geq (\mu + 10)r_2$, $r_2 > 45r'_1 + 2l$. Then, the size of each connected component of $\mathcal{O}^{(3)}$ is less than $k^{5r_2 - r'_2}$. The total size of $\mathcal{O}^{(3)}$ is less than $k^{-r'_2/2}$.*

Proof. Indeed, each set $\mathcal{W}_m^{(2)}$ contains no more than $k^{4r_2 + r_1}$ discs $\mathcal{O}_{mj}^{(3)}$. Therefore, the size of $\mathcal{O}^{(3)} \cap \mathcal{W}_m^{(2)}$ is less than $k^{-r'_2 + 5r_2}$. Considering that $k^{-r'_2 + 5r_2}$ is much smaller than the length of $\Delta_m^{(2)}$, we obtain that there is no connected components which go across the whole set $\mathcal{W}_m^{(2)}$ and the size of each connected component of $\mathcal{O}^{(3)}$ is less than $k^{5r_2 - r'_2}$. Considering that the number of intervals $\Delta_m^{(2)}$ is less than $k^{45r'_1 + 2l + \delta}$, we obtain the required estimate for the total size of $\mathcal{O}^{(3)}$. $\square$

Lemma 4.23. *Let $\varphi \in \mathcal{W}^{(2)}$ and C_3 be the circle $|z - k^{2l}| = k^{-2r'_2 k^{2\gamma r_1}}$. Then*

$$\left\| \left(P(r_1)(H(\vec{\mathfrak{z}}^{(2)}(\varphi)) - z)P(r_1) \right)^{-1} \right\| \leq 4^2 k^{2r'_2 k^{2\gamma r_1}}.$$

Proof. The proof is similar to the proof of Lemma 3.21 if we take into account (123) and (138). We notice also, that since in the proof of the lemma we use the estimates from the previous step along with some perturbation arguments: first, the series decomposition (cf. (120) and (123)), and second, the shift from $\vec{\mathfrak{z}}^{(1)}$ to $\vec{\mathfrak{z}}^{(2)}$, we accumulate additional factor 4. $\square$

5 Step III

Let k_* be sufficiently large to satisfy the estimates:

$$k_* \geq k_1(V, \delta, \tau), \quad k_*^{\delta/8} > 10^8 + \|V\| + \mu + 2l,$$

$k_1(V, \delta, \tau)$ being introduced in the formulation of Theorem 4.1. We also assume that k_* is such that all constants c in previous estimates (e.g. (184), (185)) satisfy $c < k_*^{\delta/8}$. Since now on we consider $k > k_*$. This restriction on k won't change in all consecutive steps. We introduce a new notation $O_T(\cdot)$: let $f(k) = O_T(k^{-\gamma})$ mean that $|f(k)| < Tk^{-\gamma}$ when $k > k_*$.

5.1 Operator $H^{(3)}$. Perturbation Formulas

Let $P(r_2)$ be an orthogonal projector onto $\Omega(r_2) := \{\mathbf{m} : \|\vec{p}_{\mathbf{m}}\| \leq k^{r_2}\}$ and $H^{(3)} = P(r_2)HP(r_2)$. From now on we assume

$$k^\delta < r_2 < k^{\gamma 10^{-7} r_1}. \tag{186}$$

Note that $45r'_1 + 2l < k^\delta < k^{\gamma 10^{-7} r_1}$ for all $k > k_*$, since $10^8 < r_1 < k^{\delta/8}$. Let $\beta = 2l - 2 - 41\mu\delta$ and

$$5\mu r_2 < r'_2 < \frac{\beta}{128} k^{\delta_0 r_1 - \delta - 3}. \quad (187)$$

We consider $H^{(3)}(\vec{\varkappa}^{(2)}(\varphi))$ as a perturbation of $\tilde{H}^{(2)}(\vec{\varkappa}^{(2)}(\varphi))$:

$$\tilde{H}^{(2)} := \tilde{P}_j^{(2)} H \tilde{P}_j^{(2)} + \left(P(r_2) - \tilde{P}_j^{(2)} \right) H_0,$$

where $H = H(\vec{\varkappa}^{(2)}(\varphi))$, $H_0 = H_0(\vec{\varkappa}^{(2)}(\varphi))$ and $\tilde{P}_j^{(2)}$ is the projection $P^{(2)}$, see (165), corresponding to $\varphi_0(j)$ in the interval $\Delta_j^{(2)}$ containing φ . Note that the operator $\tilde{H}^{(2)}$ has a block structure, the block $\tilde{P}_j^{(2)} H \tilde{P}_j^{(2)}$ being composed of smaller blocks $P_i H P_i$, $i = 0, \dots, 5$, see (169), (170). Let

$$W^{(2)} = H^{(3)} - \tilde{H}^{(2)} = P(r_2) V P(r_2) - \tilde{P}_j^{(2)} V \tilde{P}_j^{(2)}, \quad (188)$$

$$g_r^{(3)}(\vec{\varkappa}) := \frac{(-1)^r}{2\pi i r} \text{Tr} \oint_{C_3} \left(W^{(2)}(\tilde{H}^{(2)}(\vec{\varkappa}) - zI)^{-1} \right)^r dz, \quad (189)$$

$$G_r^{(3)}(\vec{\varkappa}) := \frac{(-1)^{r+1}}{2\pi i} \oint_{C_3} (\tilde{H}^{(2)}(\vec{\varkappa}) - zI)^{-1} \left(W^{(2)}(\tilde{H}^{(2)}(\vec{\varkappa}) - zI)^{-1} \right)^r dz, \quad (190)$$

where C_3 is the circle $|z - k^{2l}| = \varepsilon_0^{(3)}$, $\varepsilon_0^{(3)} = k^{-2r'_2 k^{2\gamma r_1}}$.

Theorem 5.1. *Suppose $k > k_*$, φ is in the real $k^{-r'_2 - \delta}$ -neighborhood of $\omega^{(3)}(k, \delta, \tau)$ and $\varkappa \in \mathbb{R}$, $|\varkappa - \varkappa^{(2)}(\varphi)| \leq \varepsilon_0^{(3)} k^{-2l+1-\delta}$, $\vec{\varkappa} = \varkappa(\cos \varphi, \sin \varphi)$. Then, there exists a single eigenvalue of $H^{(3)}(\vec{\varkappa})$ in the interval $\varepsilon_3(k, \delta, \tau) = \left(k^{2l} - \varepsilon_0^{(3)}, k^{2l} + \varepsilon_0^{(3)} \right)$. It is given by the absolutely converging series:*

$$\lambda^{(3)}(\vec{\varkappa}) = \lambda^{(2)}(\vec{\varkappa}) + \sum_{r=2}^{\infty} g_r^{(3)}(\vec{\varkappa}). \quad (191)$$

For coefficients $g_r^{(3)}(\vec{\varkappa})$ the following estimates hold:

$$|g_r^{(3)}(\vec{\varkappa})| < k^{-\frac{\beta}{5} k^{r_1 - \delta} - \beta(r-1)}. \quad (192)$$

The corresponding spectral projection is given by the series:

$$\mathcal{E}^{(3)}(\vec{\varkappa}) = \mathcal{E}^{(2)}(\vec{\varkappa}) + \sum_{r=1}^{\infty} G_r^{(3)}(\vec{\varkappa}), \quad (193)$$

$\mathcal{E}^{(2)}(\vec{\varkappa})$ being the spectral projection of $H^{(2)}(\vec{\varkappa})$. The operators $G_r^{(3)}(\vec{\varkappa})$ satisfy the estimates:

$$\|G_r^{(3)}(\vec{\varkappa})\|_1 < k^{-\frac{\beta}{10} k^{r_1 - \delta} - \beta r}, \quad (194)$$

$$G_r^{(3)}(\vec{\varkappa})_{\text{ss}'} = 0, \text{ when } 2rk^{\gamma r_1 + 3} + 3k^{r_1} < \|\vec{p}_{\text{s}}\| + \|\vec{p}_{\text{s}'}\|. \quad (195)$$

Corollary 5.2. *For the perturbed eigenvalue and its spectral projection the following estimates hold:*

$$\lambda^{(3)}(\vec{\varkappa}) = \lambda^{(2)}(\vec{\varkappa}) + O_2 \left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta} \right), \quad (196)$$

$$\|\mathcal{E}^{(3)}(\vec{\varkappa}) - \mathcal{E}^{(2)}(\vec{\varkappa})\|_1 < 2k^{-\frac{\beta}{10}k^{r_1-\delta}-\beta}. \quad (197)$$

$$|\mathcal{E}^{(3)}(\vec{\varkappa})_{\mathbf{s}\mathbf{s}'}| < k^{-d^{(3)}(\mathbf{s},\mathbf{s}')} , \quad \text{when } \|\vec{p}_{\mathbf{s}}\| > 4k^{r_1} \text{ or } \|\vec{p}_{\mathbf{s}'}\| > 4k^{r_1}, \quad (198)$$

$$d^{(3)}(\mathbf{s},\mathbf{s}') = \frac{1}{8}(\|\vec{p}_{\mathbf{s}}\| + \|\vec{p}_{\mathbf{s}'}\|)k^{-\gamma r_1-3}\beta + \frac{1}{10}\beta k^{r_1-\delta}.$$

Formulas (196) and (197) easily follow from (191), (192) and (193), (194). The estimate (198) follows from (193), (194), (195) and (119).

Proof. Let us consider the perturbation series

$$(H^{(3)} - z)^{-1} = \sum_{r=0}^{\infty} (\tilde{H}^{(2)} - z)^{-1} \left(-W^{(2)}(\tilde{H}^{(2)} - z)^{-1} \right)^r, \quad (199)$$

here and below all the operators are computed at $\vec{\varkappa}$. Further, we consider $\vec{\varkappa}$ and, therefore, the operators, as analytic functions of φ in $\mathcal{W}_j^{(2)}$, assuming $\varkappa$ is fixed. By (188) and (169), $W^{(2)} = V - \sum_{i=0}^5 P_i V P_i$. By assumption on $\varkappa$ and Lemmas 4.21 and 4.23,

$$\|(\tilde{H}^{(2)}(\vec{\varkappa}) - z)^{-1}\| < 2 \cdot 4^2 k^{2r'_2 k^{2\gamma r_1}}. \quad (200)$$

To check the convergence it is enough to show that

$$\left\| \left(\tilde{H}^{(2)} - z \right)^{-1} W^{(2)} \right\| < k^{-\beta}. \quad (201)$$

Then,

$$\|(H^{(3)}(\vec{\varkappa}) - z)^{-1}\| < 4^3 k^{2r'_2 k^{2\gamma r_1}}. \quad (202)$$

Let us prove (201). Operator $\tilde{H}^{(2)}$ has a block structure, different blocks being separated by the $\|\cdot\|$ distance greater than k^δ . This means that not only the blocks themselves, but also the blocks multiplied by $W^{(2)}$ have non-zero action on orthogonal subspaces. The operator $\tilde{H}^{(2)}$ acts as H_0 “outside” the blocks. Because of the block structure and the estimate $|z - k^{2l}| = k^{-2r'_2 k^{2\gamma r_1}} = o(k^{-\beta})$, it suffices to prove:

$$\left\| \left(P(r_2) - \tilde{P}_j^{(2)} \right) (H_0 - k^{2l})^{-1} V \right\| < \frac{1}{3} k^{-\beta}, \quad (203)$$

$$\left\| P(r_1) (\tilde{H}^{(2)} - z)^{-1} V (P(r_2) - P(r_1)) \right\| < \frac{1}{3} k^{-\beta}, \quad (204)$$

$$\left\| P_{nr} \left(\tilde{H}^{(2)} - k^{2l} \right)^{-1} V (P(r_2) - P_{nr}) \right\| < \frac{1}{3} k^{-\beta}, \quad (205)$$

$$\left\| P_s \left(\tilde{H}^{(2)} - k^{2l} \right)^{-1} V(P(r_2) - P_s) \right\| < \frac{1}{3} k^{-\beta}, \quad (206)$$

$$\left\| P_w \left(\tilde{H}^{(2)} - k^{2l} \right)^{-1} V(P(r_2) - P_w) \right\| < \frac{1}{3} k^{-\beta}, \quad (207)$$

$$\left\| P_g \left(\tilde{H}^{(2)} - k^{2l} \right)^{-1} V(P(r_2) - P_g) \right\| < \frac{1}{3} k^{-\beta}, \quad (208)$$

$$\left\| P_b \left(\tilde{H}^{(2)} - k^{2l} \right)^{-1} V(P(r_2) - P_b) \right\| < \frac{1}{3} k^{-\beta}, \quad (209)$$

By definition of $\tilde{P}_j^{(2)}$,

$$\left\| \left(P(r_2) - \tilde{P}_j^{(2)} \right) (H_0 - k^{2l})^{-1} \right\| < k^{-2l+2+40\mu\delta}.$$

The estimate (203) easily follows.

Let us prove (204). By Lemma 4.15,

$$P(r_1)V(P(r_2) - P(r_1)) = P(r_1)^\partial V(P(r_2) - P(r_1)),$$

where $P(r_1)^\partial$ is the projection on the boundary of $\Pi(r_1)$. Therefore, it suffice to prove:

$$\left\| P(r_1)(H^{(2)} - z)^{-1} P(r_1)^\partial \right\| < k^{-\beta-\delta/2}, \quad (210)$$

the obvious relation $P(r_1)\tilde{H}^{(2)} = H^{(2)}$ has been taken into account. As in the proof of Theorem 4.1, we consider $H^{(2)}$ as a perturbation of $\tilde{H}^{(1)}$, $H^{(2)} = \tilde{H}^{(1)} + W$. Taking into account that $\tilde{H}^{(1)}$ has a k^δ -block structure and V is a trigonometric polynomial, we obtain

$$P(\delta) \left(\left(\tilde{H}^{(1)} - z \right)^{-1} W \right)^s P^\partial(r_1) = 0, \quad \text{when } 1 \leq s \leq S, \quad S := \frac{1}{4} k^{r_1-\delta}.$$

Hence,

$$P(r_1)(H^{(2)} - z)^{-1} P^\partial(r_1) = \sum_{s=0}^{S-1} P \left(\tilde{H}^{(1)} - z \right)^{-1} A_*^s P^\partial(r_1) + P(r_1) \left(\tilde{H}^{(2)} - z \right)^{-1} A_*^S P^\partial(r_1), \quad (211)$$

where $A_* = -P \left(\tilde{H}^{(1)} - z \right)^{-1} W$, $P = P(r_1) - P(\delta)$. Considering as in the proof of Theorem 4.1,² we obtain: $\left\| P \left(\tilde{H}^{(1)} - z \right)^{-1} \right\| < ck^{-\beta-\delta} < k^{-\beta-\delta/2}$. It follows: $\|A_*\| < \frac{1}{4} k^{-\beta}$. By Theorem 4.1 and the definition of C_3 , $\left\| \left(\tilde{H}^{(2)} - z \right)^{-1} \right\| < (\varepsilon_0^{(3)})^{-1}$. Substituting

²We replace $P(r_1)$ by P , this compensates for the smallness of C_3 .

the last three estimates into (211) and taking into account that $(\varepsilon_0^{(3)})^{-1} < k^{\beta S/2}$, we obtain (210) and, therefore, (204) for all $\varphi \in \mathcal{W}_j^{(2)}$.

Next, we prove (205). by Lemma 4.15,

$$P_{nr}V(P(r_2) - P_{nr}) = P_{nr}^\partial V(P(r_2) - P_{nr}),$$

where P_{nr}^∂ is the projection on the boundary of Π_{nr} . Therefore, it suffices to prove

$$\left\| P_{nr} \left(\tilde{H}^{(2)} - k^{2l} \right)^{-1} P_{nr}^\partial \right\| < k^{-\beta-\delta/2}. \quad (212)$$

Note that Lemma 3.14 holds for any $r_1 > \delta$ (the restriction on r_1 is introduced later). Therefore, the estimates (77)–(80) hold for $\mathbf{m} \in \Omega(r_2)$. By the definition of Π_{nr} , $\varepsilon_0 > k^{-10r'_1}$ and $p_{\mathbf{m}} > k^{-5r'}$ in these estimates. It follows (see Corollary 3.15),

$$\left\| P_{nr} \left(\tilde{H}^{(2)} - k^{2l} \right)^{-1} P_{nr} \right\| < k^{40r'_1}. \quad (213)$$

Considering as in the proof of Theorem 4.1 (see the proof of (122)), we obtain (212) and, hence, (205) for all $\varphi \in \mathcal{W}_j^{(2)}$.

Next, we prove (206). Denote by $\hat{H}$ the reduction of the operator H onto a particular simple cluster i.e. $\hat{H} = P_s H P_s$ where $(P_s)_{\mathbf{mm}} = 1$ if $\mathbf{m}$ belongs to the simple cluster and $(P_s)_{\mathbf{mm}} = 0$ otherwise. By Lemma 4.21,

$$\|(\hat{H} - k^{2l})^{-1}\| \leq ck^{2\mu r_2 + r'_2}, \quad (214)$$

By Lemma 4.15,

$$P_s(\hat{H} - k^{2l})^{-1}V(P(r_2) - P_s) = P_s(\hat{H} - k^{2l})^{-1}P_s^\partial V(P(r_2) - P_s). \quad (215)$$

To obtain (206), it is enough to show

$$\left\| P_s(\hat{H} - k^{2l})^{-1}P_s^\partial \right\| < k^{-\beta-\delta/4}. \quad (216)$$

We are going to construct the perturbation formula for $P_s(\hat{H} - k^{2l})^{-1}P_s^\partial$. Let $\hat{H}_0 = P_{s,nr}H P_{s,nr} + (P_s - P_{s,nr})H_0$, where $P_{s,nr} = P_s P_{nr} = P_{nr} P_s$. The operator $\hat{H}_0$ has k^δ -block structure. It is analogous to the operator $\tilde{H}^{(1)}$ in the proof of Theorem 4.1. The perturbation formula for $P_s(\hat{H} - k^{2l})^{-1}P_s^\partial$ has the form:

$$\begin{aligned} P_s(\hat{H} - k^{2l})^{-1}P_s^\partial &= \sum_{r=0}^{R_s} P_s(\hat{H}_0 - k^{2l})^{-1} \left[-W_s(\hat{H}_0 - k^{2l})^{-1} \right]^r P_s^\partial \\ &+ P_s(\hat{H} - k^{2l})^{-1} \left[-W_s(\hat{H}_0 - k^{2l})^{-1} \right]^{R_s+1} P_s^\partial, \\ W_s &= \hat{H} - \hat{H}_0 = P_s V P_s - P_{s,nr} V P_{s,nr}, \quad R_s = \left[\frac{1}{8} k^{\frac{r_1}{2} - \delta} \right] - 1. \end{aligned} \quad (217)$$

When $\vec{p}_{\mathbf{m}'}$ belongs to the boundary of the white cluster, the $\|\cdot\|$ -distance from $\vec{p}_{\mathbf{m}'}$ to the point $\vec{p}_{\mathbf{m}} : 0 < p_{\mathbf{m}} < k^{-5r'_1}$ is $k^{r_1/2}$. Notice that $(\hat{H}_0 - k^{2l})_{\mathbf{m}\mathbf{m}'}^{-1} = 0$ if $\|\vec{p}_{\mathbf{m}} - \vec{p}_{\mathbf{m}'}\| > 8k^\delta$, since $\hat{H}_0$ has a k^δ structure. Considering that $R_s < \frac{1}{8}k^{\frac{r_1}{2}-\delta}$ (so, we never reach the central point of Π_s), we obtain that the finite sum in (217) is analytic inside $\mathcal{W}_j^{(2)}$ and is bounded by $2k^{-\beta-\delta/2}$, see (212). Moreover,

$$\left\| P_s \left[W_s (\hat{H}_0 - k^{2l})^{-1} \right]^{R_s+1} P_s^\partial \right\| \leq k^{-(R_s+1)(\beta+\delta/2)}. \quad (218)$$

Substituting (214) into (217) and taking into account (218) we get

$$\left\| P_s (\hat{H} - k^{2l})^{-1} P_s^\partial \right\| \leq k^{-\beta-\delta/4}, \quad (219)$$

when $\varphi \in \mathcal{W}^{(3)}$, $2r'_2 < \frac{1}{8}k^{r_1/2-\delta}\beta$.

Now, we prove (207). Here and in what follows we will often use the same notation for objects formally different but playing similar roles in different parts of the proof. We hope it will not lead to confusion but rather make it easier to keep the whole construction and further inductive arguments in mind. Denote by $\hat{H}$ the reduction of the operator H onto a particular white cluster i.e. $\hat{H} = PHP$ where $P_{\mathbf{m}\mathbf{m}} = 1$ if $\mathbf{m}$ belongs to the white cluster and $P_{\mathbf{m}\mathbf{m}} = 0$ otherwise. By Lemma 4.21,

$$\|(\hat{H} - k^{2l})^{-1}\| \leq ck^{2\mu r_2 + r'_2 k^{\frac{\gamma r_1}{6} - \delta_0 r_1}}. \quad (220)$$

By Lemma 4.15,

$$P_w (\hat{H} - k^{2l})^{-1} V(P(r_2) - P_w) = P_w (\hat{H} - k^{2l})^{-1} P_w^\partial V(P(r_2) - P_w). \quad (221)$$

To obtain (207), it is enough to show

$$\left\| P_w (\hat{H} - k^{2l})^{-1} P_w^\partial \right\| < k^{-\beta-\delta/4}. \quad (222)$$

We are going to construct the perturbation formula for $P_w (\hat{H} - k^{2l})^{-1} P_w^\partial$. Let $\hat{H}_0 = P_{w,nr} H P_{w,nr} + (P_w - P_{w,nr}) H_0$, where $P_{w,nr} = P_w P_{nr} = P_{nr} P_w$. The perturbation formula for $P_w (\hat{H} - k^{2l})^{-1} P_w^\partial$ has the form :

$$\begin{aligned} P_w (\hat{H} - k^{2l})^{-1} P_w^\partial &= \sum_{r=0}^{R_w} P_w (\hat{H}_0 - k^{2l})^{-1} \left[-W (\hat{H}_0 - k^{2l})^{-1} \right]^r P_w^\partial \\ &+ P_w (\hat{H} - k^{2l})^{-1} \left[-W (\hat{H}_0 - k^{2l})^{-1} \right]^{R_w+1} P_w^\partial, \\ W &= \hat{H} - \hat{H}_0, \quad R_w = \left\lceil \frac{1}{8} k^{\frac{\gamma r_1}{6} - \delta} \right\rceil - 1. \end{aligned} \quad (223)$$

When $\vec{p}_{\mathbf{m}'}$ belongs to the boundary of the white cluster, the $\|\cdot\|$ -distance from $\vec{p}_{\mathbf{m}'}$ to the closest point in $\mathcal{M}^{(2)}$ is $k^{\gamma r_1/6}$. Notice that $(\hat{H}_0 - k^{2l})_{\mathbf{m}\mathbf{m}'}^{-1} = 0$ if $\|\vec{p}_{\mathbf{m}} - \vec{p}_{\mathbf{m}'}\| > 8k^\delta$. Considering that $R_w < \frac{1}{8}k^{\frac{\gamma r_1}{6}-\delta}$ (so, we never reach the points in $\mathcal{M}^{(2)}$), we obtain that the finite sum in (223) is analytic inside $\mathcal{W}_j^{(2)}$ and is bounded by $2k^{-\beta-\delta/2}$, see (212). Moreover,

$$\left\| P_w \left[W(\hat{H}_0 - k^{2l})^{-1} \right]^{R+1} P_w^\partial \right\| \leq k^{-(R_w+1)(\beta+\delta)}. \quad (224)$$

Substituting (220) into (223) and taking into account (224) we get

$$\left\| P_w(\hat{H} - k^{2l})^{-1} P_w^\partial \right\| \leq 3k^{-\beta-\delta/2} < k^{-\beta-\delta/4}, \quad (225)$$

since $\varphi \in \mathcal{W}^{(3)}$, $2r'_2 < \frac{1}{8}k^{\delta_0 r_1 - \delta}\beta$.

Now, we prove (208). Denote a component of the grey region by Π and its boundary (see convention above) by $\partial\Pi$. Corresponding projectors are denoted by P and P^∂ respectively. Denote by $\hat{H}$ the reduction of the operator H onto a particular grey cluster i.e. $\hat{H} = PHP$. By Lemma 4.15,

$$P(\hat{H} - k^{2l})^{-1}V(P(r_2) - P) = P(\hat{H} - k^{2l})^{-1}P^\partial V(P(r_2) - P). \quad (226)$$

To obtain (208), it is enough to show

$$\left\| P(\hat{H} - k^{2l})^{-1}P^\partial \right\| < k^{-\beta-\delta/4}. \quad (227)$$

We are going to construct the perturbation formula for $P(\hat{H} - k^{2l})^{-1}P^\partial$. Recall, that the size of the neighborhood of grey boxes is $D = k^{\frac{\gamma r_1}{2} + 2\delta_0 r_1}$. Let P_i be a projector corresponding to a white or non-resonant cluster laying inside $\frac{D}{2}$ -neighborhood of $\partial\Pi$, the size of these clusters being much smaller than the size of the neighborhood. For definiteness, let $i = 1, \dots, \hat{I}$. Let $P^{(int)}$ be the projector onto all points in Π which are at least $D/2$ away of the boundary (internal points). Note that $P_i P^{(int)} = 0$. At last, put

$$P_0 := P - P^{(int)} - \sum_{i=1}^{\hat{I}} P_i. \quad (228)$$

Denote (cf. the case of a white cluster)

$$\hat{H}_0 := \sum_{i=1}^{\hat{I}} P_i H P_i + P^{(int)} H P^{(int)} + H_0 P_0, \quad (229)$$

$$W = \hat{H} - \hat{H}_0 = PVP - \sum_{i=1}^{\hat{I}} P_i V P_i - P^{(int)} V P^{(int)}. \quad (230)$$

We are going to use perturbation arguments between $\hat{H}_0$ and $\hat{H}$. Let R be the smallest natural number for which

$$\mathcal{A}_R := P^{(int)} \left[W(\hat{H}_0 - k^{2l})^{-1} \right]^{R+1} P^\partial \neq 0, \quad W := \hat{H} - \hat{H}_0. \quad (231)$$

It is proven in Appendix 4 that $R > \frac{1}{64} k^{(\frac{\gamma}{2} + 2\delta_0)r_1 - \delta}$. Therefore,

$$\begin{aligned} (\hat{H} - k^{2l})^{-1} P^\partial &= \sum_{r=0}^{R-1} (\hat{H}_0 - k^{2l})^{-1} \left[-(P - P^{(int)}) W(\hat{H}_0 - k^{2l})^{-1} \right]^r P^\partial \\ &+ (\hat{H} - k^{2l})^{-1} \left[-(P - P^{(int)}) W(\hat{H}_0 - k^{2l})^{-1} \right]^R P^\partial. \end{aligned} \quad (232)$$

The first term in the RHS of (232) contains only non-resonant and white clusters. Thus, we can use the estimates obtained before in the case of non-resonant and white clusters (see (205), (207)). To estimate the second term we, first, notice that

$$\left\| \left[-(P - P^{(int)}) W(\hat{H}_0 - k^{2l})^{-1} \right]^R P^\partial \right\| \leq k^{-\beta R} \leq k^{-\frac{\beta}{64} k^{(\frac{\gamma}{2} + 2\delta_0)r_1 - \delta}}. \quad (233)$$

By Lemma 4.21 ,

$$\|(\hat{H} - k^{2l})^{-1}\| \leq c k^{2\mu r_2 + r'_2 k^{(\frac{\gamma}{2} + \delta_0)r_1}}. \quad (234)$$

Now, considering that $2r'_2 < \frac{\beta}{64} k^{\delta_0 r_1 - \delta}$ and combining the estimates above, we obtain (227) and, therefore, (208).

We prove (209) in the analogous way. Indeed, denote a component of the black region by Π and its boundary (see convention above) by $\partial\Pi$. Corresponding projectors are denoted by P and P^∂ respectively. Again, $\hat{H} := PHP$ and, by Lemma 4.15,

$$P(\hat{H} - k^{2l})^{-1} V(P(r_2) - P) = P(\hat{H} - k^{2l})^{-1} P^\partial V(P(r_2) - P). \quad (235)$$

To obtain (209), it is enough to show

$$\left\| P(\hat{H} - k^{2l})^{-1} P^\partial \right\| < k^{-\beta - \delta}. \quad (236)$$

We are going to construct the perturbation formula for $P(\hat{H} - k^{2l})^{-1} P^\partial$. Recall, that the size of the neighborhood of black boxes is $D = k^{\gamma r_1 + \delta_0 r_1}$. Put $\hat{H} = PHP$ (cf. the case of white and grey clusters). Let P_i be a projector corresponding to a grey, white or non-resonant cluster laying inside $\frac{D}{2}$ -neighborhood of $\partial\Pi$, the size of these clusters being much smaller than the size of the neighborhood. For definiteness, let $i = 1, \dots, \hat{I}$. Let $P^{(int)}$ be the projector onto all points in Π which are at least $D/2$ away of the boundary (internal points). Again, we define P_0 , $\hat{H}_0$ and W by formulas (228), (229) and (230). We are going to use perturbation arguments between $\hat{H}_0$ and $\hat{H}$. Let R be the smallest positive

integer for which (231) holds in the case of a black cluster. It is proven in Appendix 5 that $R > \frac{1}{64}k^{(\gamma r_1 + \delta_0 r_1 - \delta)}$. Next, we use (232). The first term in the RHS of (232) contains only non-resonant, white and grey clusters. Thus, we can use the estimates (205)-(208) obtained before in the case of non-resonant, white and grey clusters. To estimate the second term we, first, notice that

$$\left\| \left[-(P - P^{(int)})W(\hat{H}_0 - k^{2l})^{-1} \right]^R P^\partial \right\| < k^{-\beta R} < k^{-\frac{\beta}{64}k^{\gamma r_1 + \delta_0 r_1 - \delta}}. \quad (237)$$

By Lemma 4.21,

$$\|(\hat{H} - k^{2l})^{-1}\| \leq ck^{2\mu r_2 + r'_2 k^{\gamma r_1 + 3}}. \quad (238)$$

Now, choosing $2r'_2 < \frac{\beta}{64}k^{\delta_0 r_1 - \delta - 3}$ and combining the estimates above we obtain (236) and, therefore, (209).

Estimates (203) – (209) provide convergence of the series for the resolvent. Integrating the resolvent over the contour we get (191) and (193).

Proof of (194) is analogous to that of (115) in Theorem 4.1. Indeed, we consider the operator $A = W^{(2)}(\tilde{H}^{(2)} - z)^{-1}$ and represent it as $A = A_0 + A_1 + A_2$, where $A_0 = (P(r_2) - \mathcal{E}^{(2)}(\vec{\mathcal{Z}}))A(P(r_2) - \mathcal{E}^{(2)}(\vec{\mathcal{Z}}))$, $A_1 = (P(r_2) - \mathcal{E}^{(2)}(\vec{\mathcal{Z}}))A\mathcal{E}^{(2)}(\vec{\mathcal{Z}})$, $A_2 = \mathcal{E}^{(2)}(\vec{\mathcal{Z}})A(P(r_2) - \mathcal{E}^{(2)}(\vec{\mathcal{Z}}))$. Note that $\mathcal{E}^{(2)}(\vec{\mathcal{Z}})W^{(2)}\mathcal{E}^{(2)}(\vec{\mathcal{Z}}) = 0$, because of (188). We see that

$$\oint_{C_3} (\tilde{H}^{(2)} - z)^{-1} A_0^r dz = 0,$$

since the integrand is a holomorphic function inside C_3 . Therefore,

$$G_r^{(3)}(\vec{\mathcal{Z}}) = \frac{(-1)^r}{2\pi i} \sum_{j_1, \dots, j_r=0,1,2, j_1^2 + \dots + j_r^2 \neq 0} \oint_{C_3} (\tilde{H}^{(2)} - z)^{-1} A_{j_1} \dots A_{j_r} dz. \quad (239)$$

At least one of indices in each term is equal to 1 or 2. Let us show that

$$\|A_2\|_1 < \frac{1}{2}k^{-\frac{\beta}{10}k^{r_1 - \delta - \beta}}. \quad (240)$$

First, we notice that

$$\begin{aligned} \mathcal{E}^{(2)}W^{(2)}(P(r_2) - \mathcal{E}^{(2)}) &= \mathcal{E}^{(2)}W^{(2)}(P(r_2) - P(r_1)) = \\ \mathcal{E}^{(2)}P^\partial(r_1)W^{(2)}(P(r_2) - P(r_1)) &= \mathcal{E}^{(2)}P^\partial(r_1)W^{(2)} \end{aligned}$$

by (188). Hence, $A_2 = \mathcal{E}^{(2)}P^\partial(r_1)A(P(r_2) - \mathcal{E}^{(2)})$. Using (119), we obtain $\|\mathcal{E}^{(2)}P^\partial(r_1)\| < k^{-\frac{\beta}{10}k^{r_1 - \delta}}$. Considering that $\mathcal{E}^{(2)}$ is a one-dimensional projection, we obtain the same estimate for $\mathbf{S}_1$ -norm. Now (240) easily follows. Applying the same trick as in the proof of (115) we obtain (194).

Let us obtain the estimate for $g_r^{(3)}(\vec{\varkappa})$. Obviously,

$$g_r^{(3)}(\vec{\varkappa}) = \frac{(-1)^r}{2\pi i r} \sum_{j_1, \dots, j_r=0,1,2, j_1^2+\dots+j_r^2 \neq 0} \text{Tr} \oint_{C_3} A_{j_1} \dots A_{j_r} dz. \quad (241)$$

Note that each term contains both A_1 and A_2 , since we compute the trace of the integral. Using (240) and repeating arguments from the proof of (113), we obtain (192).

The estimate (195) follows from the fact that the biggest white, grey or black component has the size not greater than $k^{\gamma r_1+3}$. Therefore the biggest block of $\tilde{H}^{(2)}$ not coinciding with $P(r_1)HP(r_1)$ has the size not greater than $k^{\gamma r_1+3}$.

□

It is easy to see that coefficients $g_r^{(3)}(\vec{\varkappa})$ and operators $G_r^{(3)}(\vec{\varkappa})$ can be analytically extended into the complex $k^{-r'_2-\delta}$ neighborhood of $\omega^{(3)}$ (in fact, into $k^{-r'_2-\delta}$ -neighborhood of $\mathcal{W}^{(3)}$) as functions of φ and to the complex $(\varepsilon_0^{(3)} k^{-2l+1-\delta})$ -neighborhood of $\varkappa = \varkappa^{(2)}(\varphi)$ as functions of $\varkappa$, estimates (192), (194) being preserved. Now, we use formulae (189), (191) to extend $\lambda^{(3)}(\vec{\varkappa}) = \lambda^{(3)}(\varkappa, \varphi)$ as an analytic function. Obviously, series (191) is differentiable. Using Cauchy integral and Lemma 4.3 we get the following lemma.

Lemma 5.3. *Under conditions of Theorem 5.1 the following estimates hold when $\varphi \in \omega^{(3)}(k, \delta)$ or its complex $k^{-r'_2-\delta}$ -neighborhood and $\varkappa \in \mathbb{C} : |\varkappa - \varkappa^{(2)}(\varphi)| < \varepsilon_0^{(3)} k^{-2l+1-\delta}$.*

$$\lambda^{(3)}(\vec{\varkappa}) = \lambda^{(2)}(\vec{\varkappa}) + O_2 \left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta} \right), \quad (242)$$

$$\frac{\partial \lambda^{(3)}}{\partial \varkappa} = \frac{\partial \lambda^{(2)}}{\partial \varkappa} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta} M_1 \right), \quad M_1 := \frac{k^{2l-1+\delta}}{\varepsilon_0^{(3)}}, \quad (243)$$

$$\frac{\partial \lambda^{(3)}}{\partial \varphi} = \frac{\partial \lambda^{(2)}}{\partial \varphi} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta+r'_2+\delta} \right), \quad (244)$$

$$\frac{\partial^2 \lambda^{(3)}}{\partial \varkappa^2} = \frac{\partial^2 \lambda^{(2)}}{\partial \varkappa^2} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta} M_1^2 \right), \quad (245)$$

$$\frac{\partial^2 \lambda^{(3)}}{\partial \varkappa \partial \varphi} = \frac{\partial^2 \lambda^{(2)}}{\partial \varkappa \partial \varphi} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta+r'_2+\delta} M_1 \right), \quad (246)$$

$$\frac{\partial^2 \lambda^{(3)}}{\partial \varphi^2} = \frac{\partial^2 \lambda^{(2)}}{\partial \varphi^2} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta+2r'_2+2\delta} \right). \quad (247)$$

Corollary 5.4. *All “ O_2 ”-s on the right hand sides of (242)-(247) can be written as $O_1 \left(k^{-\frac{1}{10}\beta k^{r_1-\delta}} \right)$.*

5.2 Isoenergetic Surface for Operator $H^{(3)}$

Lemma 5.5. 1. For every $\lambda := k^{2l}$, $k > k_*$, and φ in the real $\frac{1}{2}k^{-r'_2-\delta}$ -neighborhood of $\omega^{(3)}(k, \delta, \tau)$, there is a unique $\varkappa^{(3)}(\lambda, \varphi)$ in the interval

$$I_2 := [\varkappa^{(2)}(\lambda, \varphi) - \varepsilon_0^{(3)}k^{-2l+1-\delta}, \varkappa^{(2)}(\lambda, \varphi) + \varepsilon_0^{(3)}k^{-2l+1-\delta}],$$

such that

$$\lambda^{(3)}(\vec{\varkappa}^{(3)}(\lambda, \varphi)) = \lambda, \quad \vec{\varkappa}^{(3)}(\lambda, \varphi) := \varkappa^{(3)}(\lambda, \varphi)\vec{\nu}(\varphi). \quad (248)$$

2. Furthermore, there exists an analytic in φ continuation of $\varkappa^{(3)}(\lambda, \varphi)$ to the complex $\frac{1}{2}k^{-r'_2-\delta}$ -neighborhood of $\omega^{(3)}(k, \delta, \tau)$ such that $\lambda^{(3)}(\vec{\varkappa}^{(3)}(\lambda, \varphi)) = \lambda$. Function $\varkappa^{(3)}(\lambda, \varphi)$ can be represented as $\varkappa^{(3)}(\lambda, \varphi) = \varkappa^{(2)}(\lambda, \varphi) + h^{(3)}(\lambda, \varphi)$, where

$$|h^{(3)}(\varphi)| = O_1\left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta-2l+1}\right), \quad (249)$$

$$\frac{\partial h^{(3)}}{\partial \varphi} = O_2\left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta-2l+1+r'_2+\delta}\right), \quad \frac{\partial^2 h^{(3)}}{\partial \varphi^2} = O_4\left(k^{-\frac{1}{5}\beta k^{r_1-\delta}-\beta-2l+1+2r'_2+2\delta}\right). \quad (250)$$

Proof. The proof is completely analogous to that of Lemma 3.11, estimates (242)–(247) being used. $\square$

Let us consider the set of points in $\mathbb{R}^2$ given by the formula: $\vec{\varkappa} = \vec{\varkappa}^{(3)}(\varphi)$, $\varphi \in \omega^{(3)}(k, \delta, \tau)$. By Lemma 5.5 this set of points is a slight distortion of $\mathcal{D}_2$. All the points of this curve satisfy the equation $\lambda^{(3)}(\vec{\varkappa}^{(3)}(\varphi)) = k^{2l}$. We call it isoenergetic surface of the operator $H^{(3)}$ and denote by $\mathcal{D}_3$.

5.3 Preparation for Step IV

5.3.1 Properties of the Quasiperiodic Lattice. Continuation

Let

$$\mathcal{S}^{(2)}(k, \xi) := \{\vec{\varkappa} \in \mathbb{R}^2 : \|(H^{(2)}(\vec{\varkappa}) - k^{2l})^{-1}\| > k^\xi\}. \quad (251)$$

The main purpose of this section is to estimate the number of points $\vec{\varkappa}_0 + \vec{p}_{\mathbf{m}}$, $\|\vec{p}_{\mathbf{m}}\| < k^{r_2}$ in $\mathcal{S}^{(2)}(k, \xi)$, $\vec{\varkappa}_0$ being fixed. In fact, we prove a more subtle result, see Lemma 5.13.

We consider $\vec{p}_{\mathbf{m}} = 2\pi(\mathbf{s}_1 + \alpha\mathbf{s}_2)$ with integer vectors $\mathbf{s}_j$ such that $|\mathbf{s}_j| \leq 4k^{r_2}$. We repeat the arguments from the beginning of Section 4.3. Namely, let $(q, p) \in \mathbb{Z}^2$ be a pair such that $0 < q \leq 4k^{r_2}$ and

$$|\alpha q + p| \leq 16k^{-r_2}. \quad (252)$$

We choose a pair (p, q) which gives the best approximation. In particular, p and q are mutually simple. Put $\epsilon_q := \alpha + \frac{p}{q}$. We have

$$k^{-2r_2\mu} \leq |\epsilon_q| \leq 16q^{-1}k^{-r_2}. \quad (253)$$

We write $\mathbf{s}_2$ in the form

$$\mathbf{s}_2 = q\mathbf{s}'_2 + \mathbf{s}''_2 \quad (254)$$

with integer vectors $\mathbf{s}'_2$ and $\mathbf{s}''_2$, $0 \leq (\mathbf{s}''_2)_j < q$ for $j = 1, 2$. Hence, $|(\mathbf{s}'_2)_j| \leq 4k^{r_2}/q + 1$. It follows

$$(2\pi)^{-1}\vec{p}_{\mathbf{m}} = (\mathbf{s}_1 - p\mathbf{s}'_2) + \left(-\frac{p}{q}\mathbf{s}''_2 + \epsilon_q\mathbf{s}''_2\right) + \epsilon_q q\mathbf{s}'_2.$$

Denote $\mathbf{s} := \mathbf{s}_1 - p\mathbf{s}'_2$. Then $|\mathbf{s}| \leq 8k^{r_2}$. The number of different vectors $\tilde{\mathbf{s}} := -\frac{p}{q}\mathbf{s}''_2 + \epsilon_q\mathbf{s}''_2$ is not greater than $(2q)^2$. For each fixed pair $\tilde{\mathbf{s}}, \mathbf{s}$ we obtain a lattice parameterized by $\mathbf{s}'_2$. We call this lattice a cluster corresponding to given $\tilde{\mathbf{s}}, \mathbf{s}$. Each cluster, obviously, is a square lattice with the step $\epsilon_q q$. It contains no more than $(9k^{r_2}q^{-1})^2$ elements, since $|(\mathbf{s}'_2)_j| \leq 4k^{r_2}q^{-1} + 1$, $j = 1, 2$. The size of each cluster is less than $5|\epsilon_q|k^{r_2}$. As before we have the following statements.

Lemma 5.6. *Suppose that ϵ_q satisfies the inequality*

$$|\epsilon_q| \leq \frac{1}{64}q^{-1}k^{-r_2}. \quad (255)$$

Then, the size of each cluster is less than $\frac{1}{8q}$. The distance between clusters is greater than $\frac{1}{2q}$.

Lemma 5.7. *The number of vectors $\vec{p}_{\mathbf{m}}$, satisfying the inequalities $\|\vec{p}_{\mathbf{m}}\| < 2k^{r_2}$, $p_{\mathbf{m}} < |\epsilon_q|qk^{r_2/3}$, does not exceed $k^{2r_2/3}$.*

Lemma 5.8. *Suppose q in the inequality (141) satisfies the estimate $q > k^{2r_2/3}$. Then, the number of vectors $\vec{p}_{\mathbf{m}}$, $\|\vec{p}_{\mathbf{m}}\| < 2k^{r_2}$, satisfying the inequality $p_{\mathbf{m}} < k^{-2r_2/3}$ does not exceed $2^{12} \cdot k^{2r_2/3}$.*

We consider the matrix $H^{(2)}(\vec{\mathcal{Z}}) = P(\gamma r_1)H(\vec{\mathcal{Z}})P(\gamma r_1)$ where $\vec{\mathcal{Z}} \in \mathbb{R}^2$, $P(\gamma r_1)$ is the orthogonal projection corresponding to $\Omega(\gamma r_1)$.³ We construct the block structure in $H^{(2)}(\vec{\mathcal{Z}})$ analogous to that in Step II. The difference is that now we consider any $\vec{\mathcal{Z}} \in \mathbb{R}^2$, not only $\vec{\mathcal{Z}}$ being close to $\vec{\mathcal{Z}}^{(1)}(\varphi)$. Indeed, we call $\mathbf{m} \in \Omega(\gamma r_1)$ non-resonant if (cf. (69))

$$\left| |\vec{\mathcal{Z}} + \vec{p}_{\mathbf{m}}|^2 - k^2 \right| > k^{-40\mu\delta}. \quad (256)$$

Obviously, this estimate is stable in the $k^{-41\mu\delta-1}$ -neighborhood of a given $\vec{\mathcal{Z}}$. Hence, the definition of a non-resonant $\mathbf{m}$ is stable in this neighborhood up to a multiplier $1 + o(1)$ in the r.h.s. of (256). Around each resonant $\mathbf{m}$ we construct k^δ -boxes/clusters (see (73)). Let $P_{\mathbf{m}}$ be the projection on the k^δ -cluster containing $\mathbf{m}$. If

$$\left\| (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}) - k^{2l})P_{\mathbf{m}})^{-1} \right\| < k^{4\gamma r'_1}, \quad (257)$$

³It is a slight abuse of notations, since $H^{(2)}$ in Step II was defined for $\gamma = 1$.

then we call the k^δ -cluster effectively non-resonant (cf. (103)) for a given $\vec{\mathcal{Z}}$. Note, that the above estimate and, therefore, the definition of an effectively non-resonant k^δ -cluster is stable in the $k^{-4\gamma r'_1 - 2l + 1 - \delta}$ -neighborhood of a given $\vec{\mathcal{Z}}$. The k^δ -clusters, where (257) is not valid, are called effectively resonant k^δ -clusters. Thus, we have constructed a block structure in $H^{(2)}(\vec{\mathcal{Z}})$, which is stable in the $k^{-4\gamma r'_1 - 2l + 1 - \delta}$ -neighborhood of a given $\vec{\mathcal{Z}}$.

Definition 5.9. We denote by $J(\vec{\mathcal{Z}})$ the number of the effectively resonant k^δ -clusters in $H^{(2)}(\vec{\mathcal{Z}})$ for a given $\vec{\mathcal{Z}}$. Further (with a slight abuse of notations) we consider $J(\vec{\mathcal{Z}})$ to be constant in the $2k^{-4\gamma r'_1 - 2l + 1 - \delta}$ -neighborhood of a given $\vec{\mathcal{Z}}$.

Let $\vec{\mathcal{Z}} = \mathbf{a}\tau_1 + \mathbf{b}$, $|\mathbf{a}| = 1$, $|\mathbf{b}| < 4k^{\gamma r_1}$. We consider $H^{(2)}(\vec{\mathcal{Z}})$ as a function of τ_1 in the complex $2k^{-\rho_1}$ -neighborhood of zero, $\rho_1 = 4\gamma r'_1 + 2l - 1 + \delta$.

Lemma 5.10. The resolvent $(H^{(2)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}$ has no more than $8J(\mathbf{b})$ poles in the complex $2k^{-\rho_1}$ -neighborhood of zero. It satisfies the following estimate in the complex $k^{-\rho_1}$ -neighborhood of zero:

$$\|(H^{(2)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}\| < k^{16\rho_1} \left(\frac{k^{-\rho_1}}{\varepsilon_0} \right)^{8J(\vec{\mathcal{Z}})}, \quad (258)$$

where $\varepsilon_0 = \min\{k^{-2\rho_1}, \varepsilon\}$, ε being the distance to the nearest pole.

Proof. Recall (Definition 5.9) that $J(\vec{\mathcal{Z}})$ may be considered to be constant in $2k^{-\rho_1}$ -neighborhood of $\mathbf{b}$. Hence, $J(\vec{\mathcal{Z}}) = J(\mathbf{b})$ for such $\vec{\mathcal{Z}}$ -s. Let us consider the collection of all k^δ -clusters $P_{\mathbf{m}}H(\vec{\mathcal{Z}})P_{\mathbf{m}}$ for $H^{(2)}(\vec{\mathcal{Z}})$, τ_1 being in $2k^{-\rho_1}$ -neighborhood of zero. Note that the collection is the same for all such $\vec{\mathcal{Z}}$. We construct the corresponding block operator $\tilde{H}^{(1)}(\vec{\mathcal{Z}})$:

$$\tilde{H}^{(1)}(\vec{\mathcal{Z}}) = \sum P_{\mathbf{m}}H P_{\mathbf{m}} + H_0(I - \sum P_{\mathbf{m}}).$$

If a k^δ -cluster $P_{\mathbf{m}}H(\vec{\mathcal{Z}})P_{\mathbf{m}}$ is effectively non-resonant, then its resolvent, obviously, has no poles τ_1 in the $2k^{-\rho_1}$ -neighborhood of $\tau_1 = 0$. Considering that a k^δ -cluster contains no more than 4 squares (Appendix 2), we obtain, that the resolvent of each effectively resonant k^δ -cluster has no more than 8 poles τ_{1j} in the $2k^{-40\mu\delta - \delta}$ -neighborhood of $\tau_1 = 0$. Indeed, the relation opposite to (256) can hold for no more than four different $\mathbf{m}$ -s. Each function $|\vec{\mathcal{Z}} + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2$ is a quadratic polynomial with respect to τ_1 . It is easy to see that $||\vec{\mathcal{Z}} + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} - k^{2l}| > k^{2l-2-80\mu\delta-4\delta}$ when $\tau_1 \notin \mathbf{D}_0$, $\mathbf{D}_0$ being the $k^{-40\mu\delta-2\delta}$ -neighborhood of the roots τ_{1j} of the polynomials. Obviously, $\mathbf{D}_0$ consists of at most 8 discs. We consider only those connected components of $\mathbf{D}_0$ which are inside $2k^{-40\mu\delta-\delta}$ disk around $\tau_1 = 0$ (components of $\mathbf{D}_0$ are much smaller than the size of the disk). The perturbation series for the resolvent of $P_{\mathbf{m}}H(\vec{\mathcal{Z}})P_{\mathbf{m}}$ with respect to $P_{\mathbf{m}}H_0(\vec{\mathcal{Z}})P_{\mathbf{m}}$ ($V = 0$) converges on the boundaries of these components and the following estimate holds there:

$$\|(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}) - k^{2l})P_{\mathbf{m}})^{-1}\| < k^{-2l+2+80\mu\delta+5\delta}.$$

Hence, the resolvents of $P_{\mathbf{m}}H_0(\vec{z})P_{\mathbf{m}}$ and $P_{\mathbf{m}}H(\vec{z})P_{\mathbf{m}}$ have the same number of poles inside each component of $\mathbf{D}_0$. This means that the resolvent of each effectively resonant cluster $P_{\mathbf{m}}H(\vec{z})P_{\mathbf{m}}$ has no more than 8 poles τ_{1j} in the $k^{-40\mu\delta-\delta}$ -neighborhood of $\tau_1 = 0$. By the maximum principle,

$$\| (P_{\mathbf{m}}(H(\vec{z}) - k^{2l})P_{\mathbf{m}})^{-1} \| < k^{-2l+2+80\mu\delta+5\delta} \left(\frac{16k^{-40\mu\delta-2\delta}}{k^{-2\rho_1}} \right)^8 < \frac{1}{4}k^{16\rho_1}$$

at the distance greater than $k^{-2\rho_1}$ from the poles. Therefore, resolvent $(\tilde{H}^{(1)}(\vec{z}) - k^{2l})^{-1}$ has no more than $8J(\vec{z})$ poles τ_{1j} in the complex $k^{-40\mu\delta-\delta}$ -neighborhood of $\tau_1 = 0$ and satisfies the estimate

$$\|(\tilde{H}^{(1)}(\vec{z}) - k^{2l})^{-1}\| < \frac{1}{4}k^{16\rho_1} \quad (259)$$

at the distance greater than $k^{-2\rho_1}$ from the poles. Let us consider the union of $k^{-2\rho_1}$ neighborhoods of these poles. It may consist from several connected components. We are interested only in those intersecting with the $2k^{-\rho_1}$ disk around $\tau_1 = 0$. We denote their union by $\mathbf{D}$. Using a rough estimate $J(\vec{z}) < k^{4\gamma r_1}$ gives that $\mathbf{D}$ belongs to the $3k^{-\rho_1}$ -neighborhood of zero. Thus, (259) holds outside $\mathbf{D}$. Considering as before (see the proof of Theorem 4.1, (120)–(125)), we can show that the perturbation series for the resolvent $(H^{(2)}(\vec{z}) - k^{2l})^{-1}$ with respect to $(\tilde{H}^{(1)}(\vec{z}) - k^{2l})^{-1}$ converges on the boundary of $\mathbf{D}$ and

$$\|(H^{(2)}(\vec{z}) - k^{2l})^{-1}\| < k^{16\rho_1}$$

outside $\mathbf{D}$, the resolvent has no more than $8J(\vec{z})$ poles in $\mathbf{D}$. Using again the maximum principle we obtain (258). $\square$

Note that each connected component of $\mathcal{S}^{(2)}(k, \xi)$, see (251), is bounded by the curves $D(\vec{z}, k^{2l} \pm k^{-\xi}) = 0$, where $D(\vec{z}, \lambda) = \det (H^{(2)}(\vec{z}) - \lambda)$.

Lemma 5.11. *Let $\mathbf{l}$ be a segment of a straight line in $\mathbb{R}^2$,*

$$\mathbf{l} := \{ \vec{z} = \mathbf{a}\tau_1 + \mathbf{b}, \tau_1 \in (0, \eta) \}, \quad |\mathbf{a}| = 1, \quad |\mathbf{b}| < 4k^{\gamma r_1}, \quad 0 < \eta < k^{-5\gamma r'_1}. \quad (260)$$

Suppose both ends of $\mathbf{l}$ belong to a connected component of $\mathcal{S}^{(2)}(k, \xi)$. If ξ is sufficiently large, namely, $\xi \geq 30J(\mathbf{b}) \ln_k \frac{1}{\eta}$, then, there is an inner part $\mathbf{l}'$ of the segment, which is not in $\mathcal{S}^{(2)}(k, \xi)$.

Corollary 5.12. *Let $\vec{z} \in \mathcal{S}^{(2)}(k, \xi)$ and $\xi/30J(\vec{z}) > 10\gamma r'_1$. Then the distance from $\vec{z}$ to the boundary of $\mathcal{S}^{(2)}(k, \xi)$ is less than $k^{-\xi/30J(\vec{z})}$.*

Proof of the corollary. Let us consider a segment of the length $\eta = k^{-\xi/30J(\vec{z})}$ starting at $\vec{z}$. By the statement of the lemma it intersects a boundary $D(\vec{z}, k^{2l} \pm k^{-\xi}) = 0$.

Proof. Choose $\varepsilon = \eta^2$ in (258). Using the hypothesis of the lemma, we obtain that the right-hand side of (258) is less than k^ε outside the discs. Let us estimate the total

size (sum of the sizes) of the discs. Indeed, the size of each disc is 2ε and the number of discs is, obviously, less than $16k^{4\gamma r_1}$. Therefore, the total size admits the estimate from above: $32\varepsilon k^{4\gamma r_1} < \eta$, since $\eta < k^{-5\gamma r_1'}$. This means there is a part $\mathbf{l}'$ of $\mathbf{l}$ outside these discs. By (258), this part is outside $\mathcal{S}^{(2)}(k, \xi)$, when ξ is as described in the statement of the lemma. $\square$

Let $\vec{\mathfrak{x}}_0 \in \mathbb{R}^2$ be fixed and $\mathcal{N}(k, r_2, \vec{\mathfrak{x}}_0, J_0)$ be the following subset of the lattice $\vec{\mathfrak{x}}_0 + \vec{p}_{\mathbf{n}}$, $\mathbf{n} \in \Omega(r_2)$:

$$\mathcal{N}(k, r_2, \vec{\mathfrak{x}}_0, J_0) = \{ \vec{\mathfrak{x}}_0 + \vec{p}_{\mathbf{n}} : \mathbf{n} \in \Omega(r_2) : J(\vec{\mathfrak{x}}_0 + \vec{p}_{\mathbf{n}}) \leq J_0 \},$$

J being defined by Definition 5.9. Thus, $\mathcal{N}$ includes only such $\mathbf{n}$ that the surrounding $k^{\gamma r_1}$ -block contains less than J_0 of effectively resonant k^δ -clusters. Let $N(k, r_2, \vec{\mathfrak{x}}_0, J_0, \xi)$ be the number of points $\vec{\mathfrak{x}}_0 + \vec{p}_{\mathbf{n}}$ in $\mathcal{S}^{(2)}(k, \xi) \cap \mathcal{N}(k, r_2, \vec{\mathfrak{x}}_0, J_0)$.

Lemma 5.13. *If $r_2 > 10\gamma r_1'$ and $\xi > 60\mu r_2 J_0$, then*

$$N(k, r_2, \vec{\mathfrak{x}}_0, J_0, \xi) \leq k^{\frac{2}{3}r_2 + 43l\gamma r_1}. \quad (261)$$

Proof. Let us call a subset $\tilde{\mathcal{S}}$ of $\mathcal{S}^{(2)}(k, \xi)$ elementary if it can be described by a formula of the type:

$$\tilde{\mathcal{S}} := \{ \vec{\mathfrak{x}} : a < \varkappa_1 < b, f_1(\varkappa_1) < \varkappa_2 < f_2(\varkappa_1) \},$$

where the curves $\varkappa_2 = f_i(\varkappa_1)$, $i = 1, 2$, belong to the boundary of $\mathcal{S}^{(2)}(k, \xi)$, have the lengths less than 1, functions $f_i(\varkappa_1)$ are monotone, continuously differentiable and have no inflection points. We assume that the boundaries $\varkappa_2 = f_i(\varkappa_1)$ are parameterized by $\varkappa_1$ for definiteness. The set where $\varkappa_1 = f_i(\varkappa_2)$, $a < \varkappa_2 < b$, is completely analogous.

Next, we show that the number of points in $\tilde{\mathcal{S}} \cap \mathcal{N}(k, r_2, \vec{\mathfrak{x}}_0, J_0)$ does not exceed $2^{14}k^{\frac{2}{3}r_2}$. Indeed, let us consider a segment $\vec{p}_{\mathbf{n}-\mathbf{n}'}$ between two points $\vec{\mathfrak{x}}_0 + \vec{p}_{\mathbf{n}}$ and $\vec{\mathfrak{x}}_0 + \vec{p}_{\mathbf{n}'}$ in $\tilde{\mathcal{S}}$. Obviously, $\|\vec{p}_{\mathbf{n}-\mathbf{n}'}\| < 2k^{r_2}$ and $p_{\mathbf{n}-\mathbf{n}'} > k^{-\mu r_2}$. The direction of the segment cannot be parallel to the axis $\varkappa_2$ by Corollary 5.12. We enumerate the points $\vec{\mathfrak{x}}_0 + \vec{p}_{\mathbf{n}} \in \tilde{\mathcal{S}} \cap \mathcal{N}(k, r_2, \vec{\mathfrak{x}}_0, J_0)$ in the order of the increasing first coordinate and connect subsequent points by segments. Consider all segments with the length greater or equal to $\frac{1}{64}k^{-\frac{2r_2}{3}}$. The number of such segments does not exceed $128k^{\frac{2r_2}{3}}$, since they are much longer than the width of $\tilde{\mathcal{S}}$ (Corollary 5.12). It remains to estimate the number of segments with the length less than $\frac{1}{64}k^{-\frac{2r_2}{3}}$.

First, we prove that no more than two segments $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$, $\vec{p}_{\mathbf{n}_2-\mathbf{n}'_2}$ can be parallel to each other, if they are in the same elementary component $\tilde{\mathcal{S}}$. Indeed, both ends of $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$ are in $\tilde{\mathcal{S}}$. By the previous lemma there is a piece of the segment which is not in $\tilde{\mathcal{S}}$ (we notice that now we use the lemma for $k^{-\mu r_2} < \eta < \frac{1}{64}k^{-\frac{2r_2}{3}}$). Hence, the segment intersects one of the curves $\varkappa_2 = f_i(\varkappa_1)$ twice. It follows, that there is a point on the curve, where the curve is parallel to the segment. Suppose another segment $\vec{p}_{\mathbf{n}_2-\mathbf{n}'_2}$ intersects the same curve. Then, there is a point on the curve, where the curve is parallel to the second segment.

Since the curve is concave, it can not be the case. Therefore, $\vec{p}_{\mathbf{n}_2-\mathbf{n}'_2}$ intersects another curve. It follows that no more than two segments $\vec{p}_{\mathbf{n}_1-\mathbf{n}'_1}$, $\vec{p}_{\mathbf{n}_2-\mathbf{n}'_2}$ can be parallel to each other, if they are in the same elementary component $\tilde{S}$.

To finish the proof of the lemma we consider two cases. Suppose q in the inequality (252) satisfies the estimate $q > k^{2r_2/3}$. Then, by Lemma 5.8, the number of vectors $\vec{p}_{\mathbf{n}}$, $||\vec{p}_{\mathbf{n}}|| < 2k^{r_2}$, satisfying the inequality $p_{\mathbf{n}} < \frac{1}{64}k^{-2r_2/3}$ does not exceed $2^{12}k^{2r_2/3}$. Since each of them can be used only twice, the total number of short segments does not exceed $2^{13}k^{2r_2/3}$.

Let $q \leq k^{2r_2/3}$. If $|\epsilon_q| > \frac{1}{64}q^{-1}k^{-r_2}$. Then, obviously, $\frac{1}{64}k^{-2r_2/3} < |\epsilon_q|qk^{r_2/3}$. Applying Lemma 5.7, we obtain that the number of segments with the length less than $\frac{1}{64}k^{-2r_2/3}$ is less than $k^{2r_2/3}$. Since each of them can be used only twice, the total number of short segments does not exceed $2k^{2r_2/3}$. It remains to consider the case $q \leq k^{2r_2/3}$, $|\epsilon_q| \leq \frac{1}{64}q^{-1}k^{-r_2}$. By Lemma 5.6, clusters are well separated. Considering that the distance between clusters is greater than $\frac{1}{2q}$ and the size of each cluster is less than $\frac{1}{8q}$, we obtain that no more than $8q$ clusters can intersect $\tilde{S}$. Indeed, the distance between two clusters is greater than $\frac{1}{2q}$. By Corollary 5.12, the set $\tilde{S}$ belongs to the $k^{-\xi/30}J_0$ -neighborhood of each curve $\varkappa_2 = f_i(\varkappa_1)$, $i = 1, 2$. Using the hypothesis of the lemma we easily get that the size of the neighborhood is $o(q^{-1})$. If a cluster intersects $\tilde{S}$, its $\frac{1}{4q}$ -neighborhood intersects both curves $\varkappa_2 = f_i(\varkappa_1)$, $i = 1, 2$. Since the distance between clusters is greater than $\frac{1}{2q}$, the distance along the curve between its intersection with $\frac{1}{4q}$ -neighborhoods of different clusters is greater than $\frac{1}{4q}$. Considering that the lengths of the curves is less than 1, we obtain that no more than $8q$ clusters can intersect $\tilde{S}$. Next, the segments with the length less than $\frac{1}{2}k^{-2r_2/3}$ cannot connect different clusters, since the distance between clusters is greater than $\frac{1}{2q} \geq \frac{1}{2}k^{-2r_2/3}$. Therefore, any segment of the length less than $\frac{1}{2}k^{-2r_2/3}$ is inside one cluster. The part of the shorter curve inside the clusters has the length L_{in} which is less than the double size of a cluster $10|\epsilon_q|k^{r_2}$ (the curve is concave) multiplied by the number of clusters $8q$, i.e., $L_{in} < 80|\epsilon_q|qk^{r_2}$. If we consider the segments with the length greater than $|\epsilon_q|qk^{r_2/3}$, then the number of such segments is less than $L_{in}/|\epsilon_q|qk^{r_2/3}$, i.e., it is less than $80k^{2r_2/3}$. By Lemma 5.7, the total number of segments of the length less than $|\epsilon_q|qk^{r_2/3}$ is less than $k^{2r_2/3}$. Each of them can be used only twice. Thus, the total number of segments is less than $162k^{2r_2/3}$.

We proved that the number of segments in $\tilde{S}$ does not exceed $2^{14}k^{2r_2/3}$. Therefore, the number of points in $\tilde{S} \cap \mathcal{N}(k, r_2, \vec{\varkappa}_0, J_0)$ does not exceed $2^{14}k^{\frac{2}{3}r_2} + 1$. Considering that $k^{\gamma l r_1} > 2^{15}$, we obtain that the number of points inside $\tilde{S} \cap \mathcal{N}(k, r_2, \vec{\varkappa}_0, J_0)$ does not exceed $k^{\frac{2}{3}r_2 + \gamma l r_1}$.

If we show that $\mathcal{S}^{(2)}(k, \xi)$ is the union of no more than $k^{42l\gamma r_1}$ elementary components $\tilde{S}$, then estimate (261) easily follows. Indeed, let us consider the boundary of $\mathcal{S}^{(2)}(k, \xi)$. It is described by curves $D(\vec{\varkappa}, k^{2l} \pm k^{-\xi}) = 0$, $\vec{\varkappa} \in \mathbb{R}^2$. We break each curve into elementary components as described in Appendix 6. By Lemma 10.9 the number of such pieces is less than $k^{17l\gamma r_1}$. With each elementary piece of the boundary we associate the part of

the adjacent connected component of $\mathcal{S}^{(2)}(k, \xi)$, which is in the $k^{-\xi/30J_0}$ -neighborhood of the elementary piece. By Corollary 5.12, every point in $\mathcal{S}^{(2)}(k, \xi)$ belongs to such a component, some components overlapping. Let us consider one of these components $\hat{\mathcal{S}}$. By construction, it is adjacent to a boundary elementary component, which can be described in the form $\varkappa_1 = f_1(\varkappa_2)$ or $\varkappa_2 = f_1(\varkappa_1)$. Let us assume for definiteness that it is described by the formula $\varkappa_2 = f_1(\varkappa_1)$. By Corollary 5.12, there is another boundary (described by $\varkappa_2 = f_2(\varkappa_1)$) of $\hat{\mathcal{S}}$ in the $k^{-\xi/30J_0}$ -neighborhood of $\varkappa_2 = f_1(\varkappa_1)$. It also can be split into no more than $k^{17l\gamma r_1}$ elementary components. Further, each elementary component contains no more than $k^{8l\gamma r_1}$ points $\vec{z} : D(\vec{z}, k^{2l} + k^{-\xi}) = D(\vec{z}, k^{2l} - k^{-\xi}) = 0$, unless the last equality is an identity on this component (Bezout Theorem). We use these points to break each elementary component into at most $k^{8l\gamma r_1}$ parts. Correspondingly, we split the set $\hat{\mathcal{S}}$ by lines $\varkappa_1 = C$ into at most $k^{25l\gamma r_1}$ components $\tilde{\mathcal{S}}$. The second boundary of $\tilde{\mathcal{S}}$ also can be parameterized by $\varkappa_2$, since $D_{\varkappa_2} \neq 0$ on an elementary component of the boundary. By the definition of an elementary component of the boundary (Appendix 6), both functions $\varkappa_2 = f_i(\varkappa_1)$ are monotone, continuously differentiable and don't have inflection points, the length of the corresponding curves being less than 1. Moreover, neither boundary contains intersections with other pieces of the boundary of $\mathcal{S}^{(2)}(k, \xi)$. Thus, $\mathcal{S}^{(2)}(k, \xi)$ is the union of at most $k^{42l\gamma r_1}$ elementary components $\tilde{\mathcal{S}}$, each being bounded by lines $\varkappa_i = a, b$ and elementary pieces of the boundary of $\mathcal{S}^{(2)}(k, \xi)$, which do not intersect with other pieces of the boundary of $\mathcal{S}^{(2)}(k, \xi)$. $\square$

5.3.2 Model Operator for Step IV

Let $r_3 > r_2$. We repeat for r_3 the construction from the subsection 4.4.1, which was done for arbitrary $r_2 > r_1$. Let $\mathbf{m} \in \Omega(r_3)$. We denote the $k^{\gamma r_1}$ -component containing $\mathbf{m}$ by $\tilde{\Pi}(\mathbf{m})$ and the corresponding projector by $\tilde{P}(\mathbf{m})$. For $\mathbf{m}$ belonging to the same $k^{\gamma r_1}$ -component, $\tilde{\Pi}(\mathbf{m})$ and $\tilde{P}(\mathbf{m})$ are the same. Put

$$\mathcal{M}^{(3)} := \mathcal{M}^{(3)}(\varphi_0, r_3) = \{\mathbf{m} \in \mathcal{M}^{(2)}(\varphi_0, r_3) \cup \Omega_s^{(2)}(r_3) : \varphi_0 \in \mathcal{O}_{\mathbf{m}}^{(3)}(r'_2, 1)\}, \quad (262)$$

where $\Omega_s^{(2)}(r_3)$ is the extension of $\Omega_s^{(2)}(r_2)$ to $\Omega(r_3)$,

$$\Omega_s^{(2)}(r_3) = \{\mathbf{m} \in \Omega(r_3), 0 < p_{\mathbf{m}} \leq k^{-5r'_1}\}, \quad (263)$$

$\mathcal{O}_{\mathbf{m}}^{(3)}(r'_2, \tau)$ is the union of the disks of the radius $\tau k^{-r'_2}$ with the centers at poles of the resolvent $(\tilde{P}(\mathbf{m})(H(\vec{z}^{(2)}(\varphi)) - k^{2l}I)\tilde{P}(\mathbf{m}))^{-1}$ in the $k^{-44r'_1-2l-\delta}$ -neighborhood of φ_0 . (Here $\mathcal{M}^{(2)}(\varphi_0, r_3)$ is defined as in (162) with r_3 instead of r_2). For $\mathbf{m}$ belonging to the same $k^{\gamma r_1}$ -component, the sets $\mathcal{O}_{\mathbf{m}}^{(3)}(r'_2, \tau)$ are identical. We say that $\mathbf{m} \in \mathcal{M}^{(3)}$ is $k^{\gamma r_1}$ -resonant. The corresponding $k^{\gamma r_1}$ -clusters we call resonant too.

Let $\varphi_0 \in \omega^{(3)}(k, \delta, 1)$. By construction of the non-resonant set $\omega^{(3)}(k, \delta, 1)$, we have $\mathcal{M}^{(3)} \cap \Omega(r_2) = \emptyset$.

Further we use the property of the set $\mathcal{M}^{(3)}$ formulated in the next lemma which is an analogue of the Lemma 4.11.

Lemma 5.14. *Let $r'_2 > 2k^{(\gamma+\delta_0)10^{-4}r_1-2\delta}$.⁴ Let $1/20 < \gamma' < 20$, $\mathbf{m}_0 \in \Omega(r_3)$ and $\Pi_{\mathbf{m}_0}$ be the $k^{\gamma'r_2}$ -neighborhood (in $\|\cdot\|$ -norm) of $\mathbf{m}_0$. Then the set $\Pi_{\mathbf{m}_0}$ contains less than $k^{\frac{2}{3}\gamma'r_2+50l\gamma r_1}$ elements of $\mathcal{M}^{(3)}$.*

Proof. If $\mathbf{m} \in \mathcal{M}^{(3)}$, then there is a φ_* such that $|\varphi_* - \varphi_0| < k^{-r'_2}$ and

$$\det \left(\tilde{P}(\mathbf{m})(H(\tilde{\mathcal{Z}}^{(2)}(\varphi_*)) - k^{2l}I)\tilde{P}(\mathbf{m}) \right) = 0,$$

where $\tilde{P}(\mathbf{m})$ is the projection corresponding to the $k^{\gamma r_1}$ -cluster $\tilde{\Pi}(\mathbf{m})$, which includes $\mathbf{m}$. The cluster $\tilde{\Pi}(\mathbf{m})$ can be simple, white, grey or black. Since φ_0 is close to φ_* , perturbation arguments give:

$$\left\| (\tilde{P}(\mathbf{m})(H(\tilde{\mathcal{Z}}^{(2)}(\varphi_0)) - k^{2l}I)\tilde{P}(\mathbf{m}))^{-1} \right\| \geq \frac{1}{4l}k^\xi, \quad \xi \geq r'_2 - 2l + 1. \quad (264)$$

We will apply Lemma 5.13 to $\tilde{\Pi}(\mathbf{m})$ with $\xi = r'_2 - 2l + 1$ in order to prove the lemma in hand in the same way we proved Lemma 4.11, using Lemmas 4.8, 4.10. There are some technical complications though. Here is a detailed proof.

We start with considering simple boxes $\tilde{\Pi}(\mathbf{m})$, $\mathbf{m} \in \mathcal{M}^{(3)} \cap \Omega_s(r_3)$. Each box has the $\|\cdot\|$ -size $2k^{r_1/2}$ and contains no other than $\mathbf{m}$ elements of $\mathcal{M}^{(2)}(\varphi_0, r_3) \cup \Omega_s(r_3)$. Indeed, $\tilde{\mathcal{Z}}^{(2)}(\varphi_0)$ satisfies the conditions of Lemma 3.18. This means that the k^δ -cluster around each $\mathbf{q}$: $\|\vec{p}_{\mathbf{q}}\| < k^{r_1}$ is non-resonant. Since $\tilde{\mathcal{Z}} = \tilde{\mathcal{Z}}^{(2)}(\varphi_0) + \vec{p}_{\mathbf{m}}$ is a small perturbation of $\tilde{\mathcal{Z}}^{(2)}(\varphi_0)$, the k^δ -box around each $\mathbf{m} + \mathbf{q}$: $\|\vec{p}_{\mathbf{q}}\| < k^{r_1}$ is non-resonant too. This means $\mathbf{m} + \mathbf{q} \notin \mathcal{M}^{(2)}$. Further, $\mathbf{m} + \mathbf{q} \notin \Omega_s(r_3)$ by (28), since $\mathbf{m} \in \Omega_s(r_3)$ and $\|\vec{p}_{\mathbf{q}}\| < k^{r_1}$. Thus, $\mathbf{m} + \mathbf{q} \notin \mathcal{M}^{(2)}(\varphi_0, r_3) \cup \Omega_s(r_3)$. Next, we apply Lemma 5.13 with $\tilde{\mathcal{Z}}_0 = \tilde{\mathcal{Z}}^{(2)}(\varphi_0) + \vec{p}_{\mathbf{m}_0}$, $J_0 = 1$, $\xi = k^{r'_2-2l+1}$, to conclude that the number of simple boxes $\tilde{\Pi}(\mathbf{m})$, $\mathbf{m} \in \mathcal{M}^{(3)} \cap \Omega_s(r_3)$ does not exceed $k^{\frac{2}{3}\gamma'r_2+43l\gamma r_1}$. Indeed, we rewrite $\tilde{\mathcal{Z}}^{(2)}(\varphi_0) + \vec{p}_{\mathbf{m}}$ in the form: $\tilde{\mathcal{Z}}^{(2)}(\varphi_0) + \vec{p}_{\mathbf{m}} = \tilde{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}}$, $\mathbf{n} = \mathbf{m} - \mathbf{m}_0 \in \Omega(\gamma'r_2)$. By (251), $\tilde{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}} \in \mathcal{S}^{(2)}(k, \xi)$ (the operator in formula (251) having the size $2k^{\frac{1}{2}r_1}$ and $\xi = r'_2 - 2l + 1$, see (264)). Since, $\tilde{\Pi}(\mathbf{m})$ is simple, $\tilde{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}} \in \mathcal{N}(k, \gamma'r_2, \tilde{\mathcal{Z}}_0, 1)$ (here, γ is taken to be equal to $1/2$ in the definition of $\mathcal{S}^{(2)}(k, \xi)$). Thus, $\tilde{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}} \in \mathcal{S}^{(2)}(k, \xi) \cap \mathcal{N}(k, \gamma'r_2, \tilde{\mathcal{Z}}_0, 1)$. By Lemma 5.13, the number of such $\tilde{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}}$ does not exceed $k^{2\gamma'r_2/3+43l\frac{1}{2}r_1}$. Therefore, the number of $\mathcal{M}^{(3)}$ elements in simple boxes also does not exceed $k^{2\gamma'r_2/3+43l\frac{1}{2}r_1}$.

Next, let us consider white clusters $\tilde{\Pi}(\mathbf{m})$, such that $\xi \geq k^{\frac{1}{6}\gamma r_1-2\delta}$. Generally speaking, $\tilde{\Pi}(\mathbf{m})$ has a shape (in $\mathbb{Z}^4$) more complicated than a rectangular. However, each such cluster can be put in a box of the size $3k^{\gamma r_1/2+2\delta_0}$, the box containing less than $k^{\frac{1}{6}\gamma r_1-\delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$ and the box resolvent satisfying (264) with $\xi = k^{\frac{1}{6}\gamma r_1-2\delta}$ (Lemma 10.10). Applying Lemma 5.13 to such boxes ($\tilde{\mathcal{Z}}_0 = \tilde{\mathcal{Z}}^{(2)}(\varphi_0) + \vec{p}_{\mathbf{m}_0}$, $J_0 = k^{\frac{1}{6}\gamma r_1-\delta_0 r_1}$, $\xi = k^{\frac{1}{6}\gamma r_1-2\delta}$), we obtain that the number of $\Pi_{\mathbf{m}_0}$ points $\mathbf{m}$ in such boxes does not exceed $k^{2\gamma'r_2/3+43l\gamma r_1}$. Similarly, we can treat grey boxes when $\xi \geq k^{\frac{1}{2}\gamma r_1+2\delta_0 r_1-2\delta}$ (Lemma 10.11), black boxes

⁴ We also notice that this condition is consistent with the restriction $r'_2 < \frac{\beta}{128}k^{\delta_0 r_1-\delta-3}$ in (187).

when $\xi \geq k^{\gamma r_1 + \delta_0 r_1 - 2\delta}$ (Lemma 10.12). However, in some cases ξ does not satisfy the previous estimates from below. For such ξ a somewhat more complicated construction is needed. Indeed, let us consider $(\tilde{P}(H(\vec{\mathcal{Z}}^{(2)}(\varphi_0)) - k^{2l}I)\tilde{P})^{-1}$ for $\tilde{\Pi}$ being white, grey or black cluster containing a point(s) of $\mathcal{M}^{(3)}$. A cluster $\tilde{\Pi}$ consists of blocks with the minimal size $k^{\gamma r_1/6}$. Let us create a substructure inside $\tilde{\Pi}$. Namely, we construct white, grey and black clusters corresponding to a smaller γ which we denote by $\tilde{\gamma}$, $\tilde{\gamma} = 10^{-4}\gamma$. Note, that there are no simple small clusters inside $\tilde{\Pi}$, since $\tilde{\Pi}$ is not simple. The size of these new clusters is much smaller than $k^{\gamma r_1/6}$. However, they have properties analogous to those of the bigger clusters (γ). These new clusters we call subclusters. We assert that at least one subcluster satisfies one of the following estimates (depending on whether this subcluster is white, grey or black):

$$\|(P_{w,sub}(H(\vec{\mathcal{Z}}^{(2)}(\varphi_0)) - k^{2l}I)P_{w,sub})^{-1}\| > k^{k^{\frac{\tilde{\gamma}r_1}{6}-2\delta}}, \quad (265)$$

$$\|(P_{g,sub}(H(\vec{\mathcal{Z}}^{(2)}(\varphi_0)) - k^{2l}I)P_{g,sub})^{-1}\| > k^{k^{(\frac{\tilde{\gamma}}{2}+2\tilde{\delta}_0)r_1-2\delta}}, \quad (266)$$

$$\|(P_{b,sub}(H(\vec{\mathcal{Z}}^{(2)}(\varphi_0)) - k^{2l}I)P_{b,sub})^{-1}\| > k^{k^{(\tilde{\gamma}+\tilde{\delta}_0)r_1-2\delta}}, \quad (267)$$

where $\tilde{\delta}_0 = \tilde{\gamma}/100$ (cf. definition of δ_0). Indeed, if all subclusters satisfy the inequalities opposite to the inequalities above, then the perturbation series for the resolvent of the bigger cluster (γ) (with respect to the block operator consisting of subclusters) converges, see the proof of Theorem 5.1, in particular the proof of (206) – (209). Hence, we have

$$\|(\tilde{P}(H(\vec{\mathcal{Z}}^{(2)}(\varphi_0)) - k^{2l}I)\tilde{P})^{-1}\| \leq k^{k^{(\tilde{\gamma}+\tilde{\delta}_0)r_1-2\delta}},$$

which contradicts to (264) under the hypothesis of the lemma $r'_2 > 2k^{(\gamma+\delta_0)10^{-4}r_1-2\delta}$.

Now, let us consider a resonant $k^{\gamma r_1}$ -cluster $\tilde{\Pi}$, see (264), and the substructure inside. Note that each subcluster satisfying (265)-(267) can be treated the same way we treated the bigger clusters for large ξ . Namely, let us consider all $k^{\gamma r_1}$ -clusters $\tilde{\Pi}$ for which there exists a white subcluster satisfying (265). By Lemma 10.11 each such subcluster can be put in a box of the size $3k^{\tilde{\gamma}r_1/2+2\tilde{\delta}_0}$, the box resolvent satisfying (265). Such box has less than $k^{(\frac{\tilde{\gamma}}{6}-\tilde{\delta}_0)r_1}$ points $\mathbf{m}$ of $\mathcal{M}^{(2)}$. Now, applying Lemma 5.13 with $\vec{\mathcal{Z}}_0 = \vec{\mathcal{Z}}^{(2)}(\varphi_0) + \vec{p}_{\mathbf{m}_0}$, $J_0 = k^{(\frac{\tilde{\gamma}}{6}-\tilde{\delta}_0)r_1}$, $\xi = k^{\frac{\tilde{\gamma}r_1}{6}-2\delta}$, we obtain that the number of points $\mathbf{m}$ in white subclusters (265) does not exceed $k^{\frac{2\gamma' r_2}{3}+43l\tilde{\gamma}r_1}$. Here we notice that condition of Lemma 5.13 holds, since $r_2 < k^{\tilde{\delta}_0 r_1 - 3\delta}$ by (186). It follows that the number of $k^{\gamma r_1}$ -clusters $\tilde{\Pi}(\mathbf{m})$, containing at least one white subcluster (265), does not exceed $k^{\frac{2\gamma' r_2}{3}+43l\tilde{\gamma}r_1}$.

Next, we consider all $k^{\gamma r_1}$ -clusters $\tilde{\Pi}(\mathbf{m})$ for which there exists a grey subcluster, satisfying (266), but no white subclusters satisfying (265). Applying Lemma 10.11 and

Lemma 5.13 with $J_0 = k^{(\frac{\tilde{\gamma}}{2} + \tilde{\delta}_0)r_1}$ and $\xi = k^{(\frac{\tilde{\gamma}}{2} + 2\tilde{\delta}_0)r_1 - 2\delta}$, we obtain that the number of such $k^{\gamma r_1}$ -clusters $\tilde{\Pi}(\mathbf{m})$ in $\Pi_{\mathbf{m}_0}$ does not exceed $k^{\frac{2\gamma' r_2}{3} + 43l\gamma r_1}$.

Similarly, applying Lemma 10.12 and Lemma 5.13 with $\xi = k^{(\tilde{\gamma} + \tilde{\delta}_0)r_1 - 2\delta}$ and $J_0 = ck^{\tilde{\gamma}r_1 + 3}$, we obtain that the number of $k^{\gamma r_1}$ -clusters $\tilde{\Pi}(\mathbf{m})$, containing a black subcluster (267) (and no grey or white subclusters, satisfying (265), (266)), does not exceed $k^{\frac{2\gamma' r_2}{3} + 43l\gamma r_1}$. Here, we also used $r_2 < k^{\tilde{\delta}_0 r_1 - 3 - 3\delta}$.

Combining these estimates, we see that the number of clusters $\tilde{\Pi}$, containing at least one point of $\mathcal{M}^{(3)}$ does not exceed $k^{\frac{2\gamma' r_2}{3} + 43l\gamma r_1}$. Taking into account that each $k^{\gamma r_1}$ -cluster has a size not greater than $k^{\frac{3\gamma r_1}{2} + 3}$ and, hence, contains less than $k^{6\gamma r_1 + 12}$ elements, we obtain that the total number of elements of $\mathcal{M}^{(3)}$ in $\Pi_{\mathbf{m}_0}$, does not exceed $k^{\frac{2\gamma' r_2}{3} + 50l\gamma r_1}$.

□

We continue with constructing $k^{\gamma r_1}$ -clusters in $\Omega(r_3)$, $r_3 > r_2$, the same way we did it for $\Omega(r_2)$ in Section 4.4.1. We call a $k^{\gamma r_1}$ -cluster resonant if it contains $\mathbf{m} \in \mathcal{M}^{(3)}$, see (262). Next, we repeat the construction after Lemma 4.11. More precisely, let us split $\Omega(r_3) \setminus \Omega(r_2)$ into $k^{\gamma r_2}$ -boxes, $\gamma = \frac{1}{5}$.

1. *Simple region.* Let $\Omega_s^{(3)}(r_3) \subset \Omega_s^{(2)}(r_3)$ be defined by the formula:

$$\Omega_s^{(3)}(r_3) = \{\mathbf{m} \in \Omega(r_3), 0 < p_{\mathbf{m}} \leq k^{-r'_2 k^{2\gamma r_1}}\}. \quad (268)$$

It is easy to see that $\Omega_s^{(3)}(r_3) \subset \mathcal{M}(\varphi_0, r_3)$, since $p_{\mathbf{m}}$ is small, see (71), (69). Next, if $\mathbf{m} \in \Omega_s^{(3)}(r_3)$, then there are no other elements of $\mathcal{M}(\varphi_0, r_3)$ in the k^δ -box around $\mathbf{m}$. Further, $\mathbf{m}$ itself can belong or do not belong to $\mathcal{M}^{(2)}(\varphi_0, r_3)$, but there are no other elements of $\mathcal{M}^{(2)}(\varphi_0, r_3)$ in the k^{r_1} -box around such $\mathbf{m}$. The proof of these facts is analogous to that in Step III, see “Simple region”, page 47. Next, if $\mathbf{m} \in \Omega_s^{(3)}(r_3)$, then there are no other elements of $\Omega_s^{(3)}(r_3)$ in the surrounding $||| \cdot |||$ -box of the size k^{r_2} , see (28). Last, $\mathbf{m}$ can belong or do not belong to $\mathcal{M}^{(3)}(\varphi_0, r_3)$, but there are no other elements from $\mathcal{M}^{(3)}(\varphi_0, r_3)$ in the k^{r_2} -box around such $\mathbf{m}$. Indeed, $\vec{\mathcal{Z}}^{(2)}(\varphi_0)$ satisfies the conditions of Lemma 4.21. This means that the $k^{\gamma r_1}$ -cluster around each $\mathbf{q}$: $0 < |||\vec{p}_{\mathbf{q}}||| < k^{r_2}$ is non-resonant. Since $\vec{\mathcal{Z}} = \vec{\mathcal{Z}}^{(2)}(\varphi_0) + \vec{p}_{\mathbf{m}}$ is a small perturbation of $\vec{\mathcal{Z}}^{(2)}(\varphi_0)$, the $k^{\gamma r_1}$ -box around each $\mathbf{m} + \mathbf{q}$: $0 < |||\vec{p}_{\mathbf{q}}||| < k^{r_2}$ is non-resonant too. This means $\mathbf{m} + \mathbf{q} \notin \mathcal{M}^{(3)}(\varphi_0, r_3)$.

For each $\mathbf{m} \in \Omega_s^{(3)}(r_3)$ we consider its $k^{r_2/2}$ -neighborhood. The union of such boxes we call the simple region and denote it by $\Pi_s(r_3)$. The corresponding projection is $P_s(r_3)$. Note that the distance from the simple region to the nearest point of $\mathcal{M}^{(3)}$ is greater than $\frac{1}{2}k^{r_2}$.

2. *Black, grey and white regions* are defined in the same way as in the construction after Lemma 4.11 with r_3 instead of r_2 , r_2 instead of r_1 , $\mathcal{M}^{(3)}$ instead of $\mathcal{M}^{(2)}$ and the restriction $p_{\mathbf{m}} > k^{-r'_2 k^{2\gamma r_1}}$ instead of $p_{\mathbf{m}} > k^{-5r'_1}$. We continue to use notation

$P_b, P_g, P'_g, P_w, P'_w$ and $\Pi_b, \Pi_g, \Pi'_g, \Pi_w, \Pi'_w$. Sometimes, where it can lead to confusion we will write $P_b(r_3)$ etc. to distinguish these objects from the ones introduced in Step II.

3. *Non-resonant region.* Now, the non-resonant region consists of two parts: the simpler part which was non-resonant already in the previous step and the part which is new for the current step. Namely, first we consider k^δ -neighborhoods of all points in the set $\mathcal{M}(r_3, \varphi_0) \setminus \left(\mathcal{M}(r_2, \varphi_0) \cup \mathcal{M}^{(2)}(r_3, \varphi_0) \cup \Omega_s^{(2)}(r_3) \right)$. The union of this neighborhoods we denote $\Pi_{nr, \delta}$. The corresponding projection is $P_{nr, \delta}$. These k^δ -clusters can be treated by means of the second step. We also consider all points in the set $\mathcal{M}^{(2)}(r_3, \varphi_0) \cup \Omega_s^{(2)}(r_3) \setminus \left(\mathcal{M}^{(2)}(r_2, \varphi_0) \cup \mathcal{M}^{(3)}(r_3, \varphi_0) \cup \Omega_s^{(3)}(r_3) \right)$. We construct simple, white, grey and black clusters around them exactly as in preparation to Step III. The union of these clusters we denote Π_{nr, r_1} . The corresponding projection is P_{nr, r_1} . The set $\Pi_{nr} := \Pi_{nr, \delta} \cup \Pi_{nr, r_1}$ is called the non-resonant set with P_{nr} being the corresponding projection. The part of the non-resonant region which is outside $\Pi_s \cup \Pi_b \cup \Pi_g \cup \Pi_w$, we denote Π'_{nr} and the corresponding projection by P'_{nr} .

We put as before

$$P_r(r_3) := P_s(r_3) + P_b(r_3) + P'_g(r_3) + P'_w(r_3), \quad P^{(3)} := P_r(r_3) + P'_{nr}(r_3) + P(r_2). \quad (269)$$

We also continue to use the similar agreement in the notation which we set in Step II. We just note that now we use r_2 rather than r_1 to establish equivalence between the boxes.

We continue construction from Step II. Repeating the arguments from the proofs of Lemmas 4.12, 4.13, 4.14 with obvious changes (in particular, using Lemma 5.14 instead of Lemma 4.11) we obtain the following results.

- Lemma 5.15.** 1. Each Π_b^j contains no more than $k^{\gamma r_2/2 - \delta_0 r_2 + 150l\gamma r_1}$ black boxes.
2. The size of Π_b^j in $||| \cdot |||$ norm is less than $k^{3\gamma r_2/2 + 150l\gamma r_1}$.
3. Each Π_b^j contains no more than $k^{\gamma r_2 + 150l\gamma r_1}$ elements of $\mathcal{M}^{(3)}$. Moreover, any box of $||| \cdot |||$ -size $k^{3\gamma r_2/2 + 150l\gamma r_1}$ containing Π_b^j has no more than $k^{\gamma r_2 + 150l\gamma r_1}$ elements of $\mathcal{M}^{(3)}$ inside.

- Lemma 5.16.** 1. Each Π_g^j contains no more than $k^{\gamma r_2/3 + 2\delta_0 r_2}$ grey boxes.

2. The size of Π_g^j in $||| \cdot |||$ norm is less than $k^{5\gamma r_2/6 + 4\delta_0 r_2}$.
3. Each Π_g^j contains no more than $k^{\gamma r_2/2 + \delta_0 r_2}$ elements of $\mathcal{M}^{(3)}$.

- Lemma 5.17.** 1. The size of Π_w^j in $||| \cdot |||$ norm is less than $k^{\gamma r_2/3 - \delta_0 r_2}$.

2. Each Π_w^j contains no more than $k^{\gamma r_2/6 - \delta_0 r_2}$ points of $\mathcal{M}^{(3)}$.

The construction of the rest of Section 4.4.1 stays unchanged. Let us introduce corresponding notation, formulate the results and provide some comments.

Next lemmas are the analogues of Lemmas 4.17, 4.18, 4.19.

Lemma 5.18. *Let $\varphi_0 \in \omega^{(3)}(k, \delta, \tau)$, $|\varphi - \varphi_0| < k^{-k^{r_1}}$. Then,*

$$\left\| \left(P_{nr} (H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l} I) P_{nr} \right)^{-1} \right\| < k^{r'_2 k^{2\gamma r_1}} k^{r'_2} \leq k^{k^{3\gamma r_1}}. \quad (270)$$

Proof. Construction in Section 4.4 is made for an arbitrary large r_2 . Here we repeat it for r_3 . We use Lemma 4.17 for $\Pi_{nr, \delta}$ Lemma 4.18 for white, grey and black clusters ($\varepsilon_0 = k^{-r'_2}$). We also use Lemma 4.19 ($p_{\mathbf{m}} > k^{-r'_2 k^{2\gamma r_1}}$, $\varepsilon_0 = k^{-r'_2}$), for simple clusters in Π_{nr, r_1} . We also use (249). All together the estimates for the clusters resolvents yield (270). The estimate (270) is stable when $|\varphi - \varphi_0| < k^{-k^{r_1}}$, since $k^{-k^{r_1} + 2l} = o(k^{-k^{3\gamma r_1}})$. $\square$

Lemma 5.19. *Let $\varphi_0 \in \omega^{(3)}(k, \delta, \tau)$, and $|\varphi - \varphi_0| < k^{-k^{r_1}}$, $i = 1, 2, 3$. Then,*

1. *The number of poles of the resolvent $\left(P_i (H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l} I) P_i \right)^{-1}$ in the disc $|\varphi - \varphi_0| < k^{-k^{r_1}}$ is no greater than $N_i^{(2)}$, where $N_1^{(2)} = k^{\gamma r_2 + 150l\gamma r_1}$, $N_2^{(2)} = k^{\gamma r_2/2 + \delta_0 r_2}$, $N_3^{(2)} = k^{\gamma r_2/6 - \delta_0 r_2}$.*
2. *Let ε be the distance to the nearest pole of the resolvent in $\mathcal{W}^{(3)}$ and $\varepsilon_0 = \min\{\varepsilon, k^{-r'_2}\}$. Then the following estimates hold:*

$$\left\| \left(P_i (H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l} I) P_i \right)^{-1} \right\| < k^{2r'_2 k^{2\gamma r_1}} k^{r'_2} \left(\frac{k^{-r'_2}}{\varepsilon_0} \right)^{N_i^{(2)}} \leq k^{k^{3\gamma r_1}} \left(\frac{k^{-r'_2}}{\varepsilon_0} \right)^{N_i^{(2)}}, \quad (271)$$

$$\left\| \left(P_i (H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l} I) P_i \right)^{-1} \right\|_1 < k^{2r'_2 k^{2\gamma r_1}} k^{r'_2 + 8\gamma r_2} \left(\frac{k^{-r'_2}}{\varepsilon_0} \right)^{N_i^{(2)}} \leq k^{k^{3\gamma r_1}} \left(\frac{k^{-r'_2}}{\varepsilon_0} \right)^{N_i^{(2)}}. \quad (272)$$

Proof. The proof of this lemma is analogous to that of Lemma 4.18 up to the replacement of $\mathcal{M}^{(2)}$ by $\mathcal{M}^{(3)}$, $\mathcal{O}_{\mathbf{m}}^{(2)}$ by $\mathcal{O}_{\mathbf{m}}^{(3)}$, and the shift of indices: δ to r_1 , r_1 to r_2 , etc. We apply Lemmas 5.15–5.17 instead of 4.12–4.14 and Lemmas 4.18, 4.19 with $\varepsilon_0 = k^{-r'_2}$ and $p_{\mathbf{m}} > k^{-r'_2 k^{2\gamma r_1}}$ instead of Lemma 3.14. We also note that $N_i^{(1)} < k^{2\gamma r_1}$ in (174), (175). $\square$

Lemma 5.20. *Let $\varphi_0 \in \omega^{(3)}(k, \delta, \tau)$. Then, the operator $\left(P_s^j(H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l}I)P_s^j\right)^{-1}$ has no more than one pole in the disk $|\varphi - \varphi_0| < k^{-k^{r_1}}$. Moreover,*

$$\left\| \left(P_s^j(H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l}I)P_s^j \right)^{-1} \right\| < \frac{8k^{-2l+1}}{p_{\mathbf{m}}\varepsilon_0}, \quad (273)$$

$$\left\| \left(P_s^j(H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l}I)P_s^j \right)^{-1} \right\|_1 < \frac{8k^{-2l+1+4r_2}}{p_{\mathbf{m}}\varepsilon_0}, \quad (274)$$

$\varepsilon_0 = \min\{\varepsilon, k^{-r'_2}\}$, where ε is the distance to the pole of the operator.

Note that $p_{\mathbf{m}} > k^{-\mu r_3}$ when $\mathbf{m} \in \Omega(r_3)$. The analogues of Lemma 4.15 and Corollary 4.16 also hold.

5.3.3 Resonant and Nonresonant Sets for Step IV

We divide $[0, 2\pi)$ into $[2\pi k^{k^{r_1}}] + 1$ intervals $\Delta_m^{(3)}$ with the length not bigger than $k^{-k^{r_1}}$. If a particular interval belongs to $\mathcal{O}^{(3)}$ we ignore it; otherwise, let $\varphi_0(m) \notin \mathcal{O}^{(3)}$ be a point inside the $\Delta_m^{(3)}$. Let

$$\mathcal{W}_m^{(3)} = \{\varphi \in \mathcal{W}^{(3)} : |\varphi - \varphi_0(m)| < 4k^{-k^{r_1}}\}. \quad (275)$$

Clearly, neighboring sets $\mathcal{W}_m^{(3)}$ overlap (because of the multiplier 4 in the inequality), they cover $\hat{\mathcal{W}}^{(3)}$, which is the restriction of $\mathcal{W}^{(3)}$ to the $2k^{-k^{r_1}}$ -neighborhood of $[0, 2\pi)$. For each $\varphi \in \hat{\mathcal{W}}^{(3)}$ there is an m such that $|\varphi - \varphi_0(m)| < 4k^{-k^{r_1}}$. We consider the poles of the resolvent $(P^{(3)}(H(\vec{\mathcal{Z}}^{(3)}(\varphi)) - k^{2l}I)P^{(3)})^{-1}$ in $\hat{\mathcal{W}}_m^{(3)}$ and denote them by $\varphi_{mj}^{(3)}$, $j = 1, \dots, M_m$. As before, the resolvent has a block structure. The number of blocks clearly cannot exceed the number of elements in $\Omega(r_3)$, i.e. k^{4r_3} . Using the estimates for the number of poles for each block, the estimate being provided by Lemma 5.19, Part 1, we can roughly estimate the number of poles of the resolvent by $k^{4r_3+r_2}$. Next, let $r'_3 > k^{r_1}$ and $\mathcal{O}_{mj}^{(4)}$ be the disc of the radius $k^{-r'_3}$ around $\varphi_{mj}^{(3)}$.

Definition 5.21. *The set*

$$\mathcal{O}^{(4)} = \cup_{mj} \mathcal{O}_{mj}^{(4)} \quad (276)$$

we call the forth resonant set. The set

$$\mathcal{W}^{(4)} = \hat{\mathcal{W}}^{(3)} \setminus \mathcal{O}^{(4)} \quad (277)$$

is called the forth non-resonant set. The set

$$\omega^{(4)} = \mathcal{W}^{(4)} \cap [0, 2\pi) \quad (278)$$

is called the forth real non-resonant set.

The following statements can be proven in the same way as Lemmas 4.21, 4.22 and 4.23.

Lemma 5.22. *Let $r'_3 > \mu r_3 > k^{r_1}$, $\varphi \in \mathcal{W}^{(4)}$, $\varphi_0(m)$ corresponds to an interval $\Delta_m^{(3)}$ containing $\Re\varphi$. Let Π be one of the components $\Pi_s^j(\varphi_0(m))$, $\Pi_b^j(\varphi_0(m))$, $\Pi_g^j(\varphi_0(m))$, $\Pi_w^j(\varphi_0(m))$ and $P(\Pi)$ be the projection corresponding to Π . Let also $\varkappa \in \mathbb{C} : |\varkappa - \varkappa^{(3)}(\varphi)| < k^{-r'_3 k^{2\gamma r_2}}$. Then,*

$$\left\| (P(\Pi) (H(\varkappa(\varphi)) - k^{2l} I) P(\Pi))^{-1} \right\| < k^{2\mu r_3 + r'_3 N^{(2)}}, \quad (279)$$

$$\left\| (P(\Pi) (H(\varkappa(\varphi)) - k^{2l} I) P(\Pi))^{-1} \right\|_1 < k^{(2\mu+1)r_3 + r'_3 N^{(2)}}, \quad (280)$$

$N^{(2)}$ corresponding to the color of Π ($N^{(2)} = 1$, $k^{\gamma r_2 + 150l\gamma r_1}$, $k^{\gamma r_2/2 + \delta_0 r_2}$, $k^{\gamma r_2/6 - \delta_0 r_2}$ for simple, black, grey and white clusters, correspondingly).

By total size of the set $\mathcal{O}^{(4)}$ we mean the sum of the sizes of its connected components.

Lemma 5.23. *Let $r'_3 \geq (\mu + 10)r_3$, $r_3 > k^{r_1}$. Then, the size of each connected component of $\mathcal{O}^{(4)}$ is less than $32k^{4r_3 - r'_3}$. The total size of $\mathcal{O}^{(4)}$ is less than $k^{-r'_3/2}$.*

Lemma 5.24. *Let $\varphi \in \mathcal{W}^{(3)}$ and C_4 be the circle $|z - k^{2l}| = k^{-2r'_3 k^{2\gamma r_2}}$. Then*

$$\left\| (P(r_2)(H(\varkappa^{(3)}(\varphi)) - z)P(r_2))^{-1} \right\| \leq 4^3 k^{2r'_3 k^{2\gamma r_2}}.$$

We prove this lemma using (202).

6 STEP IV

6.1 Operator $H^{(4)}$. Perturbation Formulas

Let $P(r_3)$ be an orthogonal projector onto $\Omega(r_3) := \{\mathbf{m} : \|\vec{p}_{\mathbf{m}}\| \leq k^{r_3}\}$ and $H^{(4)} = P(r_3)HP(r_3)$. From now on, we assume

$$k^{r_1} < r_3 < k^{\gamma 10^{-7} r_2}, \quad k^{2\gamma 10^{-4} r_2} < r'_3 < k^{\delta_0 r_2/2}. \quad (281)$$

We consider $H^{(4)}(\varkappa^{(3)}(\varphi))$ as a perturbation of $\tilde{H}^{(3)}(\varkappa^{(3)}(\varphi))$:

$$\tilde{H}^{(3)} := \tilde{P}_j^{(3)} H \tilde{P}_j^{(3)} + \left(P(r_3) - \tilde{P}_j^{(3)} \right) H_0,$$

where $H = H(\varkappa^{(3)}(\varphi))$, $H_0 = H_0(\varkappa^{(3)}(\varphi))$ and $\tilde{P}_j^{(3)}$ is the projection $P^{(3)}$ corresponding to $\varphi_0(j)$ in the interval $\Delta_j^{(3)}$ containing φ , see (269). Note that the operator $\tilde{H}^{(3)}$ has a

block structure, the block $\tilde{P}_j^{(3)} H \tilde{P}_j^{(3)}$ being composed of smaller blocks $P_i H P_i$, $i = 0, \dots, 5$. By analogy with (188)–(190),

$$W^{(3)} = H^{(4)} - \tilde{H}^{(3)} = P(r_3) V P(r_3) - \tilde{P}_j^{(3)} V \tilde{P}_j^{(3)}, \quad (282)$$

$$g_r^{(4)}(\vec{\kappa}) := \frac{(-1)^r}{2\pi i r} \text{Tr} \oint_{C_4} \left(W^{(3)} (\tilde{H}^{(3)}(\vec{\kappa}) - zI)^{-1} \right)^r dz, \quad (283)$$

$$G_r^{(4)}(\vec{\kappa}) := \frac{(-1)^{r+1}}{2\pi i} \oint_{C_4} (\tilde{H}^{(3)}(\vec{\kappa}) - zI)^{-1} \left(W^{(3)} (\tilde{H}^{(3)}(\vec{\kappa}) - zI)^{-1} \right)^r dz, \quad (284)$$

where C_4 is the circle $|z - k^{2l}| = \varepsilon_0^{(4)}$, $\varepsilon_0^{(4)} = k^{-2r'_3 k^{2\gamma r_2}}$.

The proof of the following statements is analogous to the one in the previous step (see Theorem 5.1, Corollary 5.2 and Lemma 5.3) up to the replacement of r_3 by r_4 , r_2 by r_3 , r_1 by r_2 , etc.

Theorem 6.1. *Suppose $k > k_*$, φ is in the real $k^{-r'_3 - \delta}$ -neighborhood of $\omega^{(4)}(k, \delta, \tau)$ and $\kappa \in \mathbb{R}$, $|\kappa - \kappa^{(3)}(\varphi)| \leq \varepsilon_0^{(4)} k^{-2l+1-\delta}$, $\vec{\kappa} = \kappa(\cos \varphi, \sin \varphi)$. Then, there exists a single eigenvalue of $H^{(4)}(\vec{\kappa})$ in the interval $\varepsilon_4(k, \delta, \tau) = \left(k^{2l} - \varepsilon_0^{(4)}, k^{2l} + \varepsilon_0^{(4)} \right)$. It is given by the absolutely converging series:*

$$\lambda^{(4)}(\vec{\kappa}) = \lambda^{(3)}(\vec{\kappa}) + \sum_{r=2}^{\infty} g_r^{(4)}(\vec{\kappa}). \quad (285)$$

For coefficients $g_r^{(4)}(\vec{\kappa})$ the following estimates hold:

$$|g_r^{(4)}(\vec{\kappa})| < k^{-\frac{\beta}{5} k^{r_2 - r_1 - \beta(r-1)}}. \quad (286)$$

The corresponding spectral projection is given by the series:

$$\mathcal{E}^{(4)}(\vec{\kappa}) = \mathcal{E}^{(3)}(\vec{\kappa}) + \sum_{r=1}^{\infty} G_r^{(4)}(\vec{\kappa}), \quad (287)$$

$\mathcal{E}^{(3)}(\vec{\kappa})$ being the spectral projection of $H^{(3)}(\vec{\kappa})$. The operators $G_r^{(4)}(\vec{\kappa})$ satisfy the estimates:

$$\|G_r^{(4)}(\vec{\kappa})\|_1 < k^{-\frac{\beta}{10} k^{r_2 - r_1 - \beta r}}, \quad (288)$$

$$G_r^{(4)}(\vec{\kappa})_{ss'} = 0, \text{ when } 2rk^{\gamma r_2 + 150l\gamma r_1} + 3k^{r_2} < \| |\vec{p}_s| \| + \| |\vec{p}_{s'}| \|. \quad (289)$$

Corollary 6.2. *For the perturbed eigenvalue and its spectral projection the following estimates hold:*

$$\lambda^{(4)}(\vec{\kappa}) = \lambda^{(3)}(\vec{\kappa}) + O_2 \left(k^{-\frac{1}{5}\beta k^{r_2 - r_1 - \beta}} \right), \quad (290)$$

$$\|\mathcal{E}^{(4)}(\vec{\mathcal{X}}) - \mathcal{E}^{(3)}(\vec{\mathcal{X}})\|_1 < 2k^{-\frac{\beta}{10}k^{r_2-r_1}-\beta}. \quad (291)$$

$$|\mathcal{E}^{(4)}(\vec{\mathcal{X}})_{\mathbf{s}\mathbf{s}'}| < k^{-d^{(4)}(\mathbf{s},\mathbf{s}')} \text{, when } \|\vec{p}_{\mathbf{s}}\| > 4k^{r_2} \text{ or } \|\vec{p}_{\mathbf{s}'}\| > 4k^{r_2}, \quad (292)$$

$$d^{(4)}(\mathbf{s},\mathbf{s}') = \frac{1}{8}(\|\vec{p}_{\mathbf{s}}\| + \|\vec{p}_{\mathbf{s}'}\|)k^{-\gamma r_2-150l\gamma r_1}\beta + \frac{1}{10}\beta k^{r_2-r_1}.$$

Lemma 6.3. *Under conditions of Theorem 6.1 the following estimates hold when $\varphi \in \omega^{(4)}(k, \delta)$ or its complex $k^{-r'_3-\delta}$ -neighborhood and $\varkappa \in \mathbb{C} : |\varkappa - \varkappa^{(3)}(\varphi)| < \varepsilon_0^{(4)}k^{-2l+1-\delta}$.*

$$\lambda^{(4)}(\vec{\mathcal{X}}) = \lambda^{(3)}(\vec{\mathcal{X}}) + O_2\left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta}\right), \quad (293)$$

$$\frac{\partial \lambda^{(4)}}{\partial \varkappa} = \frac{\partial \lambda^{(3)}}{\partial \varkappa} + O_2\left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta} M_2\right), \quad M_2 := \frac{k^{2l-1+\delta}}{\varepsilon_0^{(4)}}, \quad (294)$$

$$\frac{\partial \lambda^{(4)}}{\partial \varphi} = \frac{\partial \lambda^{(3)}}{\partial \varphi} + O_2\left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta+r'_3+\delta}\right), \quad (295)$$

$$\frac{\partial^2 \lambda^{(4)}}{\partial \varkappa^2} = \frac{\partial^2 \lambda^{(3)}}{\partial \varkappa^2} + O_2\left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta} M_2^2\right), \quad (296)$$

$$\frac{\partial^2 \lambda^{(4)}}{\partial \varkappa \partial \varphi} = \frac{\partial^2 \lambda^{(3)}}{\partial \varkappa \partial \varphi} + O_2\left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta+r'_3+\delta} M_2\right), \quad (297)$$

$$\frac{\partial^2 \lambda^{(4)}}{\partial \varphi^2} = \frac{\partial^2 \lambda^{(3)}}{\partial \varphi^2} + O_2\left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta+2r'_3+2\delta}\right). \quad (298)$$

Corollary 6.4. *All “ O_2 ”-s on the right hand sides of (293)-(298) can be written as $O_1\left(k^{-\frac{1}{10}\beta k^{r_2-r_1}}\right)$.*

Remark 6.5. *In the proof of Theorem 6.1 and similar statements in every further step of the induction we obtain the estimate of the form (201). It is important to notice that the right hand side of these estimates is always exactly $k^{-\beta}$. It can't become better since it comes from the estimate of the free resolvent on the set of points satisfying $\|\vec{\mathcal{X}} + \vec{p}_{\mathbf{m}}\|_{\mathbb{R}}^{2l} - k^{2l} \geq k^{2l-2-40\mu\delta}$. What changes is the first term in the perturbation series, see e.g. (192), (194) vs (286), (288).*

6.2 Isoenergetic Surface for Operator $H^{(4)}$

The following statement is an analogue of Lemma 5.5.

Lemma 6.6. *1. For every $\lambda := k^{2l}$, $k > k_*$, and φ in the real $\frac{1}{2}k^{-r'_3-\delta}$ -neighborhood of $\omega^{(4)}(k, \delta, \tau)$, there is a unique $\varkappa^{(4)}(\lambda, \varphi)$ in the interval $I_3 := [\varkappa^{(3)}(\lambda, \varphi) - \varepsilon_0^{(4)}k^{-2l+1-\delta}, \varkappa^{(3)}(\lambda, \varphi) + \varepsilon_0^{(4)}k^{-2l+1-\delta}]$, such that*

$$\lambda^{(4)}(\vec{\mathcal{X}}^{(4)}(\lambda, \varphi)) = \lambda, \quad \vec{\mathcal{X}}^{(4)}(\lambda, \varphi) := \varkappa^{(4)}(\lambda, \varphi)\vec{\nu}(\varphi). \quad (299)$$

2. Furthermore, there exists an analytic in φ continuation of $\varkappa^{(4)}(\lambda, \varphi)$ to the complex $\frac{1}{2}k^{-r'_3-\delta}$ -neighborhood of $\omega^{(4)}(k, \delta, \tau)$ such that $\lambda^{(4)}(\vec{\varkappa}^{(4)}(\lambda, \varphi)) = \lambda$. Function $\varkappa^{(4)}(\lambda, \varphi)$ can be represented as $\varkappa^{(4)}(\lambda, \varphi) = \varkappa^{(3)}(\lambda, \varphi) + h^{(4)}(\lambda, \varphi)$, where

$$|h^{(4)}(\varphi)| = O_1 \left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta-2l+1} \right), \quad (300)$$

$$\frac{\partial h^{(4)}}{\partial \varphi} = O_2 \left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta-2l+1+r'_3+\delta} \right), \quad \frac{\partial^2 h^{(4)}}{\partial \varphi^2} = O_4 \left(k^{-\frac{1}{5}\beta k^{r_2-r_1}-\beta-2l+1+2r'_3+2\delta} \right). \quad (301)$$

Let us consider the set of points in $\mathbb{R}^2$ given by the formula: $\vec{\varkappa} = \vec{\varkappa}^{(4)}(\varphi)$, $\varphi \in \omega^{(4)}(k, \delta, \tau)$. By Lemma 6.6 this set of points is a slight distortion of $\mathcal{D}_3$. All the points of this curve satisfy the equation $\lambda^{(4)}(\vec{\varkappa}^{(4)}(\varphi)) = k^{2l}$. We call it isoenergetic surface of the operator $H^{(4)}$ and denote by $\mathcal{D}_4$.

7 Induction

7.1 Inductive formulas for r_n

Now, we are ready to introduce the induction. In fact, STEP IV has been the first inductive step. Here, for the sake of convenience, we reformulate the main statements from the previous step in terms of r_n , $n \geq 3$, and provide necessary comments. First, we choose

$$k^{r_{n-2}} < r_n < k^{\gamma 10^{-7} r_{n-1}}, \quad k^{2\gamma 10^{-4} r_{n-1}} < r'_n < k^{\delta_0 r_{n-1}/2}, \quad n \geq 3. \quad (302)$$

7.2 Preparation for Step $n+1$, $n \geq 4$

7.2.1 Properties of the Quasiperiodic Lattice. Induction

Here we prove the inductive version of the results from Section 5.3.1. We consider $\vec{p}_{\mathbf{m}} = 2\pi(\mathbf{s}_1 + \alpha \mathbf{s}_2)$ with integer vectors $\mathbf{s}_j$ such that $|\mathbf{s}_j| \leq 4k^{r_{n-1}}$. We repeat the arguments from the beginning of Section 4.3. Namely, let $(q, p) \in \mathbb{Z}^2$ be a pair such that $0 < q \leq 4k^{r_{n-1}}$ and

$$|\alpha q + p| \leq 16k^{-r_{n-1}}. \quad (303)$$

We choose a pair (p, q) which gives the best approximation. In particular, p and q are mutually simple. Put $\epsilon_q := \alpha + \frac{p}{q}$. We have

$$k^{-2r_{n-1}\mu} \leq |\epsilon_q| \leq 16q^{-1}k^{-r_{n-1}}. \quad (304)$$

The analogs of Lemmas 5.6–5.8 hold with $n-1$ instead of 2.

We consider the matrix $H^{(n-1)}(\vec{z}) = P(\gamma r_{n-2})H(\vec{z})(\gamma r_{n-2})$ where $\vec{z} \in \mathbb{R}^2$, $P(\gamma r_{n-2})$ is the orthogonal projection corresponding to $\Omega(\gamma r_{n-2})$.⁵ We construct the block structure in $H^{(n-1)}(\vec{z})$ analogous to that in Step $n-1$. The difference is that now we consider any $\vec{z} \in \mathbb{R}^2$, not only $\vec{z}$ being close to $\vec{z}^{(n-2)}(\varphi)$. Correspondingly, we define non-resonant $\mathbf{m}$ not in terms of φ , but in more general terms of inequalities providing convergence of perturbation series. Indeed, we call $\mathbf{m} \in \Omega(\gamma r_{n-2})$ non-resonant if (cf. (69))

$$||\vec{z} + \vec{p}_{\mathbf{m}}|^2 - k^2| > k^{-40\mu\delta}. \quad (305)$$

Obviously, this estimate is stable in the $k^{-41\mu\delta-1}$ -neighborhood of a given $\vec{z}$. Hence, the definition of a non-resonant $\mathbf{m}$ is stable in this neighborhood up to a multiplier $1 + o(1)$ in the r.h.s. of (305). Around each resonant $\mathbf{m}$ we construct k^δ -boxes/clusters (see (73)). Let $P_{\mathbf{m}}$ be the projection on the k^δ -cluster containing $\mathbf{m}$. If

$$|| (P_{\mathbf{m}}(H(\vec{z}) - k^{2l})P_{\mathbf{m}})^{-1} || < k^{4\gamma r'_1} \quad (306)$$

(cf. (103)), then we call the k^δ -cluster effectively non-resonant for a given $\vec{z}$. Note, that the above estimate and, therefore, the definition of an effectively non-resonant k^δ -cluster is stable in the $k^{-4\gamma r'_1 - 2l + 1 - \delta}$ -neighborhood of a given $\vec{z}$. The k^δ -clusters, where (306) is not valid, are called effectively resonant k^δ -clusters. Around each effectively resonant k^δ -cluster, we construct $k^{\gamma r_1}$ -clusters. We sort these clusters into four types: simple, white and black clusters as in Section 4.4.1, using the term “ $\mathbf{m}$ is effectively resonant” instead of “ $\mathbf{m} \in \mathcal{M}^{(2)}$ ”. There is no need to consider a special case of simple clusters here. Note that Lemmas 4.12 – 4.14 are valid for an arbitrary $\vec{z}$, since they are based on Lemmas 4.8 4.10 proven for an arbitrary $\vec{z}$. By analogy with (184), a $k^{\gamma r_1}$ -cluster is called effectively non-resonant if

$$|| (P_{\mathbf{m}}(H(\vec{z}) - k^{2l})P_{\mathbf{m}})^{-1} || < k^{2\mu r_2 + r'_2 N_i^{(1)}}, \quad (307)$$

where $N_i^{(1)}$ corresponds to the color of a $k^{\gamma r_1}$ -cluster, $N_i^{(1)} = k^{\gamma r_1 + 3}, k^{\gamma r_1/2 + \delta_0 r_1}$ or $k^{\gamma r_1/6 - \delta_0 r_1}$. If $n = 4$ we stop here. If $n > 4$, we surround effectively resonant $k^{\gamma r_1}$ -clusters by blocks of the next size, etc. The analogs of Lemmas 4.12 – 4.14 are valid, see Lemmas 5.15 – 5.17, 7.7 – 7.9. Eventually, the $k^{\gamma r_{n-3}}$ -cluster is effectively non-resonant if

$$|| (P_{\mathbf{m}}(H(\vec{z}) - k^{2l})P_{\mathbf{m}})^{-1} || < k^{\mu r_{n-2} + r'_{n-2} N_i^{(n-3)}}, \quad (308)$$

where $N_i^{(n-3)}$ is $N_i^{(n-3)} = k^{\gamma r_{n-3} + 150l\gamma r_{n-4}}, k^{\gamma r_{n-3}/2 + \delta_0 r_{n-3}}, k^{\gamma r_{n-3}/6 - \delta_0 r_{n-3}}$, depending on the color of the cluster (cf. (307), (279)). Further we put $150l\gamma r_0 = 3$. This will make (307) to be a special case of (308) ($n = 4$). Thus, we have constructed a block structure in $H^{(n-1)}(\vec{z})$, which is stable in the $k^{-\rho_{n-2}}$ -neighborhood of a given $\vec{z}$, where $\rho_1 = 4\gamma r'_1 + 2l - 1 + \delta$ and

$$\rho_{n-2} = \mu r_{n-2} + r'_{n-2} k^{\gamma r_{n-3} + 150\gamma r_{n-4}} + 2l - 1 + \delta, \text{ when } n \geq 4.$$

It is not difficult to show that $\rho_{n-2} < r_{n-1}$.

⁵It is a slight abuse of notations, since $H^{(n-1)}$ in Step $n-1$ was defined for $\gamma = 1$.

Definition 7.1. We denote by $J(\vec{\mathcal{Z}})$ the number of the effectively resonant $k^{\gamma_{r_{n-3}}}$ -clusters in $H^{(n-1)}(\vec{\mathcal{Z}})$ for a given $\vec{\mathcal{Z}}$. Further (with a slight abuse of notations) we consider $J(\vec{\mathcal{Z}})$ to be constant in the $k^{-\rho_{n-2}}$ -neighborhood of a given $\vec{\mathcal{Z}}$.

Let $\vec{\mathcal{Z}} = \mathbf{a}\tau_1 + \mathbf{b}$, $|\mathbf{a}| = 1$, $|\mathbf{b}| < 4k^{\gamma_{r_{n-2}}}$. We consider $H^{(n-1)}(\vec{\mathcal{Z}})$ as a function of τ_1 in the complex $k^{-\rho_{n-2}}$ -neighborhood of zero.

Lemma 7.2. The resolvent $(H^{(n-1)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}$ has no more than $k^{2\gamma_{r_{n-3}}}J(\mathbf{b})$ poles τ_{1j} in the complex $2k^{-\rho_{n-2}}$ -neighborhood of zero. It satisfies the following estimate in the complex $k^{-\rho_{n-2}}$ -neighborhood of zero.:

$$\|(H^{(n-1)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}\| < k^{\rho_{n-2}}k^{2\gamma_{r_{n-3}}} \left(\frac{k^{-\rho_{n-2}}}{\varepsilon_0} \right)^{J(\vec{\mathcal{Z}})k^{2\gamma_{r_{n-3}}}}, \quad (309)$$

where $\varepsilon_0 = \min\{k^{-2\rho_{n-2}}, \varepsilon\}$, ε being the distance to the nearest pole $\tau_{1,j}$.

Proof. The lemma is proved by induction. For $n = 3$, see Lemma 5.10. Let us consider the case $n \geq 4$. Recall (Definition 7.1) that $J(\vec{\mathcal{Z}})$ may be considered to be constant in $2k^{-\rho_{n-2}}$ -neighborhood of $\tau_1 = 0$. Hence, $J(\vec{\mathcal{Z}}) = J(\mathbf{b})$ for such $\vec{\mathcal{Z}}$ -s.

Let us consider the collection of all $k^\delta, \dots, k^{\gamma_{r_{n-3}}}$ -clusters $P_{\mathbf{m}}$ for $H^{(n-1)}(\vec{\mathcal{Z}})$. Note that the collection is the same for all such $\vec{\mathcal{Z}}$. We construct the corresponding block operator $\tilde{H}^{(n-2)}(\vec{\mathcal{Z}})$:

$$\tilde{H}^{(n-2)}(\vec{\mathcal{Z}}) = \sum P_{\mathbf{m}} H P_{\mathbf{m}} + H_0(I - \sum P_{\mathbf{m}}).$$

If a $k^{\gamma_{r_{n-3}}}$ -cluster $P_{\mathbf{m}} H(\vec{\mathcal{Z}}) P_{\mathbf{m}}$ is effectively non-resonant, then its resolvent, obviously, has no poles τ_1 in the $2k^{-\rho_{n-2}}$ -neighborhood of $\tau_1 = 0$. The resolvent of each effectively resonant $k^{\gamma_{r_{n-3}}}$ -cluster $P_{\mathbf{m}} H(\vec{\mathcal{Z}}) P_{\mathbf{m}}$ has no more than $N_i^{(n-3)} k^{2\gamma_{r_{n-4}}}$ ($k^{2\gamma_{r_0}}$ is taken to be equal to 8 for $n = 4$) poles τ_{1j} in the $k^{-\rho_{n-3}}$ -neighborhood of $\tau_1 = 0$. It follows from this lemma for the previous step and also Lemmas 4.12–4.14, 5.15–5.17 and 7.7–7.9 for previous steps, which give the estimates for $J(\vec{\mathcal{Z}})$ in the previous steps, based on the color of clusters. Let us consider the union of $k^{-2\rho_{n-2}}$ neighborhoods of these poles and denote it by $\mathbf{D}_{\mathbf{m}}$. By this lemma for $n - 1$, instead of n , each $k^{\gamma_{r_{n-3}}}$ cluster satisfies the estimate

$$\|(P_{\mathbf{m}}(H^{(n-1)}(\vec{\mathcal{Z}}) - k^{2l})P_{\mathbf{m}})^{-1}\| < k^{\rho_{n-3}}k^{2\gamma_{r_{n-4}}}k^{2\rho_{n-2}N_i^{(n-3)}}k^{2\gamma_{r_{n-4}}}$$

outside $\mathbf{D}_{\mathbf{m}}$, $N_i^{(n-3)}$ corresponding the color of the cluster. Note that $\max_i N_i^{(n-3)} = N_1^{(n-3)} < k^{\gamma_{r_{n-3}}+150l\gamma_{r_{n-4}}}$. Therefore, the resolvent $(\tilde{H}^{(n-2)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}$ has no more than $J(\vec{\mathcal{Z}})N_1^{(n-3)}k^{2\gamma_{r_{n-4}}}$ poles τ_{1j} in the complex $k^{-\rho_{n-3}}$ -neighborhood of $\tau_1 = 0$. Let $\mathbf{D} = \cup_{\mathbf{m}} \mathbf{D}_{\mathbf{m}}$, the union being taken over all $\mathbf{m}$ corresponding to all resonant clusters. The number of $\mathbf{m}$ -s in the union, obviously, does not exceed $k^{4\gamma_{r_{n-2}}}$, which is the number of different $\mathbf{m}$ in $H^{(n-1)}(\vec{\mathcal{Z}})$. Therefore, the size of each connected component of $\mathbf{D}$ is less than $k^{-2\rho_{n-2}}k^{4\gamma_{r_{n-2}}} = o(k^{-\rho_{n-2}})$. We are interested only in those components of $\mathbf{D}$, which

are completely in the disk of the radius $2k^{-\rho_{n-2}}$ around $\tau_1 = 0$. Considering as before ⁶, we can show that the perturbation series for the resolvent $(H^{(n-1)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}$ with respect to $(\tilde{H}^{(n-2)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}$ converges on the boundary of $\mathbf{D}$. The resolvents have the same number of poles inside each component of $\mathbf{D}$. Hence, $(H^{(n-1)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}$ has no more than $J(\vec{\mathcal{Z}})N_1^{(n-3)}k^{2\gamma r_{n-4}}$ poles in $\mathbf{D}$. It is easy to see that $J(\vec{\mathcal{Z}})N_1^{(n-3)}k^{2\gamma r_{n-4}} < J(\vec{\mathcal{Z}})k^{2\gamma r_{n-3}}$. The resolvent satisfies the following estimate outside $\mathbf{D}$:

$$\|(H^{(n-1)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}\| < k^{\rho_{n-3}}k^{2\gamma r_{n-4}}k^{2\rho_{n-2}N_1^{(n-3)}k^{2\gamma r_{n-4}}} < k^{\rho_{n-2}}k^{2\gamma r_{n-3}}.$$

Using the maximum principle we obtain (309). $\square$

Next, we introduce

$$\mathcal{S}^{(n-1)}(k, \xi) := \{\vec{\mathcal{Z}} \in \mathbb{R}^2 : \|(H^{(n-1)}(\vec{\mathcal{Z}}) - k^{2l})^{-1}\| > k^\xi\}. \quad (310)$$

It is easy to see that each connected component of $\mathcal{S}^{(n-1)}(k, \xi)$ is bounded by the curves $D(\vec{\mathcal{Z}}, k^{2l} \pm k^{-\xi}) = 0$, where $D(\vec{\mathcal{Z}}, \lambda) = \det(H^{(n-1)}(\vec{\mathcal{Z}}) - \lambda)$.

Lemma 7.3. *Let $\mathbf{l}$ be a segment of a straight line in $\mathbb{R}^2$,*

$$\mathbf{l} := \{\vec{\mathcal{Z}} = \mathbf{a}\tau_1 + \mathbf{b}, \tau_1 \in (0, \eta)\}, \quad |\mathbf{a}| = 1, \quad |\mathbf{b}| < 4k^{\gamma r_{n-2}}, \quad 0 < \eta < k^{-\rho_{n-2}}\}. \quad (311)$$

Suppose both ends of $\mathbf{l}$ belong to a connected component of $\mathcal{S}^{(n-1)}(k, \xi)$. If ξ is sufficiently large, namely, $\xi \geq 2k^{2\gamma r_{n-3}}J(\mathbf{b})\log_k \frac{1}{\eta}$, then, there is an inner part $\mathbf{l}'$ of the segment, which is not in $\mathcal{S}^{(n-1)}(k, \xi)$.

Corollary 7.4. *Let $\vec{\mathcal{Z}} \in \mathcal{S}^{(n-1)}(k, \xi)$ and $\xi > k^{2\gamma r_{n-3}}J(\vec{\mathcal{Z}})\rho_{n-2}$. Then the distance from $\vec{\mathcal{Z}}$ to the boundary of $\mathcal{S}^{(n-1)}(k, \xi)$ is less than $k^{-\tilde{\xi}}$, $\tilde{\xi} = \xi k^{-2\gamma r_{n-3}}J(\vec{\mathcal{Z}})^{-1}$.*

Proof of the corollary. Let us consider a segment of the length $\eta = k^{-\tilde{\xi}}$ starting at $\vec{\mathcal{Z}}$. By the statement of the lemma it intersects a boundary $D(\vec{\mathcal{Z}}, k^{2l} \pm k^{-\xi}) = 0$.

Proof. Choose $\varepsilon = \eta^2$. Using the hypothesis of the lemma, we obtain that the right-hand side of (309) is less than k^ξ outside the discs. Let us estimate the total size (sum of the sizes) of the discs. Indeed, the size of each disc is $2\eta^2$ and the number of discs is, obviously, less $16k^{4\gamma r_{n-2}}$. Therefore, the total size admits the estimate from above: $32\eta^2k^{4\gamma r_{n-2}} = o(\eta)$, since $\eta < k^{-\rho_{n-2}}$. This means there is a part $\mathbf{l}'$ of $\mathbf{l}$ outside these discs. By (309), this part is outside $\mathcal{S}^{(n-1)}(k, \xi)$, when ξ is as described in the statement of the lemma. $\square$

Let $\vec{\mathcal{Z}}_0 \in \mathbb{R}^2$ be fixed and $\mathcal{N}(k, r_{n-1}, \vec{\mathcal{Z}}_0, J_0)$ be the following subset of the lattice $\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}}$, $\mathbf{n} \in \Omega(r_{n-1})$:

$$\mathcal{N}(k, r_{n-1}, \vec{\mathcal{Z}}_0, J_0) = \{\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}} : \mathbf{n} \in \Omega(r_{n-1}) : J(\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}}) \leq J_0\},$$

⁶see the proof of Theorem 5.1 with r_{n-2} instead of r_2 , r_{n-3} instead of r_1 and $k^{\gamma r_{n-3}+150l\gamma r_{n-4}}$ instead of $k^{\gamma r_1+3}$, when one considers black clusters.

J being defined by Definition 7.1. Thus, $\mathcal{N}$ includes only such $\mathbf{n}$ that the surrounding $k^{\gamma r_{n-2}}$ -block contains less than J_0 of effectively resonant points. Let $N(k, r_{n-1}, \vec{\mathcal{Z}}_0, J_0, \xi)$ be the number of points $\vec{\mathcal{Z}}_0 + \vec{p}_{\mathbf{n}}$ in $\mathcal{S}^{(n-1)}(k, \xi) \cap \mathcal{N}(k, r_{n-1}, \vec{\mathcal{Z}}_0, J_0)$.

Lemma 7.5. *If $\xi > 4\mu r_{n-1} J_0 k^{2\gamma r_{n-3}}$, then*

$$N(k, r_{n-1}, \vec{\mathcal{Z}}_0, J_0, \xi) \leq k^{\frac{2}{3}r_{n-1} + 43l\gamma r_{n-2}}. \quad (312)$$

Proof. The proof of the lemma is completely analogous to that of 5.13 up to replacement of 2 by $n - 1$. Instead of Corollary 5.12 we use Corollary 7.4 and the inequality $\rho_{n-2} < r_{n-1}$. $\square$

7.2.2 Model Operator for Step $n + 1$

We repeat for r_n the construction from the subsection 4.4.1, which was done for arbitrary large r_{n-1} . We start with introducing a new notation by analogy with (263) and (268):

$$\Omega_s^{(j)}(r_n) = \{\mathbf{m} \in \Omega(r_n), 0 < p_{\mathbf{m}} < k^{-r'_{j-1}} k^{2\gamma r_{j-2}}\}, \quad j \geq 2, \quad (313)$$

where $k^{2\gamma r_{j-2}}$ is taken to be just 5 when $j = 2$. Note that $\Omega_s^{(j+1)} \subset \Omega_s^{(j)}$ and $\Omega_s^{(j)} = \emptyset$ when $j > n$. Next, let $\mathbf{m} \in \Omega(r_n)$. We denote the $k^{\gamma r_{n-2}}$ -component containing $\mathbf{m}$ by $\tilde{\Pi}(\mathbf{m})$ and the corresponding projector by $\tilde{P}(\mathbf{m})$. For $\mathbf{m}$ belonging to the same $k^{\gamma r_{n-2}}$ -component, $\tilde{\Pi}(\mathbf{m})$ and $\tilde{P}(\mathbf{m})$ are the same. We define $\mathcal{M}^{(n)}$ by the recurrent formula, which starts with $\mathcal{M}^{(3)}$, see (262):

$$\mathcal{M}^{(n)} := \mathcal{M}^{(n)}(\varphi_0, r_n) = \{\mathbf{m} \in \mathcal{M}^{(n-1)}(\varphi_0, r_n) \cup \Omega_s^{(n-1)}(r_n) : \varphi_0 \in \mathcal{O}_{\mathbf{m}}^{(n)}(r'_{n-1}, 1)\}, \quad (314)$$

where $\mathcal{O}_{\mathbf{m}}^{(n)}(r'_{n-1}, \tau)$ is the union of the disks of the radius $\tau k^{-r'_{n-1}}$ with the centers at poles of the resolvent $(\tilde{P}(\mathbf{m})(H(\vec{\mathcal{Z}}^{(n-1)}(\varphi)) - k^{2l}I)\tilde{P}(\mathbf{m}))^{-1}$ in the $k^{-44r'_{n-2}-2l-\delta}$ -neighborhood of φ_0 . For $\mathbf{m}$ belonging to the same $k^{\gamma r_{n-1}}$ -component, the sets $\mathcal{O}_{\mathbf{m}}^{(n)}(r'_{n-1}, \tau)$ are identical. We say that $\mathbf{m} \in \mathcal{M}^{(n)}$ is $k^{\gamma r_{n-2}}$ -resonant. The corresponding $k^{\gamma r_{n-2}}$ -clusters we call resonant too.

Let $\varphi_0 \in \omega^{(n)}(k, \delta, 1)$. By construction of the non-resonant set $\omega^{(n)}(k, \delta, 1)$, we have $\mathcal{M}^{(n)} \cap \Omega(r_{n-1}) = \emptyset$.

Further we use the property of the set $\mathcal{M}^{(n)}$ formulated in the next lemma which is an analogue of the Lemmas 4.11, 5.14.

Lemma 7.6. *Let $r'_{n-1} > 2k^{(\gamma+\delta_0)10^{-4}r_{n-2}-2\delta}$.⁷ Let $1/20 < \gamma' < 20$, $\mathbf{m}_0 \in \Omega(r_n)$ and $\Pi_{\mathbf{m}_0}$ be the $k^{\gamma' r_{n-1}}$ -neighborhood (in $\|\cdot\|$ -norm) of $\mathbf{m}_0$. Then the set $\Pi_{\mathbf{m}_0}$ contains less than $k^{\frac{2}{3}\gamma' r_{n-1} + 50l\gamma r_{n-2}}$ elements of $\mathcal{M}^{(n)}$.*

⁷ We also notice that this condition is consistent with the restriction (302).

Proof. The proof is similar to that of Lemma 5.14 up to the replacement of 3 by n . Instead of Lemma 3.18 we use Lemmas 4.21 ($n=4$) 5.22 ($n=5$) and 7.14 with $n-2$ instead of n when $n > 5$. We also use Lemma 7.5 instead of Lemma 5.13. We use (302) to show that the hypothesis of Lemma 7.5 holds. In particular, we use the inequality $r'_{n-1} \gg 4\mu r_{n-1} k^{2\gamma r_{n-3}}$, following from (302). $\square$

We continue with constructing $k^{\gamma r_{n-2}}$ -clusters in $\Omega(r_n)$, $r_n > r_{n-1}$, the same way we did it for $\Omega(r_2)$ in Section 4.4.1. We call a $k^{\gamma r_{n-2}}$ -cluster resonant if it contains $\mathbf{m} \in \mathcal{M}^{(n)}$, see (262), (314). Next, we repeat the construction after Lemma 4.11 up to the replacement of r_1 by r_{n-1} and δ be γr_{n-2} . Indeed, let us split $\Omega(r_n) \setminus \Omega(r_{n-1})$ into $k^{\gamma r_{n-1}}$ -boxes, $\gamma = \frac{1}{5}$.

First, let's consider $\mathbf{m} \in \Omega_s^{(n)}(r_n)$. As before (see "Simple region", page 47) one can prove that $\Omega_s^{(n)}(r_n) \subset \mathcal{M}(r_n)$; there are no other elements of $\mathcal{M}(r_n)$ in the k^δ -box around $\mathbf{m}$; $\mathbf{m}$ itself can belong or do not belong to $\mathcal{M}^{(j)}(r_n)$, but there are no other elements of $\mathcal{M}^{(j)}(r_n)$ in the $k^{r_{j-1}}$ -box around such $\mathbf{m}$, $j = 2, \dots, n$; and there are no other elements of $\Omega_s^{(n)}(r_n)$ in the $k^{r_{n-1}}$ -box around $\mathbf{m}$.

For each $\mathbf{m} \in \Omega_s^{(n)}(r_n)$ we consider its $k^{r_{n-1}/2}$ -neighborhood in $||| \cdot |||$ norm. The union of such boxes we call the simple region and denote Π_s . The corresponding projection is P_s .

Now, consider all other boxes (all elements $\vec{p}_{\mathbf{m}}$ there satisfy $p_{\mathbf{m}} > k^{-r'_{n-1}} k^{2\gamma r_{n-2}}$). We call a box black if it together with its neighbors contains more than $k^{\gamma r_{n-1}/2 + \delta_0 r_{n-1}}$ elements of $\mathcal{M}^{(n)}$, $\delta_0 = \gamma/100$. Let us consider "black" boxes together with their $k^{\gamma r_{n-1} + \delta_0 r_{n-1}}$ -neighborhoods and call this the black region. We denote the black region by Π_b . The corresponding projector is P_b . By white boxes we mean $k^{\gamma r_{n-1}}$ -boxes which together with its neighbors contain no more than $k^{\gamma r_{n-1}/2 + \delta_0 r_{n-1}}$ elements of $\mathcal{M}^{(n)}$. Every white box we split into "small" boxes of the size $k^{\gamma r_{n-1}/2 + 2\delta_0 r_{n-1}}$. We call a small box "grey" if it together with its neighbors contains more than $k^{\gamma r_{n-1}/6 - \delta_0 r_{n-1}}$ elements of $\mathcal{M}^{(n)}$. Grey small boxes together with its $k^{\gamma r_{n-1}/2 + 2\delta_0 r_{n-1}}$ -neighborhoods we call the grey region. The notation for this region is Π_g . The corresponding projector is P_g . The part of the grey region which is outside the black region, we denote Π'_g and the corresponding projection by P'_g . By a white small box we call a small box which has no more than $k^{\gamma r_{n-1}/6 - \delta_0 r_{n-1}}$ elements of $\mathcal{M}^{(n)}$. In each small white box we consider $k^{\gamma r_{n-1}/6}$ -boxes around each point of $\mathcal{M}^{(n)}$. The union of such $k^{\gamma r_{n-1}/6}$ -boxes we call the white region and denote Π_w . The corresponding projection is P_w . The part of the white region which is outside the black and grey regions, we denote Π'_w and the corresponding projection by P'_w .

We put as before

$$P_r^{(n)} := P_s^{(n)} + P_b^{(n)} + P_g^{(n)'} + P_w^{(n)'}$$

The construction of the non-resonant region is the inductive extension of that for Step IV, see Section 5.3, page 76. Indeed, we start with construction of k^δ clusters in $\Omega(r_n)$. Those of them, who are resonant, we extend to $k^{\gamma r_1}$ clusters, those of them, which are resonant we extend to $k^{\gamma r_2}$ clusters, and so on until we reach the size $k^{\gamma r_{n-2}}$. On each step we construct a colored structure (simple, black, grey, white). If $k^{\gamma r_j}$ -cluster happens to intersect $k^{\gamma r_{j+1}}$ -

cluster, we consider it to be a part of $k^{\gamma_{r_{j+1}}}$ -cluster. Thus, $k^{\gamma_{r_j}}$ -clusters are built around the points of $\mathcal{M}^{(j+1)}(r_n, \varphi_0) \cup \Omega_s^{(j+1)}(r_n) \setminus \left(\mathcal{M}^{(j+1)}(r_{n-1}, \varphi_0) \cup \mathcal{M}^{(j+2)}(r_n, \varphi_0) \cup \Omega_s^{(j+2)}(r_n) \right)$. The set of all other non-resonant $k^{\gamma_{r_j}}$ -clusters we denote by $\Pi_{nr, r_j}^{(n)'}.$ Then

$$\Pi_{nr}^{(n)} := \cup_{j=0}^{n-2} \Pi_{nr, r_j}^{(n)},$$

Those $\Pi_{nr, r_j}^{(n)}$, which intersect with $\Pi_r^{(n)}$ we attach to $\Pi_r^{(n)}$ just slightly abusing the notation (cf. Section 5.3). The part of $\Pi_{nr, r_j}^{(n)}$ which does not intersect with $\Pi_r^{(n)}$ we denote by $\Pi_{nr, r_j}^{(n)'}$. Correspondingly, the part of $\Pi_{nr}^{(n)}$ which does not intersect $\Pi_r^{(n)}$ is denoted by $\Pi_{nr}^{(n)'}$. Further,

$$P^{(n)} := P_r^{(n)} + P_{nr}^{(n)'} + P(r_{n-1}). \quad (315)$$

We continue construction from Section 4.4. Repeating the arguments from the proofs of Lemmas 4.12, 4.13, 4.14 with obvious changes (in particular, using Lemma 7.6 instead of Lemmas 4.11, 5.14) we obtain the following results. (Here and in what follows we will omit superscript (n) when it does not lead to a confusion.)

- Lemma 7.7.** 1. Each Π_b^j contains no more than $k^{\gamma_{r_{n-1}}/2 - \delta_0 r_{n-1} + 150l\gamma_{r_{n-2}}}$ black boxes.
2. The size of Π_b^j in $||| \cdot |||$ norm is less than $k^{3\gamma_{r_{n-1}}/2 + 150l\gamma_{r_{n-2}}}$.
3. Each Π_b^j contains no more than $k^{\gamma_{r_{n-1}} + 150l\gamma_{r_{n-2}}}$ elements of $\mathcal{M}^{(n)}$. Moreover, any box of $||| \cdot |||$ -size $k^{3\gamma_{r_{n-1}}/2 + 150l\gamma_{r_{n-2}}}$ containing Π_b^j has no more than $k^{\gamma_{r_{n-1}} + 150l\gamma_{r_{n-2}}}$ elements of $\mathcal{M}^{(n)}$ inside.

- Lemma 7.8.** 1. Each Π_g^j contains no more than $k^{\gamma_{r_{n-1}}/3 + 2\delta_0 r_{n-1}}$ grey boxes.
2. The size of Π_g^j in $||| \cdot |||$ norm is less than $k^{5\gamma_{r_{n-1}}/6 + 4\delta_0 r_{n-1}}$.
3. Each Π_g^j contains no more than $k^{\gamma_{r_{n-1}}/2 + \delta_0 r_{n-1}}$ elements of $\mathcal{M}^{(n)}$.

- Lemma 7.9.** 1. The size of Π_w^j in $||| \cdot |||$ norm is less than $k^{\gamma_{r_{n-1}}/3 - \delta_0 r_{n-1}}$.
2. Each Π_w^j contains no more than $k^{\gamma_{r_{n-1}}/6 - \delta_0 r_{n-1}}$ points of $\mathcal{M}^{(n)}$.

The construction of the rest of Section 4.4.1 stays unchanged. Let us introduce corresponding notation, formulate the results and provide some comments.

Next lemmas are the analogues of Lemmas 4.17, 4.18, 4.19.

Lemma 7.10. Let $\varphi_0 \in \omega^{(n)}(k, \delta, \tau)$, $|\varphi - \varphi_0| < k^{-k^{r_{n-2}}}$. Then,

$$\left\| \left(P_{nr} \left(H(\vec{\mathcal{K}}^{(n)}(\varphi)) - k^{2l} I \right) P_{nr} \right)^{-1} \right\| < k^{r'_{n-1}} k^{2\gamma_{r_{n-2}}} k^{r'_{n-1}} \leq k^{3\gamma_{r_{n-2}}}. \quad (316)$$

Lemma 7.11. Let $\varphi_0 \in \omega^{(n)}(k, \delta, \tau)$, and $|\varphi - \varphi_0| < k^{-k^{r_{n-2}}}$, $i = 1, 2, 3$. Then,

1. The number of poles of the resolvent $\left(P_i(H(\vec{\mathcal{Z}}^{(n)}(\varphi)) - k^{2l}I)P_i\right)^{-1}$ in the disc $|\varphi - \varphi_0| < k^{-k^{r_{n-2}}}$ is no greater than $N_i^{(n-1)}$, where $N_1^{(n-1)} = k^{\gamma r_{n-1} + 150l\gamma r_{n-2}}$, $N_2^{(n-1)} = k^{\gamma r_{n-1}/2 + \delta_0 r_{n-1}}$, $N_3^{(n-1)} = k^{\gamma r_{n-1}/6 - \delta_0 r_{n-1}}$.
2. Let ε be the distance to the nearest pole of the resolvent in $\mathcal{W}^{(n)}$ and let $\varepsilon_0 = \min\{\varepsilon, k^{-r'_{n-1}}\}$. Then the following estimates hold:

$$\left\| \left(P_i(H(\vec{\mathcal{Z}}^{(n)}(\varphi)) - k^{2l}I)P_i \right)^{-1} \right\| < k^{2r'_{n-1}k^{2\gamma r_{n-2}}} k^{r'_{n-1}} \left(\frac{k^{-r'_{n-1}}}{\varepsilon_0} \right)^{N_i^{(n-1)}} \leq k^{k^{3\gamma r_{n-2}}} \left(\frac{k^{-r'_{n-1}}}{\varepsilon_0} \right)^{N_i^{(n-1)}}, \quad (317)$$

$$\left\| \left(P_i(H(\vec{\mathcal{Z}}^{(n)}(\varphi)) - k^{2l}I)P_i \right)^{-1} \right\|_1 < k^{2r'_{n-1}k^{2\gamma r_{n-2}}} k^{r'_{n-1} + 8\gamma r_{n-1}} \left(\frac{k^{-r'_{n-1}}}{\varepsilon_0} \right)^{N_i^{(n-1)}} \leq k^{k^{3\gamma r_{n-2}}} \left(\frac{k^{-r'_{n-1}}}{\varepsilon_0} \right)^{N_i^{(n-1)}}. \quad (318)$$

Proof. The proof of this lemma is analogous to that of Lemma 4.18 up to the replacement of $\mathcal{M}^{(2)}$ by $\mathcal{M}^{(n)}$, $\mathcal{O}_{\mathbf{m}}^{(2)}$ by $\mathcal{O}_{\mathbf{m}}^{(n)}$, and the shift of indices: δ to r_{n-2} , r_1 to r_{n-1} , etc. We apply Lemmas 7.7–7.9 instead of 4.12–4.14. We apply Lemmas 5.19, 5.20 with $\varepsilon_0 = k^{-r'_3}$ and $p_{\mathbf{m}} > k^{-r'_3 k^{2\gamma r_2}}$ instead of Lemma 4.18, 4.19 for $n = 4$ and Lemmas 7.11, 7.12 with inductively (with $n-1$ instead of n and $\varepsilon_0 = k^{-r'_{n-1}}$, $p_{\mathbf{m}} > k^{-r'_{n-1} k^{2\gamma r'_{n-2}}}$) for further steps. $\square$

Lemma 7.12. Let $\varphi_0 \in \omega^{(n)}(k, \delta, \tau)$. Then, the operator $\left(P_s^j(H(\vec{\mathcal{Z}}^{(n)}(\varphi)) - k^{2l}I)P_s^j\right)^{-1}$ has no more than one pole in the disk $|\varphi - \varphi_0| < k^{-k^{r_{n-2}}}$. Moreover,

$$\left\| \left(P_s^j(H(\vec{\mathcal{Z}}^{(n)}(\varphi)) - k^{2l}I)P_s^j \right)^{-1} \right\| < \frac{8k^{-2l+1}}{p_{\mathbf{m}}\varepsilon_0}, \quad (319)$$

$$\left\| \left(P_s^j(H(\vec{\mathcal{Z}}^{(n)}(\varphi)) - k^{2l}I)P_s^j \right)^{-1} \right\|_1 < \frac{8k^{-2l+1+4r_{n-1}}}{p_{\mathbf{m}}\varepsilon_0}, \quad (320)$$

$\varepsilon_0 = \min\{\varepsilon, k^{-r'_{n-1}}\}$, where ε is the distance to the pole of the operator.

Note that $p_{\mathbf{m}} > k^{-2\mu r_n}$ when $\mathbf{m} \in \Omega(r_n)$. The analogues of Lemma 4.15 and Corollary 4.16 also hold.

7.2.3 Resonant and Nonresonant Sets for Step $n + 1$

We divide $[0, 2\pi)$ into $[2\pi k^{r_{n-2}}] + 1$ intervals $\Delta_m^{(n)}$ with the length not bigger than $k^{-k^{r_{n-2}}}$. If a particular interval belongs to $\mathcal{O}^{(n)}$ we ignore it; otherwise, let $\varphi_0(m) \notin \mathcal{O}^{(n)}$ be a point inside the $\Delta_m^{(n)}$. Let

$$\mathcal{W}_m^{(n)} = \{\varphi \in \mathcal{W}^{(n)} : |\varphi - \varphi_0(m)| < 4k^{-k^{r_{n-2}}}\}. \quad (321)$$

Clearly, neighboring sets $\mathcal{W}_m^{(n)}$ overlap (because of the multiplier 4 in the inequality), they cover $\hat{\mathcal{W}}^{(n)}$, which is the restriction of $\mathcal{W}^{(n)}$ to the $2k^{-k^{r_{n-2}}}$ -neighborhood of $[0, 2\pi)$. For each $\varphi \in \hat{\mathcal{W}}^{(n)}$ there is an m such that $|\varphi - \varphi_0(m)| < 4k^{-k^{r_{n-2}}}$. We consider the poles of the resolvent $(P^{(n)}(H(\vec{\varkappa}^{(n)}(\varphi)) - k^{2l})P^{(n)})^{-1}$ in $\hat{\mathcal{W}}_m^{(n)}$ and denote them by $\varphi_{mj}^{(n)}$, $j = 1, \dots, M_m$. As before, the resolvent has a block structure. The number of blocks clearly cannot exceed the number of elements in $\Omega(r_n)$, i.e. k^{4r_n} . Using the estimates for the number of poles for each block, the estimate being provided by Lemma 7.11 Part 1, we can roughly estimate the number of poles of the resolvent by $k^{4r_n + r_{n-1}}$. Next, let $\mathcal{O}_{mj}^{(n+1)}$ be the disc of the radius $k^{-r'_n}$ around $\varphi_{mj}^{(n)}$.

Definition 7.13. *The set*

$$\mathcal{O}^{(n+1)} = \cup_{mj} \mathcal{O}_{mj}^{(n+1)} \quad (322)$$

we call the $n + 1$ -th resonant set. The set

$$\mathcal{W}^{(n+1)} = \mathcal{W}^{(n)} \setminus \mathcal{O}^{(n+1)} \quad (323)$$

is called the $n + 1$ -th non-resonant set. The set

$$\omega^{(n+1)} = \mathcal{W}^{(n+1)} \cap [0, 2\pi) \quad (324)$$

is called the $n + 1$ -th real non-resonant set.

The following statements can be proven in the same way as Lemmas 4.21, 4.22 and 4.23.

Lemma 7.14. *Let $r'_n > \mu r_n > k^{r_{n-2}}$ ⁸, $\varphi \in \mathcal{W}^{(n+1)}$, $\varphi_0(m)$ corresponds to an interval $\Delta_m^{(n)}$ containing $\Re \varphi$. Let Π be one of the components $\Pi_s^j(\varphi_0(m))$, $\Pi_b^j(\varphi_0(m))$, $\Pi_g^j(\varphi_0(m))$, $\Pi_w^j(\varphi_0(m))$ and $P(\Pi)$ be the projection corresponding to Π . Let also $\varkappa \in \mathbb{C} : |\varkappa - \varkappa^{(n)}(\varphi)| < k^{-r'_n k^{2\gamma_{r_{n-1}}}}$. Then,*

$$\left\| (P(\Pi) (H(\vec{\varkappa}(\varphi)) - k^{2l} I) P(\Pi))^{-1} \right\| < k^{2\mu r_n + r'_n N^{(n-1)}}, \quad (325)$$

$$\left\| (P(\Pi) (H(\vec{\varkappa}(\varphi)) - k^{2l} I) P(\Pi))^{-1} \right\|_1 < k^{(2\mu+1)r_n + r'_n N^{(n-1)}}, \quad (326)$$

$N^{(n-1)}$ corresponding to the color of Π ($N^{(n-1)} = 1$, $k^{\gamma_{r_{n-1}} + 150l\gamma_{r_{n-2}}}$, $k^{\gamma_{r_{n-1}}/2 + \delta_0 r_{n-1}}$, $k^{\gamma_{r_{n-1}}/6 - \delta_0 r_{n-1}}$ for simple, black, grey and white clusters, correspondingly).

⁸These inequalities follow from (302).

Proof. The lemma follows from Lemmas 7.11, 7.12 and the definition of $\mathcal{W}^{(n+1)}$. $\square$

By total size of the set $\mathcal{O}^{(n+1)}$ we mean the sum of the sizes of its connected components.

Lemma 7.15. *Let $r'_n \geq (\mu + 10)r_n$, $r_n > k^{r_n-2}$. Then, the size of each connected component of $\mathcal{O}^{(n+1)}$ is less than $32k^{4r_n-r'_n}$. The total size of $\mathcal{O}^{(n+1)}$ is less than $k^{-r'_n/2}$.*

Lemma 7.16. *Let $\varphi \in \mathcal{W}^{(n)}$ and C_{n+1} be the circle $|z - k^{2l}| = k^{-2r'_n k^{2\gamma r_n-1}}$. Then*

$$\left\| \left(P(r_{n-1})(H(\vec{\mathcal{Z}}^{(n)}(\varphi)) - z)P(r_{n-1}) \right)^{-1} \right\| \leq 4^n k^{2r'_n k^{2\gamma r_n-1}}.$$

7.3 Operator $H^{(n+1)}$. Perturbation Formulas

Let $P(r_n)$ be an orthogonal projector onto $\Omega(r_n) := \{\mathbf{m} : \|\vec{p}_{\mathbf{m}}\| \leq k^{r_n}\}$ and $H^{(n+1)} = P(r_n)HP(r_n)$. We consider $H^{(n+1)}(\vec{\mathcal{Z}}^{(n)}(\varphi))$ as a perturbation of

$$\tilde{H}^{(n)} = \tilde{P}_j^{(n)} H \tilde{P}_j^{(n)} + \left(P(r_n) - \tilde{P}_j^{(n)} \right) H_0, \quad (327)$$

where $H = H(\vec{\mathcal{Z}}^{(n)}(\varphi))$, $H_0 = H_0(\vec{\mathcal{Z}}^{(n)}(\varphi))$, and $\tilde{P}_j^{(n)}$ is the projection $P^{(n)}$, see (315), corresponding to $\varphi_0(j)$ in the interval $\Delta_j^{(n)}$ containing φ . Let

$$W^{(n)} = H^{(n+1)} - \tilde{H}^{(n)} = P(r_n)VP(r_n) - \tilde{P}_j^{(n)}V\tilde{P}_j^{(n)}, \quad (328)$$

$$g_r^{(n+1)}(\vec{\mathcal{Z}}) := \frac{(-1)^r}{2\pi i r} \text{Tr} \oint_{C_{n+1}} \left(W^{(n)}(\tilde{H}^{(n)}(\vec{\mathcal{Z}}) - zI)^{-1} \right)^r dz, \quad (329)$$

$$G_r^{(n)}(\vec{\mathcal{Z}}) := \frac{(-1)^{r+1}}{2\pi i} \oint_{C_{n+1}} (\tilde{H}^{(n)}(\vec{\mathcal{Z}}) - zI)^{-1} \left(W^{(n)}(\tilde{H}^{(n)}(\vec{\mathcal{Z}}) - zI)^{-1} \right)^r dz, \quad (330)$$

where C_{n+1} is the circle $|z - k^{2l}| = \varepsilon_0^{(n+1)}$, $\varepsilon_0^{(n+1)} = k^{-2r'_n k^{2\gamma r_n-1}}$.

Recall that $\beta := 2l - 2 - 41\mu\delta$. The proof of the following statements is analogous to the one in Step III (see Theorem 5.1, Corollary 5.2, Lemma 5.3 and Lemma 5.5).

Theorem 7.17. *Suppose $k > k_*$, φ is in the real $k^{-r'_n-\delta}$ -neighborhood of $\omega^{(n+1)}(k, \delta, \tau)$ and $\varkappa \in \mathbb{R}$, $|\varkappa - \varkappa^{(n)}(\varphi)| \leq \varepsilon_0^{(n+1)} k^{-2l+1-\delta}$, $\vec{\mathcal{Z}} = \varkappa(\cos \varphi, \sin \varphi)$. Then, there exists a single eigenvalue of $H^{(n+1)}(\vec{\mathcal{Z}})$ in the interval $\varepsilon_{n+1}(k, \delta, \tau) = \left(k^{2l} - \varepsilon_0^{(n+1)}, k^{2l} + \varepsilon_0^{(n+1)} \right)$. It is given by the absolutely converging series series:*

$$\lambda^{(n+1)}(\vec{\mathcal{Z}}) = \lambda^{(n)}(\vec{\mathcal{Z}}) + \sum_{r=2}^{\infty} g_r^{(n+1)}(\vec{\mathcal{Z}}). \quad (331)$$

For coefficients $g_r^{(n+1)}(\vec{\mathcal{Z}})$ the following estimates hold:

$$|g_r^{(n+1)}(\vec{\mathcal{Z}})| < k^{-\frac{\beta}{5} k^{r_n-1-r_n-2-\beta(r-1)}}. \quad (332)$$

The corresponding spectral projection is given by the series:

$$\mathcal{E}^{(n+1)}(\vec{\mathfrak{x}}) = \mathcal{E}^{(n)}(\vec{\mathfrak{x}}) + \sum_{r=1}^{\infty} G_r^{(n+1)}(\vec{\mathfrak{x}}), \quad (333)$$

$\mathcal{E}^{(n)}(\vec{\mathfrak{x}})$ being the spectral projection of $H^{(n)}$. The operators $G_r^{(n+1)}(\vec{\mathfrak{x}})$ satisfy the estimates:

$$\|G_r^{(n+1)}(\vec{\mathfrak{x}})\|_1 < k^{-\frac{\beta}{10}k^{r_{n-1}-r_{n-2}-\beta r}}, \quad (334)$$

$$G_r^{(n+1)}(\vec{\mathfrak{x}})_{ss'} = 0, \text{ when } 2rk^{\gamma r_{n-1}+150l\gamma r_{n-2}} + 3k^{r_{n-1}} < \| |\vec{p}_s| \| + \| |\vec{p}_{s'}| \|. \quad (335)$$

Corollary 7.18. For the perturbed eigenvalue and its spectral projection the following estimates hold:

$$\lambda^{(n+1)}(\vec{\mathfrak{x}}) = \lambda^{(n)}(\vec{\mathfrak{x}}) + O_2 \left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta}} \right), \quad (336)$$

$$\| \mathcal{E}^{(n+1)}(\vec{\mathfrak{x}}) - \mathcal{E}^{(n)}(\vec{\mathfrak{x}}) \|_1 < 2k^{-\frac{\beta}{10}k^{r_{n-1}-r_{n-2}-\beta}}. \quad (337)$$

$$|\mathcal{E}^{(n+1)}(\vec{\mathfrak{x}})_{ss'}| < k^{-d^{(n+1)}(s,s')}, \text{ when } \| |\vec{p}_s| \| > 4k^{r_{n-1}} \text{ or } \| |\vec{p}_{s'}| \| > 4k^{r_{n-1}}, \quad (338)$$

$$d^{(n+1)}(s, s') = \frac{1}{8}(\| |\vec{p}_s| \| + \| |\vec{p}_{s'}| \|)k^{-\gamma r_{n-1}-150l\gamma r_{n-2}}\beta + \frac{1}{10}\beta k^{r_{n-1}-r_{n-2}}.$$

Lemma 7.19. Under conditions of Theorem 7.17 the following estimates hold when $\varphi \in \omega^{(n+1)}(k, \delta)$ or its complex $k^{-r'_n-\delta}$ neighborhood and $\mathfrak{x} \in \mathbb{C}$, $|\mathfrak{x} - \mathfrak{x}^{(n)}(\varphi)| < \varepsilon_0^{(n+1)}k^{-2l+1-\delta}$.

$$\lambda^{(n+1)}(\vec{\mathfrak{x}}) = \lambda^{(n)}(\vec{\mathfrak{x}}) + O_2 \left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta}} \right), \quad (339)$$

$$\frac{\partial \lambda^{(n+1)}}{\partial \mathfrak{x}} = \frac{\partial \lambda^{(n)}}{\partial \mathfrak{x}} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta}} M_{n-1} \right), \quad M_{n-1} := \frac{k^{2l-1+\delta}}{\varepsilon_0^{(n+1)}}, \quad (340)$$

$$\frac{\partial \lambda^{(n+1)}}{\partial \varphi} = \frac{\partial \lambda^{(n)}}{\partial \varphi} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta+r'_n+\delta}} \right), \quad (341)$$

$$\frac{\partial^2 \lambda^{(n+1)}}{\partial \mathfrak{x}^2} = \frac{\partial^2 \lambda^{(n)}}{\partial \mathfrak{x}^2} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta}} M_{n-1}^2 \right), \quad (342)$$

$$\frac{\partial^2 \lambda^{(n+1)}}{\partial \mathfrak{x} \partial \varphi} = \frac{\partial^2 \lambda^{(n)}}{\partial \mathfrak{x} \partial \varphi} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta+r'_n+\delta}} M_{n-1} \right), \quad (343)$$

$$\frac{\partial^2 \lambda^{(n+1)}}{\partial \varphi^2} = \frac{\partial^2 \lambda^{(n)}}{\partial \varphi^2} + O_2 \left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta+2r'_n+2\delta}} \right). \quad (344)$$

Corollary 7.20. All “ O_2 ”-s on the right hand sides of (339)-(344) can be written as $O_1 \left(k^{-\frac{1}{10}\beta k^{r_{n-1}-r_{n-2}}} \right)$.

7.4 Isoenergetic Surface for Operator $H^{(n+1)}$

The following statement is an analogue of Lemma 5.5.

Lemma 7.21. 1. For every $\lambda := k^{2l}$, $k > k_*$, and φ in the real $\frac{1}{2}k^{-r'_n-\delta}$ -neighborhood of $\omega^{(n+1)}(k, \delta, \tau)$, there is a unique $\varkappa^{(n+1)}(\lambda, \varphi)$ in the interval $I_n := [\varkappa^{(n)}(\lambda, \varphi) - \varepsilon_0^{(n+1)}k^{-2l+1-\delta}, \varkappa^{(n)}(\lambda, \varphi) + \varepsilon_0^{(n+1)}k^{-2l+1-\delta}]$, such that

$$\lambda^{(n+1)}(\tilde{\varkappa}^{(n+1)}(\lambda, \varphi)) = \lambda, \quad \tilde{\varkappa}^{(n+1)}(\lambda, \varphi) := \varkappa^{(n+1)}(\lambda, \varphi)\vec{\nu}(\varphi). \quad (345)$$

2. Furthermore, there exists an analytic in φ continuation of $\varkappa^{(n+1)}(\lambda, \varphi)$ to the complex $\frac{1}{2}k^{-r'_n-\delta}$ -neighborhood of $\omega^{(n+1)}(k, \delta, \tau)$ such that $\lambda^{(n+1)}(\tilde{\varkappa}^{(n+1)}(\lambda, \varphi)) = \lambda$. Function $\varkappa^{(n+1)}(\lambda, \varphi)$ can be represented as $\varkappa^{(n+1)}(\lambda, \varphi) = \varkappa^{(n)}(\lambda, \varphi) + h^{(n+1)}(\lambda, \varphi)$, where

$$|h^{(n+1)}(\varphi)| = O_1\left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta-2l+1}}\right), \quad (346)$$

$$\frac{\partial h^{(n+1)}}{\partial \varphi} = O_2\left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta-2l+1+r'_n+\delta}}\right), \quad (347)$$

$$\frac{\partial^2 h^{(n+1)}}{\partial \varphi^2} = O_4\left(k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}-\beta-2l+1+2r'_n+2\delta}}\right). \quad (348)$$

Let us consider the set of points in $\mathbb{R}^2$ given by the formula: $\tilde{\varkappa} = \tilde{\varkappa}^{(n+1)}(\varphi)$, $\varphi \in \omega^{(n+1)}(k, \delta, \tau)$. By Lemma 7.21 this set of points is a slight distortion of $\mathcal{D}_n$. All the points of this curve satisfy the equation $\lambda^{(n+1)}(\tilde{\varkappa}^{(n+1)}(\varphi)) = k^{2l}$. We call it isoenergetic surface of the operator $H^{(n+1)}$ and denote by $\mathcal{D}_{n+1}$.

8 Isoenergetic Sets. Generalized Eigenfunctions of H

8.1 Construction of Limit-Isoenergetic Set

At every step n we constructed a set $\mathcal{B}_n(\lambda)$, $\mathcal{B}_n(\lambda) \subset \mathcal{B}_{n-1}(\lambda) \subset S_1$, and a function $\varkappa^{(n)}(\lambda, \vec{\nu})$, $\vec{\nu} \in \mathcal{B}_n(\lambda)$, with the following properties. The set $\mathcal{D}_n(\lambda)$ of vectors $\tilde{\varkappa} = \varkappa^{(n)}(\lambda, \vec{\nu})\vec{\nu}$, $\vec{\nu} \in \mathcal{B}_n(\lambda)$, is a slightly distorted circle with holes, see Fig.1, Fig.2, formula (19) and Lemmas 3.11, 4.4, 5.5, 6.6, 7.21. For any $\tilde{\varkappa}^{(n)}(\lambda, \vec{\nu}) \in \mathcal{D}_n(\lambda)$ there is a single eigenvalue of $H^{(n)}(\tilde{\varkappa}^{(n)})$ equal to λ and given by the perturbation series. Let $\mathcal{B}_\infty(\lambda) = \bigcap_{n=1}^\infty \mathcal{B}_n(\lambda)$. Since $\mathcal{B}_{n+1} \subset \mathcal{B}_n$ for every n , $\mathcal{B}_\infty(\lambda)$ is a unit circle with infinite number of holes, more and more holes of smaller and smaller size appearing at each step.

Lemma 8.1. The length of $\mathcal{B}_\infty(\lambda)$ satisfies estimate (8) with $\gamma_4 = 37\mu\delta$.

Proof. Using Lemmas 3.1 (part 3), 3.19, 4.22, 5.23 and 7.15 and considering that $r_n \gg 37\delta\mu$, we easily conclude that $L(\mathcal{B}_n) = (2\pi + O(k^{-37\mu\delta}))$, $k = \lambda^{1/2l}$ uniformly in n . Since $\mathcal{B}_n$ is a decreasing sequence of sets, (8) holds. $\square$

Let us consider $\varkappa_\infty(\lambda, \vec{\nu}) = \lim_{n \rightarrow \infty} \varkappa^{(n)}(\lambda, \vec{\nu})$, $\vec{\nu} \in \mathcal{B}_\infty(\lambda)$.

Lemma 8.2. *The limit $\kappa_\infty(\lambda, \vec{v})$ exists for any $\vec{v} \in \mathcal{B}_\infty(\lambda)$ and the following estimates hold:*

$$\begin{aligned} |\kappa_\infty(\lambda, \vec{v}) - \lambda^{1/2l}| &< ck^{-4l+1+(80\mu+6)\delta}, \\ |\kappa_\infty(\lambda, \vec{v}) - \kappa^{(1)}(\lambda, \vec{v})| &< ck^{-2k^\delta Q^{-1}} k^{-4l+5+48\delta}, \\ |\kappa_\infty(\lambda, \vec{v}) - \kappa^{(n)}(\lambda, \vec{v})| &< 3k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}}}, \quad n \geq 2. \end{aligned} \quad (349)$$

Corollary 8.3. *For every $\vec{v} \in \mathcal{B}_\infty(\lambda)$ estimate (9) holds, where $\gamma_5 = (4l - 1 - (80\mu + 6)\delta)/2l > 0$.*

The lemma easily follows from the estimates (65), (138), (249), (300) and (346).

Estimates (66), (139) (250), (301) and (347) justify convergence of the series $\sum_{n=1}^\infty \frac{\partial h_n}{\partial \varphi}$, and hence, of the sequence $\frac{\partial \kappa^{(n)}}{\partial \varphi}$. We denote the limit of this sequence by $\frac{\partial \kappa_\infty}{\partial \varphi}$.

Lemma 8.4. *The estimate (23) with $\gamma_{11} = (4l - 1 - (120\mu + 7)\delta)/2l > 0$, holds for any $\vec{v} \in \mathcal{B}_\infty(\lambda)$.*

We define $\mathcal{D}_\infty(\lambda)$ by (7). Clearly, $\mathcal{D}_\infty(\lambda)$ is a slightly distorted circle of radius k with infinite number of holes. We can assign a tangent vector $\frac{\partial \kappa_\infty}{\partial \varphi} \vec{v} + \kappa_\infty \vec{\mu}$, $\vec{\mu} = (-\sin \varphi, \cos \varphi)$ to the curve $\mathcal{D}_\infty(\lambda)$, this tangent vector being the limit of corresponding tangent vectors for curves $\mathcal{D}_n(\lambda)$ at points $\vec{\kappa}^{(n)}(\lambda, \vec{v})$ as $n \rightarrow \infty$.

Next we show that $\mathcal{D}_\infty(\lambda)$ is an isoenergetic curve for H . Namely for every $\vec{\kappa} \in \mathcal{D}_\infty(\lambda)$ there is a generalized eigenfunction $\Psi_\infty(\vec{\kappa}, \vec{x})$ such that $H\Psi_\infty = \lambda\Psi_\infty$.

8.2 Generalized Eigenfunctions of H

We show that for every $\vec{\kappa}$ in a set

$$\mathcal{G}_\infty = \cup_{\lambda > \lambda_*} \mathcal{D}_\infty(\lambda), \quad \lambda_* = k_*^{2l},$$

there is a solution $\Psi_\infty(\vec{\kappa}, \vec{x})$ of the equation for eigenfunctions:

$$(-\Delta)^l \Psi_\infty(\vec{\kappa}, \vec{x}) + V(\vec{x}) \Psi_\infty(\vec{\kappa}, \vec{x}) = \lambda_\infty(\vec{\kappa}) \Psi_\infty(\vec{\kappa}, \vec{x}), \quad (350)$$

which can be represented in the form

$$\Psi_\infty(\vec{\kappa}, \vec{x}) = e^{i\langle \vec{\kappa}, \vec{x} \rangle} \left(1 + u_\infty(\vec{\kappa}, \vec{x}) \right), \quad \|u_\infty(\vec{\kappa}, \vec{x})\|_{L_\infty(\mathbb{R}^2)} = O(|\vec{\kappa}|^{-\gamma_1}), \quad (351)$$

where $u_\infty(\vec{\kappa}, \vec{x})$ is a quasi-periodic function, $\gamma_1 = 2l - 1 - 45\mu\delta > 0$; the eigenvalue $\lambda_\infty(\vec{\kappa})$ satisfies the asymptotic formula:

$$\lambda_\infty(\vec{\kappa}) = |\vec{\kappa}|^{2l} + O(|\vec{\kappa}|^{-\gamma_2}), \quad \gamma_2 = 2l - (80\mu + 6)\delta > 0. \quad (352)$$

We also show that the set $\mathcal{G}_\infty$ satisfies (6).

In fact, by (349), any $\vec{\mathcal{Z}} \in \mathcal{D}_\infty(\lambda)$ belongs to the $k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}}}$ -neighborhood of $\mathcal{D}_n(\lambda)$, $n \geq 3$. Let us consider spectral projectors $\mathcal{E}^{(n)}$, each of them being defined in a finite dimensional space of sequences with indices in $\Omega(r_{n-1})$, $r_0 := \delta$. We extend each of them to the whole space $\ell^2(\mathbb{Z}^4)$ by putting it to be zero into the orthogonal complement of $\ell^2(\Omega(r_{n-1}))$. This way they all act in space $\ell^2(\mathbb{Z}^4)$. Applying the perturbation formulae proved in the previous sections (see Corollaries 3.4, 4.2, 5.2, 6.2, 7.18), we obtain the following inequalities:

$$\begin{aligned} \|\mathcal{E}^{(1)}(\vec{\mathcal{Z}}) - \mathcal{E}_0(\vec{\mathcal{Z}})\|_1 &< ck^{-\gamma_0}, \\ \|\mathcal{E}^{(2)}(\vec{\mathcal{Z}}) - \mathcal{E}^{(1)}(\vec{\mathcal{Z}})\|_1 &< ck^{-k^\delta Q^{-1}+2-\gamma_0}, \quad \gamma_0 := 2l - 1 - 44\mu\delta, \\ \|\mathcal{E}^{(n)}(\vec{\mathcal{Z}}) - \mathcal{E}^{(n-1)}(\vec{\mathcal{Z}})\|_1 &< 2k^{-\frac{1}{10}\beta k^{r_{n-2}-r_{n-3}-\beta}}, \quad n \geq 3, \end{aligned} \quad (353)$$

$$\begin{aligned} |\lambda^{(1)}(\vec{\mathcal{Z}}) - |\vec{\mathcal{Z}}|^{2l}| &< ck^{-\gamma_2}, \quad |\lambda^{(2)}(\vec{\mathcal{Z}}) - \lambda^{(1)}(\vec{\mathcal{Z}})| < ck^{-2k^\delta Q^{-1}+4-\gamma_2}, \\ |\lambda^{(n)}(\vec{\mathcal{Z}}) - \lambda^{(n-1)}(\vec{\mathcal{Z}})| &< 2k^{-\frac{1}{5}\beta k^{r_{n-2}-r_{n-3}-\beta}}, \quad n \geq 3, \end{aligned} \quad (354)$$

where $\lambda^{(n+1)}(\vec{\mathcal{Z}})$ is the eigenvalue corresponding to $\mathcal{E}^{(n+1)}(\vec{\mathcal{Z}})$, $\mathcal{E}_0(\vec{\mathcal{Z}})$ corresponds to $V = 0$.

Remark 8.5. We see from (349), that any $\vec{\mathcal{Z}} \in \mathcal{D}_\infty(\lambda)$ belongs to the $k^{-\frac{1}{5}\beta k^{r_{n-1}-r_{n-2}}}$ -neighborhood of $\mathcal{D}_n(\lambda)$, $n \geq 3$. Applying perturbation formulae for n -th step, we easily obtain that there is an eigenvalue $\lambda^{(n)}(\vec{\mathcal{Z}})$ of $H^{(n)}(\vec{\mathcal{Z}})$ satisfying the estimate $\lambda^{(n)}(\vec{\mathcal{Z}}) = \lambda + \delta_n$, $\delta_n = o(1)$ as $n \rightarrow \infty$, the eigenvalue $\lambda^{(n)}(\vec{\mathcal{Z}})$ being given by a perturbation series of the type (331). Hence, for every $\vec{\mathcal{Z}} \in \mathcal{D}_\infty(\lambda)$ there is a limit:

$$\lim_{n \rightarrow \infty} \lambda^{(n)}(\vec{\mathcal{Z}}) = \lambda. \quad (355)$$

Let $\mathbf{v}^{(n)}$ be a unit vector corresponding to the projection $\mathcal{E}^{(n)}(\vec{\mathcal{Z}})$, $\mathcal{E}^{(n)}(\vec{\mathcal{Z}}) = (\cdot, \mathbf{v}^{(n)})\mathbf{v}^{(n)}$, $\mathbf{v}^{(n)} = \{v_s^{(n)}\}_{s \in \mathbb{Z}^4} \in \ell^2(\mathbb{Z}^4)$. By construction, $v_s^{(n)} = 0$ when $s \notin \Omega(r_{n-1})$. Let us consider the linear combination of exponents corresponding to this vector:

$$\Psi_n(\vec{\mathcal{Z}}, \vec{x}) = \sum_{s \in \Omega(r_{n-1})} v_s^{(n)} e^{i(\vec{\mathcal{Z}} + \vec{p}_s, \vec{x})}.$$

Lemma 8.6. Function $\Psi_n(\vec{\mathcal{Z}}, \vec{x})$, $n \geq 4$, satisfies the equation:

$$(-\Delta)^l \Psi_n(\vec{\mathcal{Z}}, \vec{x}) + V(\vec{x}) \Psi_n(\vec{\mathcal{Z}}, \vec{x}) = \lambda_n(\vec{\mathcal{Z}}) \Psi_n(\vec{\mathcal{Z}}, \vec{x}) + g_n(\vec{\mathcal{Z}}, \vec{x}),$$

the vector $\mathbf{g}_n$ of the Fourier coefficients of $g_n(\vec{\mathcal{Z}}, \vec{x})$ satisfying the estimate:

$$\|\mathbf{g}_n\|_{\ell^1(\mathbb{Z}^4)} < k^{-k^{\frac{1}{2}r_{n-1}}}. \quad (356)$$

Coefficients $(\mathbf{g}_n)_s$ can differ from zero only when $k^{r_{n-1}} < \|\vec{p}_s\| \leq k^{r_{n-1}} + Q$. Function $g_n(\vec{\mathcal{Z}}, \vec{x})$ obeys the estimate:

$$\|g_n\|_{L_\infty(\mathbb{R}^2)} < k^{-k^{\frac{1}{2}r_{n-1}}}. \quad (357)$$

Proof. Let $P(r_{n-1})$ be the projection in $\ell^2(\mathbb{Z}^4)$ on the subspace corresponding to $\Omega(r_{n-1})$. By construction, $P(r_{n-1})\mathbf{v}^{(n)} = \mathbf{v}^{(n)}$ and

$$H_0\mathbf{v}^{(n)} + P(r_{n-1})VP(r_{n-1})\mathbf{v}^{(n)} = \lambda_n(\vec{x})\mathbf{v}^{(n)}.$$

Since V is a trigonometric polynomial,

$$(I - P(r_{n-1}))VP(r_{n-1}) = (I - P(r_{n-1}))VP_\partial(r_{n-1}),$$

where $P_\partial(r_{n-1})_{\mathbf{m}\mathbf{m}} = 1$ only if $\mathbf{m}$ is in the Q -vicinity of the boundary. Using (338) with n instead of $n+1$, we obtain: $\|P_\partial(r_{n-1})\mathcal{E}^{(n)}\| < k^{-k^{r_{n-1}}(1-\gamma)}$ and, hence, $\|P_\partial(r_{n-1})\mathbf{v}^{(n)}\| < k^{-k^{r_{n-1}}(1-\gamma)}$. Therefore, $\|(I - P(r_{n-1}))VP(r_{n-1})\mathbf{v}^{(n)}\| < \|V\|k^{-k^{r_{n-1}}(1-\gamma)}$. It follows that $H(\vec{x})\mathbf{v}^{(n)} = \lambda_n(\vec{x})\mathbf{v}^{(n)} + \mathbf{g}_n$, where $\|\mathbf{g}_n\|_{\ell^2(\mathbb{Z}^4)} < \|V\|k^{-k^{r_{n-1}}(1-\gamma)}$. Note that elements $(\mathbf{g}_n)_s$ are equal to zero when $\|\vec{p}_s\| \leq k^{r_{n-1}}$ or $\|\vec{p}_s\| > k^{r_{n-1}} + Q$. Therefore, (356) holds. Estimate (357) follows. $\square$

Lemma 8.7. *Functions $\Psi_n(\vec{x}, \vec{x})$ satisfy the inequalities:*

$$\|\Psi_1 - \Psi_0\|_{L_\infty(\mathbb{R}^2)} < ck^{-\gamma_0+2\delta}, \quad \|(-\Delta)^l\Psi_1 - (-\Delta)^l\Psi_0\|_{L_\infty(\mathbb{R}^2)} < ck^{-\gamma_0+2\delta+2l}, \quad \Psi_0(\vec{x}) = e^{i\langle \vec{x}, \vec{x} \rangle}, \quad (358)$$

$$\begin{aligned} \|\Psi_2 - \Psi_1\|_{L_\infty(\mathbb{R}^2)} &< ck^{-k^\delta Q^{-1}+2-\gamma_0+2r_1}, \\ \|(-\Delta)^l\Psi_2 - (-\Delta)^l\Psi_1\|_{L_\infty(\mathbb{R}^2)} &< ck^{-k^\delta Q^{-1}+2-\gamma_0+(2+2l)r_1}, \end{aligned} \quad (359)$$

$$\begin{aligned} \|\Psi_n - \Psi_{n-1}\|_{L_\infty(\mathbb{R}^2)} &< k^{-\frac{1}{10}\beta k^{r_{n-2}-r_{n-3}+2r_{n-1}}}, \\ \|(-\Delta)^l\Psi_n - (-\Delta)^l\Psi_{n-1}\|_{L_\infty(\mathbb{R}^2)} &< (2\pi)^{2l}k^{-\frac{1}{10}\beta k^{r_{n-2}-r_{n-3}+(2+2l)r_{n-1}}}, \quad n \geq 3. \end{aligned} \quad (360)$$

Corollary 8.8. *All functions Ψ_n , $n = 0, 1, \dots$, obey the estimate $\|\Psi_n\|_{L_\infty(\mathbb{R}^2)} < 1 + Ck^{-\gamma_0+2\delta}$ uniformly in n .*

Proof. Using (353) and considering that $v_s^{(n)}$ are equal to zero when $\|\vec{p}_s\| > k^{r_{n-1}}$, we obtain

$$\begin{aligned} \|\mathbf{v}^{(1)} - \mathbf{v}^{(0)}\|_{\ell^1(\mathbb{Z}^4)} &< ck^{-\gamma_0+2\delta}, \quad \|\mathbf{v}^{(2)} - \mathbf{v}^{(1)}\|_{\ell^1(\mathbb{Z}^4)} < ck^{-k^\delta Q^{-1}+2-\gamma_0+2r_1}, \\ \|\mathbf{v}^{(n)} - \mathbf{v}^{(n-1)}\|_{\ell^1(\mathbb{Z}^4)} &< 3k^{-\frac{1}{10}\beta k^{r_{n-2}-r_{n-3}+2r_{n-1}-\beta}}, \quad n \geq 3. \end{aligned} \quad (361)$$

$$\begin{aligned} \|H_0(\mathbf{v}^{(1)} - \mathbf{v}^{(0)})\|_{\ell^1(\mathbb{Z}^4)} &< ck^{-\gamma_0+2\delta+2l}, \quad \|H_0(\mathbf{v}^{(2)} - \mathbf{v}^{(1)})\|_{\ell^1(\mathbb{Z}^4)} < ck^{-k^\delta Q^{-1}+2-\gamma_0+(2+2l)r_1}, \\ \|H_0(\mathbf{v}^{(n)} - \mathbf{v}^{(n-1)})\|_{\ell^1(\mathbb{Z}^4)} &< (2\pi)^{2l}k^{-\frac{1}{10}\beta k^{r_{n-2}-r_{n-3}+(2+2l)r_{n-1}-\beta}}, \quad n \geq 3. \end{aligned} \quad (362)$$

Now (358) – (360) easily follow. $\square$

Theorem 8.9. For every $\lambda > k_*^{2l}$ and $\vec{\mathcal{Z}} \in \mathcal{D}_\infty(\lambda)$ the sequence of functions $\Psi_n(\vec{\mathcal{Z}}, \vec{x})$ converges in $L_\infty(\mathbb{R}^2)$ and $W_{2,loc}^{2l}(\mathbb{R}^2)$. The limit function $\Psi_\infty(\vec{\mathcal{Z}}, \vec{x}) := \lim_{n \rightarrow \infty} \Psi_n(\vec{\mathcal{Z}}, \vec{x})$, is a quasi-periodic function:

$$\Psi_\infty(\vec{\mathcal{Z}}, \vec{x}) = \sum_{\mathbf{s} \in \mathbb{Z}^4} (v_\infty)_{\mathbf{s}} e^{i\langle \vec{\mathcal{Z}} + \vec{p}_{\mathbf{s}}, \vec{x} \rangle}, \quad (363)$$

where $\mathbf{v}_\infty = \{(v_\infty)_{\mathbf{s}}\}_{\mathbf{s} \in \mathbb{Z}^4} \in \ell^1(\mathbb{Z}^4)$ and $\|\mathbf{v}_\infty\|_{\ell^2(\mathbb{Z}^4)} = 1$. The function $\Psi_\infty(\vec{\mathcal{Z}}, \vec{x})$ satisfies the equation

$$(-\Delta)^l \Psi_\infty(\vec{\mathcal{Z}}, \vec{x}) + V(\vec{x}) \Psi_\infty(\vec{\mathcal{Z}}, \vec{x}) = \lambda \Psi_\infty(\vec{\mathcal{Z}}, \vec{x}). \quad (364)$$

It can be represented in the form

$$\Psi_\infty(\vec{\mathcal{Z}}, \vec{x}) = e^{i\langle \vec{\mathcal{Z}}, \vec{x} \rangle} (1 + u_\infty(\vec{\mathcal{Z}}, \vec{x})), \quad (365)$$

where $u_\infty(\vec{\mathcal{Z}}, \vec{x})$ is a quasi-periodic function:

$$u_\infty(\vec{\mathcal{Z}}, \vec{x}) = \sum_{n=1}^{\infty} u_n(\vec{\mathcal{Z}}, \vec{x}), \quad (366)$$

$$u_n(\vec{\mathcal{Z}}, \vec{x}) = \sum_{\mathbf{s} \in \Omega(r_{n-1})} (v_{\mathbf{s}}^{(n)} - v_{\mathbf{s}}^{(n-1)}) e^{i\langle \vec{p}_{\mathbf{s}}, \vec{x} \rangle}, \quad r_0 := \delta, \quad (367)$$

functions u_n satisfying the estimates:

$$\|u_1\|_{L_\infty(\mathbb{R}^2)} < ck^{-\gamma_0+2\delta}, \quad \|u_2\|_{L_\infty(\mathbb{R}^2)} < ck^{-k^\delta Q^{-1}+2-\gamma_0+2r_1}, \quad (368)$$

$$\|u_n\|_{L_\infty(\mathbb{R}^2)} < k^{-\frac{1}{10}\beta k^{r_{n-2}-r_{n-3}+2r_{n-1}}}, \quad n \geq 3. \quad (369)$$

Corollary 8.10. Function $u_\infty(\vec{\mathcal{Z}}, \vec{x})$ obeys the estimate (351).

Proof. Using (361), (362), we obtain that the sequence $\mathbf{v}^{(n)}$ has the limit in $\ell^1(\mathbb{Z}^4)$. We denote this limit by $\mathbf{v}_\infty$. Since, vectors $\mathbf{v}^{(n)}$ are normalized in $\ell^2(\mathbb{Z}^4)$,

$$\|\mathbf{v}_\infty\|_{\ell^2(\mathbb{Z}^4)} = 1. \quad (370)$$

By (360), we obtain that $\Psi_n(\vec{\mathcal{Z}}, \vec{x})$ is a Cauchy sequence in $L_\infty(\mathbb{R}^2)$ and $W_{2,loc}^{2l}(\mathbb{R}^2)$. Let $\Psi_\infty(\vec{\mathcal{Z}}, \vec{x}) = \lim_{n \rightarrow \infty} \Psi_n(\vec{\mathcal{Z}}, \vec{x})$. This limit is defined pointwise uniformly in $\vec{x}$ and in $W_{2,loc}^{2l}(\mathbb{R}^2)$. Noting also that $\lim \lambda_n(\vec{\mathcal{Z}}) = \lambda$, and taking into account Lemma 8.6 we obtain that (364) holds.

Let us show that Ψ_∞ is a quasi-periodic function. Obviously,

$$\Psi_\infty = \Psi_0 + \sum_{n=1}^{\infty} (\Psi_n - \Psi_{n-1}),$$

the series converging in $L_\infty(\mathbb{R}^2)$ by (360). Introducing $u_n := e^{-i\langle \vec{\mathcal{Z}}, \vec{x} \rangle} (\Psi_n - \Psi_{n-1})$, we arrive at (365), (366). Note that u_n has a form (367). Estimates (368), (369) follow from (358), (360). $\square$

Theorem 8.11. *Formulae (350), (351) and (352) hold for every $\vec{x} \in \mathcal{G}_\infty$. The set $\mathcal{G}_\infty$ is Lebesgue measurable and satisfies (6) with $\gamma_3 = 37\mu\delta$.*

Proof. By Theorem 8.9, (350), (351) hold, where $\lambda_\infty(\vec{x}) = \lambda$ for $\vec{x} \in \mathcal{D}_\infty(\lambda)$. Using (9), which is proven in Corollary 8.3, with $\varkappa_\infty = |\vec{x}|$, we easily obtain (352). It remains to prove (6). Let us consider a small region $U_n(\lambda_0)$ around an isoenergetic surface $\mathcal{D}_n(\lambda_0)$, $\lambda_0 > k_*^{2l}$. Namely, $U_n(\lambda_0) = \cup_{|\lambda - \lambda_0| < \varepsilon_0^{(n+1)}} \mathcal{D}_n(\lambda)$, $k = \lambda_0^{1/2l}$. By Theorem 7.17 the construction of the n -th non-resonant set is stable in $\varepsilon_0^{(n)}$ -neighborhood of λ_0 . Therefore, in fact, we can (and for the sake of convenience will) assume that the sets $\omega^{(n)}(\lambda)$ are chosen to be equal to $\omega^{(n)}(\lambda_0)$ for $|\lambda - \lambda_0| < \varepsilon_0^{(n)}$. Thus, $U_n(\lambda_0)$ is an open set (a distorted ring with holes) and $|U_n(\lambda_0)| = l^{-1} \varepsilon_0^{(n)} k^{-2l+1} 2\pi k (1 + O(k^{-37\mu\delta}))$. It easily follows from (67) and (346) that $U_{n+1} \subset U_n$. Definition of $\mathcal{D}_\infty(\lambda_0)$ yield: $\mathcal{D}_\infty(\lambda_0) = \cap_{n=1}^\infty U_n(\lambda_0)$. Hence, $\mathcal{G}_\infty = \cap_{n=1}^\infty \mathcal{G}_n$, where

$$\mathcal{G}_n = \cup_{\lambda > \lambda_*} \mathcal{D}_n(\lambda), \quad \lambda_* = k_*^{2l}. \quad (371)$$

Considering that $U_{n+1} \subset U_n$ for every $\lambda_0 > \lambda_*$, we obtain $\mathcal{G}_{n+1} \subset \mathcal{G}_n$. Hence, $|\mathcal{G}_\infty \cap \mathbf{B}_R| = \lim_{n \rightarrow \infty} |\mathcal{G}_n \cap \mathbf{B}_R|$. Calculating the volume of the region $\cup_{\lambda_* < \lambda < R^{2l}} U_n(\lambda)$, we easily conclude $|\mathcal{G}_n \cap \mathbf{B}_R| = |\mathbf{B}_R| (1 + O(R^{-37\mu\delta}))$ uniformly in n . Thus, we have obtained (6) with $\gamma_3 = 37\mu\delta$. $\square$

Theorem 8.12 (Bethe-Sommerfeld Conjecture). *The spectrum of operator H contains a semi-axis.*

Proof. The theorem immediately follows from the fact that the equation (364) has a bounded solution $\Psi_\infty(\vec{x}, \vec{x})$ for every sufficiently large λ . $\square$

9 Proof of Absolute Continuity of the Spectrum

The proof is somewhat analogous to that for the case limit-periodic potentials [28]. We will just refer to [28] in some places.

9.1 Operators $E_n(\mathcal{G}'_n)$, $\mathcal{G}'_n \subset \mathcal{G}_n$

Let us consider the open sets $\mathcal{G}_n$ given by (371). There is a family of eigenfunctions $\Psi_n(\vec{x}, \vec{x})$, $\vec{x} \in \mathcal{G}_n$, of the operator $H^{(n)}$, which are described by the perturbation formulas (11), (15). Let, $\mathcal{G}'_n \subset \mathcal{G}_n$, where $\mathcal{G}'_n$ is Lebesgue measurable and bounded. Let

$$E_n(\mathcal{G}'_n) F = \frac{1}{4\pi^2} \int_{\mathcal{G}'_n} (F, \Psi_n(\vec{x})) \Psi_n(\vec{x}) d\vec{x} \quad (372)$$

for any $F \in C_0^\infty(\mathbb{R}^2)$, here and below $(\cdot, \cdot)$ is the canonical scalar product in $L_2(\mathbb{R}^2)$, i.e.,

$$(F, \Psi_n(\vec{x})) = \int_{\mathbb{R}^2} F(\vec{x}) \overline{\Psi_n(\vec{x}, \vec{x})} d\vec{x}.$$

We will show that $E_n(\mathcal{G}'_n)$ is almost a projector in $L_2(\mathbb{R}^2)$ in the sense: $E_n(\mathcal{G}'_n) = E_n^*(\mathcal{G}'_n)$, $E_n^2(\mathcal{G}'_n) = E_n(\mathcal{G}'_n) + o(1)$, where $o(1)$ is in the class of bounded operators as $n \rightarrow \infty$. First, we note that (372) can be rewritten in the form:

$$E_n(\mathcal{G}'_n) = S_n(\mathcal{G}'_n) T_n(\mathcal{G}'_n), \quad (373)$$

$$\begin{aligned} T_n : L_2(\mathbb{R}^2) &\rightarrow L_2(\mathcal{G}'_n), \quad S_n : L_2(\mathcal{G}'_n) \rightarrow L_2(\mathbb{R}^2), \\ T_n F &= \frac{1}{2\pi} (F, \Psi_n(\vec{\kappa})) \quad \text{for any } F \in C_0^\infty(\mathbb{R}^2), \end{aligned} \quad (374)$$

$T_n F$ being in $L_\infty(\mathcal{G}'_n)$, and,

$$S_n f = \frac{1}{2\pi} \int_{\mathcal{G}'_n} f(\vec{\kappa}) \Psi_n(\vec{\kappa}, \vec{x}) d\vec{\kappa} \quad \text{for any } f \in L_\infty(\mathcal{G}'_n). \quad (375)$$

Note that $S_n f \in L_2(\mathbb{R}^2)$, since Ψ_n is a finite combination of exponentials $e^{i\langle \vec{\kappa} + \vec{p}_q, \vec{x} \rangle}$.

Lemma 9.1. *Let $\mathcal{G}'_n$ be bounded and $f(\vec{\kappa}), g(\vec{\kappa}) \in L_\infty(\mathcal{G}'_n)$. Then,*

$$(S_n f, S_n g)_{L_2(\mathbb{R}^2)} =_{n \rightarrow \infty} (f, g)_{L_2(\mathcal{G}'_n)} + o(1) \|f\|_{L_2(\mathcal{G}'_n)} \|g\|_{L_2(\mathcal{G}'_n)}. \quad (376)$$

where $o(1)$ goes to zero uniformly in f, g and $\mathcal{G}'_n$ as $n \rightarrow \infty$; namely, $|o(1)| < \xi_*^{-r_{n-3}(\xi_*)}$, where $\xi_* = \inf_{\vec{\xi} \in \mathcal{G}'_n} |\vec{\xi}|$.

Corollary 9.2. *The following relation holds:*

$$\left| (S_n f, S_n g)_{L_2(\mathbb{R}^2)} \right| < (1 + o(1)) \|f\|_{L_\infty(\mathcal{G}'_n)} \|g\|_{L_\infty(\mathcal{G}'_n)} |\mathcal{G}'_n|, \quad (377)$$

where $|\mathcal{G}'_n|$ is the Lebesgue measure of $\mathcal{G}'_n$.

Corollary 9.3. *The operator S_n is bounded and $\|S_n\| =_{n \rightarrow \infty} 1 + o(1)$.*

Proof. The function $\Psi_n(\vec{\kappa}, \vec{x})$ can be represented as a combination of plane waves:

$$\Psi_n(\vec{\kappa}, \vec{x}) = \sum_{\mathbf{m} \in \mathbb{Z}^2} v_{\mathbf{m}}^{(n)}(\vec{\kappa}) \exp\{i\langle \vec{\kappa} + \vec{p}_{\mathbf{m}}, \vec{x} \rangle\}, \quad (378)$$

where $v_{\mathbf{m}}^{(n)}(\vec{\kappa})$ are Fourier coefficients. By construction, $v_{\mathbf{m}}^{(n)}(\vec{\kappa}) = 0$, when $\mathbf{m} \notin \Omega(r_{n-1})$. Let $\mathbf{v}^{(n)}(\vec{\kappa})$ be the vector in $\ell^2(\mathbb{Z}^2)$ with components equal to $v_{\mathbf{m}}^{(n)}(\vec{\kappa})$. Note that the size of $\Omega(r_{n-1})$ depend on $\kappa = |\vec{\kappa}|$; to stress this fact we will use here the notations $\Omega(r_{n-1}, \kappa)$ and $r_{n-1}(\kappa)$. The Fourier transform $\widehat{\Psi}_n$ is a combination of δ -functions:

$$\widehat{\Psi}_n(\vec{\kappa}, \vec{\xi}) = 2\pi \sum_{\mathbf{m} \in \mathbb{Z}^2} v_{\mathbf{m}}^{(n)}(\vec{\kappa}) \delta(\vec{\xi} + \vec{\kappa} + \vec{p}_{\mathbf{m}})$$

From this, we easily compute the Fourier transform of $(S_n f)(\vec{x})$:

$$(\widehat{S_n f})(\vec{\xi}) = \sum_{\mathbf{m} \in \mathbb{Z}^2} v_{\mathbf{m}}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}}) f(-\vec{\xi} - \vec{p}_{\mathbf{m}}) \chi(\mathcal{G}'_n, -\vec{\xi} - \vec{p}_{\mathbf{m}}),$$

where $\chi(\mathcal{G}'_n, \cdot)$ is the characteristic function of $\mathcal{G}'_n$. Note that $v_{\mathbf{m}}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}}) \chi(\mathcal{G}'_n, -\vec{\xi} - \vec{p}_{\mathbf{m}})$ can differ from zero only when $\mathbf{m} \in \Omega(r_{n-1}, |\vec{\xi} + \vec{p}_{\mathbf{m}}|) \subset \Omega(r_{n-1}, \xi_{**})$, $\xi_{**} = \sup_{\vec{\xi} \in \mathcal{G}'_n} |\vec{\xi}|$. By Parseval's identity,

$$\begin{aligned} (S_n f, S_n g)_{L_2(\mathbb{R}^2)} &= (\widehat{S_n f}, \widehat{S_n g})_{L_2(\mathbb{R}^2)} = \\ &= \int_{\mathbb{R}^2} \sum_{\mathbf{m}, \mathbf{m}' \in \mathbb{Z}^2} T_{\mathbf{m}, \mathbf{m}'}(\vec{\xi}) f(-\vec{\xi} - \vec{p}_{\mathbf{m}}) \bar{g}(-\vec{\xi} - \vec{p}_{\mathbf{m}'}) \chi(\mathcal{G}'_n, -\vec{\xi} - \vec{p}_{\mathbf{m}}) \chi(\mathcal{G}'_n, -\vec{\xi} - \vec{p}_{\mathbf{m}'}) d\vec{\xi}, \\ T_{\mathbf{m}, \mathbf{m}'}(\vec{\xi}) &:= v_{\mathbf{m}}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}}) \overline{v_{\mathbf{m}'}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}'})}. \end{aligned}$$

Note that, in fact, the summation here is over the finite set $\mathbf{m}, \mathbf{m}' \in \Omega(r_{n-1}, \xi_{**})$. Hence we can exchange summation and integration in the above formula. Next, shifting the variable $\vec{\xi} + \vec{p}_{\mathbf{m}} \rightarrow \vec{\xi}$, denoting $\mathbf{m}' - \mathbf{m}$ by $\mathbf{m}''$ and considering that $\langle \mathbf{v}^{(n)}, \mathbf{v}^{(n)} \rangle = 1$, we obtain:

$$\begin{aligned} (\widehat{S_n f}, \widehat{S_n g})_{L_2(\mathbb{R}^2)} &= (f, g)_{L_2(\mathcal{G}'_n)} + \\ &= \sum_{\mathbf{m}'' \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} \int_{\mathbb{R}^2} B_{\mathbf{m}''}(\vec{\xi}) f(-\vec{\xi}) \bar{g}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) \chi(\mathcal{G}'_n, -\vec{\xi}) \chi(\mathcal{G}'_n, -\vec{\xi} - \vec{p}_{\mathbf{m}''}) d\vec{\xi}, \quad (379) \\ B_{\mathbf{m}''}(\vec{\xi}) &= \sum_{\mathbf{m} \in \mathbb{Z}^2} v_{\mathbf{m}}^{(n)}(-\vec{\xi}) \overline{v_{\mathbf{m}+\mathbf{m}''}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})}. \end{aligned}$$

Obviously,

$$B_{\mathbf{m}''} = \langle \mathbf{v}^{(n)}(-\vec{\xi}), \mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) \rangle, \quad (380)$$

where $\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})$ the “shifted” eigenvector: $\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) = \left(\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) \right)_{\mathbf{m}} = v_{\mathbf{m}+\mathbf{m}''}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})$. To obtain (376), it is enough to prove two estimates:

$$\sum_{\|\vec{p}_{\mathbf{m}''}\| > \xi_*^{r_{n-3}(\xi_*)}} \sup_{\vec{\xi} \in \mathcal{G}'_n} |B_{\mathbf{m}''}(\vec{\xi})| < \frac{1}{2} \xi_*^{-r_{n-3}(\xi_*)}, \quad (381)$$

$$\sum_{0 < \|\vec{p}_{\mathbf{m}''}\| \leq \xi_*^{r_{n-3}(\xi_*)}} \sup_{\vec{\xi} \in \mathcal{G}'_n} |B_{\mathbf{m}''}(\vec{\xi})| < \frac{1}{2} \xi_*^{-r_{n-3}(\xi_*)}. \quad (382)$$

To prove (381) we first check that

$$\sup_{\vec{\xi} \in \mathcal{G}'_n} |B_{\mathbf{m}''}(\vec{\xi})| < \|\vec{p}_{\mathbf{m}''}\|^{-8} \quad \text{when} \quad \|\vec{p}_{\mathbf{m}''}\| > \xi_*^{r_{n-3}(\xi_*)}. \quad (383)$$

Indeed, for every $\mathbf{m}''$ we break $\mathcal{G}'_n$ into several parts, partition being dependent on $\mathbf{m}''$:
 $\mathcal{G}'_n = \cup_{s,s'=0}^n \mathcal{G}_{ss'}$,

$$\begin{aligned} \mathcal{G}_{ss'} = & \left\{ \vec{\xi} \in \mathcal{G}'_n : |\vec{\xi}|^{r_{s-1}(|\vec{\xi}|)} \leq \frac{1}{2} \|\vec{p}_{\mathbf{m}''}\| < \gamma_s |\vec{\xi}|^{r_s(|\vec{\xi}|)} \right\} \cap \\ & \left\{ \vec{\xi} \in \mathcal{G}'_n : |\vec{\xi} + \vec{p}_{\mathbf{m}''}|^{r_{s'-1}(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|)} \leq \frac{1}{2} \|\vec{p}_{\mathbf{m}''}\| < \gamma_{s'} |\vec{\xi} + \vec{p}_{\mathbf{m}''}|^{r_{s'}(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|)} \right\}, \end{aligned} \quad (384)$$

where $r_{-1} := 0$, $r_0 = \delta$, $\gamma_s = 1$ when $s < n$, $\gamma_n = \infty$. To prove (383), it is enough to show

$$\sup_{\vec{\xi} \in \mathcal{G}_{ss'}} |B_{\mathbf{m}''}(\vec{\xi})| < \|\vec{p}_{\mathbf{m}''}\|^{-8} \quad (385)$$

for all s, s' . Assume $s, s' = n$. It follows from (384) that for any $\mathbf{m} \in \mathbb{Z}^4$ either $v_{\mathbf{m}}^{(n)}(\vec{\xi})$ or $v_{\mathbf{m}+\mathbf{m}''}^{(n)}(\vec{\xi} + \vec{p}_{\mathbf{m}''})$ is zero. Hence, $\langle \mathbf{v}^{(n)}(-\vec{\xi}), \mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) \rangle = 0$, i.e., $B_{\mathbf{m}''}(\vec{\xi}) = 0$. Next, let $0 < s < n$, $s' = n$. By (337),

$$\|\mathbf{v}^{(n)}(\vec{\xi}) - \mathbf{v}^{(s)}(\vec{\xi})\| < |\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{2}r_{s-1}(|\vec{\xi}|)}}. \quad (386)$$

It follows from the definition of $\mathcal{G}_{sn}$ that $\langle \mathbf{v}^{(s)}(\vec{\xi}), \mathbf{v}_*^{(n)}(\vec{\xi} + \vec{p}_{\mathbf{m}''}) \rangle = 0$. Therefore,

$$|B_{\mathbf{m}''}(\vec{\xi})| \leq \|\mathbf{v}^{(n)}(\vec{\xi}) - \mathbf{v}^{(s)}(\vec{\xi})\| < |\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{2}r_{s-1}(|\vec{\xi}|)}} \quad \text{when } \vec{\xi} \in \mathcal{G}_{sn}. \quad (387)$$

Using (107), (186) and (302), we obtain $|B_{\mathbf{m}''}(\vec{\xi})| \leq |\vec{\xi}|^{-10r_s(|\vec{\xi}|)}$. Considering again the definition of $\mathcal{G}_{sn}$, we get (385). Next, we consider $\mathcal{G}_{0n}$. By (46), $\mathbf{v}^{(1)} = \mathbf{v}^{(0)} + O(|\vec{\xi}|^{-2l+1+44\mu\delta})$, where $v_{\mathbf{m}}^{(0)} = \delta_{\mathbf{m},0}$. By (384), $\langle \mathbf{v}^{(0)}(\vec{\xi}), \mathbf{v}^{(n)}(\vec{\xi} + \vec{p}_{\mathbf{m}''}) \rangle = 0$. Hence, $|B_{\mathbf{m}''}(\vec{\xi})| \leq C|\vec{\xi}|^{-2l+1+44\mu\delta}$. Using again the definition of $\mathcal{G}_{0n}$ and the inequality $2l - 1 - 44\mu\delta > 8\delta$, we obtain (385). The case $s' < n$ is considered in the analogous way. Thus, (385) is proved. Summarizing (385) over $\mathbf{m}''$, we obtain (381).

Suppose $0 < \|\vec{p}_{\mathbf{m}''}\| \leq \xi_*^{r_{n-3}(\xi_*)}$. Let us estimate $B_{\mathbf{m}''}(\vec{\xi})$. Assume for definiteness that $|\vec{\xi} + \vec{p}_{\mathbf{m}''}| \leq |\vec{\xi}|$. The case of the opposite inequality is analogous up to the change of the notation $\vec{\xi} \rightarrow \vec{\xi} + \vec{p}_{-\mathbf{m}''}$, since $B_{\mathbf{m}''}(\vec{\xi}) = B_{-\mathbf{m}''}(\vec{\xi} + \vec{p}_{\mathbf{m}''})$. By (378),

$$H^{(n)}(-\vec{\xi}) \mathbf{v}^{(n)}(-\vec{\xi}) = \lambda^{(n)}(-\vec{\xi}) \mathbf{v}^{(n)}(-\vec{\xi}). \quad (388)$$

The analogous relation holds for $\mathbf{v}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})$ up to the replacement of $H^{(n)}(-\vec{\xi})$ by $H^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})$ and $\lambda^{(n)}(-\vec{\xi})$ by $\lambda^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})$:

$$H^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) \mathbf{v}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) = \lambda^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) \mathbf{v}^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}). \quad (389)$$

Note that $H^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})$ up to the shift of indices by $-\mathbf{m}''$ is equivalent to the operator $P_{\mathbf{m}''}H(-\vec{\xi})P_{\mathbf{m}''}$, where $P_{\mathbf{m}''}$ is the projection onto the box of the size $|\vec{\xi} + \vec{p}_{\mathbf{m}''}|^{r_{n-1}(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|)}$ around $-\mathbf{m}''$. Using the shifted eigenvector $\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})$, we can rewrite (389) in the form:

$$P_{\mathbf{m}''}H(-\vec{\xi})P_{\mathbf{m}''}\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) = \lambda^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}), \quad (390)$$

where $P_{\mathbf{m}''}\mathbf{v}_*^{(n)} = \mathbf{v}_*^{(n)}$. By (356),

$$H(-\vec{\xi})\mathbf{v}^{(n)}(-\vec{\xi}) = \lambda^{(n)}(-\vec{\xi})\mathbf{v}^{(n)}(-\vec{\xi}) + O\left(|\xi|^{-|\xi|^{\frac{1}{2}r_{n-1}(|\vec{\xi}|)}}\right). \quad (391)$$

Similarly,

$$\begin{aligned} H(-\vec{\xi})\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) = \\ \lambda^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) + O\left(|\xi + \vec{p}_{\mathbf{m}''}|^{-|\xi + \vec{p}_{\mathbf{m}''}|^{\frac{1}{2}r_{n-1}(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|)}}\right). \end{aligned} \quad (392)$$

Assume first $|\vec{\xi} + \vec{p}_{\mathbf{m}''}| \leq \frac{1}{2}|\vec{\xi}|$. Then $|\lambda^{(n)}(-\vec{\xi}) - \lambda^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''})| > \frac{1}{2}|\vec{\xi}|^{2l}$. Using (380), (391) and (392), we obtain:

$$B_{\mathbf{m}''} = O\left(\xi_*^{-\xi_*^{\frac{1}{2}r_{n-1}(\xi_*)}}\right) \quad \text{when } |\vec{\xi} + \vec{p}_{\mathbf{m}''}| \leq \frac{1}{2}|\vec{\xi}|. \quad (393)$$

Similar, but somewhat more subtle considerations are required when $|\vec{\xi} + \vec{p}_{\mathbf{m}''}| > \frac{1}{2}|\vec{\xi}|$. We start with introducing a parameter s . We will use it to cut $\Omega(r_{n-1}, |\vec{\xi}|)$ to approximately the same size as $\Omega(r_{n-1}, |\vec{\xi} + \vec{p}_{\mathbf{m}''}|)$. If the boxes are of approximately the same size, then $s = n - 1$. Indeed, for each $\vec{\xi}$ one of the following relations holds:

$$|\vec{\xi}|^{r_{s-1}(|\vec{\xi}|)} \leq |\vec{\xi} + \vec{p}_{\mathbf{m}''}|^{r_{n-1}(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|)} < |\vec{\xi}|^{r_s(|\vec{\xi}|)}, \quad (394)$$

where $1 \leq s \leq n - 1$ and s is defined by $\mathbf{m}''$ and $\vec{\xi}$. Note that $s < n - 1$ when $\Omega(r_{n-1}, |\vec{\xi}|)$ is essentially bigger than $\Omega(r_{n-1}, |\vec{\xi} + \vec{p}_{\mathbf{m}''}|)$. Using the second inequality in (394) and (302), we get

$$|\vec{\xi} + \vec{p}_{\mathbf{m}''}|^{r_{n-3}(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|)} < \frac{1}{8}|\vec{\xi}|^{r_{s-1}(|\vec{\xi}|)}. \quad (395)$$

Let $P_{\mathbf{0}s}$ be the projecting corresponding to $\Omega(r_s, |\vec{\xi}|)$. By (337) with s instead of n ,

$$(I - P_{\mathbf{0}s})\mathbf{v}^{(n)}(-\vec{\xi}) = O\left(|\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{2}r_{s-1}(|\vec{\xi}|)}}\right). \quad (396)$$

Let us prove the analogous estimate for $\mathbf{v}_*^{(n)}$:

$$(I - P_{\mathbf{0}s})\mathbf{v}_*^{(n)}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) = O\left(|\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{2}r_{s-1}(|\vec{\xi}|)}}\right). \quad (397)$$

Indeed, if $(P_{\mathbf{0}s})_{\mathbf{m}\mathbf{m}} = 0$, then $\|\vec{p}_{\mathbf{m}}\| > |\vec{\xi}|^{r_{s-1}(|\vec{\xi}|)}$. Using (395) and the bound on $\|\vec{p}_{\mathbf{m}''}\|$, we obtain $\|\vec{p}_{\mathbf{m}+\mathbf{m}''}\| > \frac{1}{2}|\vec{\xi}|^{r_{s-1}(|\vec{\xi}|)}$. Using (337), (338), we obtain (397). From (391),(392), considering that $\|P_{\mathbf{0}s}H\| = O\left(|\xi|^{2lr_{s-1}(|\vec{\xi}|)}\right)$ and using (396),(397), we get:

$$H^{(s)}(-\vec{\xi})P_{\mathbf{0}s}\mathbf{v}^{(n)}(-\vec{\xi}) = \lambda^{(n)}(-\vec{\xi})P_{\mathbf{0}s}\mathbf{v}^{(n)}(-\vec{\xi}) + O\left(|\xi|^{-|\xi|^{\frac{1}{4}r_{s-1}}}\right), \quad (398)$$

$$H^{(s)}(-\vec{\xi})P_{\mathbf{0}s}\mathbf{v}_*^{(n)}(-\vec{\xi}-\vec{p}_{\mathbf{m}''}) = \lambda^{(n)}(-\vec{\xi}-\vec{p}_{\mathbf{m}''})P_{\mathbf{0}s}\mathbf{v}_*^{(n)}(-\vec{\xi}-\vec{p}_{\mathbf{m}''}) + O\left(|\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{4}r_{s-1}}}\right). \quad (399)$$

Next, by Theorem 7.17 for step s , $\left|\lambda^{(n)}(-\vec{\xi}) - \lambda^{(n)}(-\vec{\xi}-\vec{p}_{\mathbf{m}''})\right| > \varepsilon_0^{(s)}/2$, where $\varepsilon_0^{(s)} = |\vec{\xi}|^{-2r'_{s-1}|\vec{\xi}|^{2\gamma r_{s-2}}}$. Indeed, $\mathbf{v}^{(n)}(-\vec{\xi})$ and $\mathbf{v}_*^{(n)}(-\vec{\xi}-\vec{p}_{\mathbf{m}''})$ are almost orthogonal since they are concentrated around $\mathbf{m} = \mathbf{0}$ and $\mathbf{m} = \mathbf{m}'' \neq \mathbf{0}$ respectively; thus $\lambda^{(n)}(-\vec{\xi}-\vec{p}_{\mathbf{m}''})$ must be outside of the interval described in Theorem 7.17, while $\lambda^{(n)}(-\vec{\xi})$ is inside twice shorter interval. Now, using (398) and (399), we obtain:

$$\langle P_{\mathbf{0}s}\mathbf{v}^{(n)}(-\vec{\xi}), P_{\mathbf{0}s}\mathbf{v}_*^{(n)}(-\vec{\xi}-\vec{p}_{\mathbf{m}''}) \rangle = O\left(|\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{4}r_{s-1}}}\right) |\vec{\xi}|^{2r'_{s-1}|\vec{\xi}|^{2\gamma r_{s-2}}} = O\left(|\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{8}r_{s-1}}}\right),$$

see (302). Using one more time (396), (397), and considering (395), we obtain $B_{\mathbf{m}''} = O\left(\xi_*^{-\xi_*^{\frac{1}{8}r_{n-3}(\xi_*)}}\right)$ for the case $|\vec{\xi} + \vec{p}_{\mathbf{m}''}| > \frac{1}{2}|\vec{\xi}|$. Using this estimate together with (393) and considering that the number of $\mathbf{m}''$ satisfying $0 < \|\vec{p}_{\mathbf{m}''}\| \leq \xi_*^{r_{n-3}(\xi_*)}$ does not exceed $16\xi_*^{4r_{n-3}(\xi_*)}$, we obtain (382). Substituting the estimates for $B_{\mathbf{m}''}$ into (379), we obtain (376).

□

It is easy to see that $T_n \subset S_n^*$. Therefore, $\|T_n\| \leq 1 + o(1)$ and can be extended to the whole space $L_2(\mathcal{G}_n)$. We still denote the extended operator by T_n , $T_n = S_n^*$. Therefore, E_n is a self-adjoint operator.

Lemma 9.4. *Let $\mathcal{G}'_n \subset \mathcal{G}''_n \subset \mathcal{G}_n$. The following relation holds as $n \rightarrow \infty$:*

$$E_n(\mathcal{G}'_n)E_n(\mathcal{G}''_n) = E_n(\mathcal{G}'_n) + o(1), \quad (400)$$

where $o(1)$ is taken in the space of bounded operators and uniform in $\mathcal{G}'_n, \mathcal{G}''_n$.

Corollary 9.5. $E_n(\mathcal{G}''_n)E_n(\mathcal{G}'_n) = E_n(\mathcal{G}'_n) + o(1)$.

This corollary is valid, since E_n is selfadjoint.

Corollary 9.6. $E_n^2(\mathcal{G}'_n) = E_n(\mathcal{G}'_n) + o(1)$ for any $\mathcal{G}'_n \subset \mathcal{G}_n$.

Proof. Let $I_n(\mathcal{G}'_n)$ be the projection from $L_2(\mathcal{G}''_n)$ to $L_2(\mathcal{G}'_n)$. It is easy to see that $T_n(\mathcal{G}'_n) = I_n(\mathcal{G}'_n)T_n(\mathcal{G}''_n)$. Hence, $T_n(\mathcal{G}'_n)S_n(\mathcal{G}''_n) = I_n(\mathcal{G}'_n)T_n(\mathcal{G}''_n)S_n(\mathcal{G}''_n)$. By (376) for set $\mathcal{G}''_n$, $T_n(\mathcal{G}''_n)S_n(\mathcal{G}''_n) = id(\mathcal{G}''_n) + o(1)$, where $id(\mathcal{G}''_n)$ is the identity in $L_2(\mathcal{G}''_n)$. It immediately follows $T_n(\mathcal{G}'_n)S_n(\mathcal{G}''_n) = I_n(\mathcal{G}'_n) + o(1)$. Substituting the last relation into the formula $E_n(\mathcal{G}'_n)E_n(\mathcal{G}''_n) = S_n(\mathcal{G}'_n)T_n(\mathcal{G}'_n)S_n(\mathcal{G}''_n)T_n(\mathcal{G}''_n)$, we obtain (400). $\square$

Let

$$\mathcal{G}_{n,\lambda} = \{\vec{x} \in \mathcal{G}_n : \lambda^{(n)}(\vec{x}) < \lambda\}. \quad (401)$$

This set is Lebesgue measurable, since $\mathcal{G}_n$ is open and $\lambda^{(n)}(\vec{x})$ is continuous on $\mathcal{G}_n$.

Lemma 9.7. $|\mathcal{G}_{n,\lambda+\varepsilon} \setminus \mathcal{G}_{n,\lambda}| \leq 2\pi\lambda^{-1+\frac{1}{t}}\varepsilon$ when $0 \leq \varepsilon \leq 1$.

The proof is based on Lemma 7.21 and completely analogous to that of Lemma 2 in [28].

By (372), $E_n(\mathcal{G}_{n,\lambda+\varepsilon}) - E_n(\mathcal{G}_{n,\lambda}) = E_n(\mathcal{G}_{n,\lambda+\varepsilon} \setminus \mathcal{G}_{n,\lambda})$. Let us obtain an estimate for this projection.

Lemma 9.8. For any $F \in C_0^\infty(\mathbb{R}^2)$ and $0 \leq \varepsilon \leq 1$,

$$\|(E_n(\mathcal{G}_{n,\lambda+\varepsilon}) - E_n(\mathcal{G}_{n,\lambda}))F\|_{L_2(\mathbb{R}^2)}^2 \leq C(F)\lambda^{-1+\frac{1}{t}}\varepsilon, \quad (402)$$

where $C(F)$ is uniform with respect to n and λ .

Proof. Let $\mathcal{G}'_n = \mathcal{G}_{n,\lambda+\varepsilon} \setminus \mathcal{G}_{n,\lambda}$. Using the definition (373) of E_n and formula (377) with $f = g = T_n F$, we obtain

$$\|E_n(\mathcal{G}'_n)F\|_{L_2(\mathbb{R}^2)}^2 < (1 + o(1))\|T_n F\|_{L_\infty(\mathcal{G}'_n)}^2 |\mathcal{G}'_n|. \quad (403)$$

Using (374) and Corollary 8.8 we easily get $\|T_n F\|_{L_\infty(\mathcal{G}'_n)} < 2\|F\|_{L_1(\mathbb{R}^2)}$. Substituting this estimate into (403) and using Lemma 9.7, we obtain (402). $\square$

9.2 Sets $\mathcal{G}_\infty$ and $\mathcal{G}_{\infty,\lambda}$

By construction, $\mathcal{G}_{n+1} \subset \mathcal{G}_n$, $\mathcal{G}_\infty = \bigcap_{n=1}^\infty \mathcal{G}_n$. Therefore, the perturbation formulas for $\lambda^{(n)}(\vec{x})$ and $\Psi_n(\vec{x})$ hold in $\mathcal{G}_\infty$ for all n . Let

$$\mathcal{G}_{\infty,\lambda} = \{\vec{x} \in \mathcal{G}_\infty : \lambda_\infty(\vec{x}) < \lambda\}. \quad (404)$$

The function $\lambda_\infty(\vec{x})$ is a Lebesgue measurable function, since it is a limit of the sequence of measurable functions. Hence, the set $\mathcal{G}_{\infty,\lambda}$ is measurable.

Lemma 9.9. The measure of the symmetric difference of two sets $\mathcal{G}_{\infty,\lambda}$ and $\mathcal{G}_{n,\lambda}$ converges to zero as $n \rightarrow \infty$ uniformly in λ in every bounded interval:

$$\lim_{n \rightarrow \infty} |\mathcal{G}_{\infty,\lambda} \Delta \mathcal{G}_{n,\lambda}| = 0.$$

The proof is completely analogous to the proof of Lemma 4 in [28].

9.3 Projections $E(\mathcal{G}_{\infty,\lambda})$

In this section, we show that the operators $E_n(\mathcal{G}_{\infty,\lambda})$ have a strong limit $E_\infty(\mathcal{G}_{\infty,\lambda})$ in $L_2(\mathbb{R}^2)$ as n tends to infinity. The operator $E_\infty(\mathcal{G}_{\infty,\lambda})$ is a spectral projection of H . It can be represented in the form $E_\infty(\mathcal{G}_{\infty,\lambda}) = S_\infty T_\infty$, where S_∞ and T_∞ are strong limits of $S_n(\mathcal{G}_{\infty,\lambda})$ and $T_n(\mathcal{G}_{\infty,\lambda})$, respectively. For any $F \in C_0^\infty(\mathbb{R}^2)$, we show:

$$E_\infty(\mathcal{G}_{\infty,\lambda}) F = \frac{1}{4\pi^2} \int_{\mathcal{G}_{\infty,\lambda}} (F, \Psi_\infty(\vec{x})) \Psi_\infty(\vec{x}, \vec{x}) d\vec{x}, \quad (405)$$

$$H E_\infty(\mathcal{G}_{\infty,\lambda}) F = \frac{1}{4\pi^2} \int_{\mathcal{G}_{\infty,\lambda}} \lambda_\infty(\vec{x}) (F, \Psi_\infty(\vec{x})) \Psi_\infty(\vec{x}, \vec{x}) d\vec{x}. \quad (406)$$

Using properties of $E_\infty(\mathcal{G}_{\infty,\lambda})$, we prove absolute continuity of the branch of the spectrum corresponding to functions $\Psi_\infty(\vec{x})$.

We consider the sequence of operators $S_n(\mathcal{G}_{\infty,\lambda})$ which are given by (375) with $\mathcal{G}'_n = \mathcal{G}_{\infty,\lambda}$.

Lemma 9.10. *We have*

$$\|(S_n(\mathcal{G}_{\infty,\lambda}) - S_{n-1}(\mathcal{G}_{\infty,\lambda}))f\|_{L_2(\mathbb{R}^2)} < C \|f\|_{L_2(\mathcal{G}_{\infty,\lambda})} \xi_*^{-\frac{1}{4} r_{n-2}(\xi_*)}. \quad (407)$$

Proof. Considering as in the proof of Lemma 9.1, we obtain

$$\begin{aligned} \|(S_n - S_{n-1})f\|_{L_2(\mathbb{R}^2)}^2 &= \left\| \widehat{S_n f} - \widehat{S_{n-1} f} \right\|_{L_2(\mathbb{R}^2)}^2 = \\ &= \int_{\mathbb{R}^2} \sum_{\mathbf{m}''} \tilde{B}_{\mathbf{m}''}(\vec{\xi}) f(-\vec{\xi}) \bar{f}(-\vec{\xi} - \vec{p}_{\mathbf{m}''}) \chi(\mathcal{G}_{\infty,\lambda}, -\vec{\xi}) \chi(\mathcal{G}_{\infty,\lambda}, -\vec{\xi} - \vec{p}_{\mathbf{m}''}) d\vec{\xi}, \end{aligned} \quad (408)$$

$$\begin{aligned} \tilde{B}_{\mathbf{m}''}(\vec{\xi}) &= \\ &= \sum_{\mathbf{m} \in \Omega(r_{n-1}, |\vec{\xi}|): \mathbf{m} + \mathbf{m}'' \in \Omega(r_{n-1}, |\vec{\xi} + \vec{p}_{\mathbf{m}''}|)} (v_{\mathbf{m}}^{(n)} - v_{\mathbf{m}}^{(n-1)}) (-\vec{\xi}) \overline{(v_{\mathbf{m}+\mathbf{m}''}^{(n)} - v_{\mathbf{m}+\mathbf{m}''}^{(n-1)})} (-\vec{\xi} - \vec{p}_{\mathbf{m}''}). \end{aligned}$$

Assume for definiteness that $|\vec{\xi} + \vec{p}_{\mathbf{m}''}| \leq |\vec{\xi}|$. If $\|\vec{p}_{\mathbf{m}''}\| > 2|\vec{\xi}|^{r_{n-1}(|\vec{\xi}|)}$, then $\tilde{B}_{\mathbf{m}''}(\vec{\xi}) = 0$. Let $\vec{\xi} : \|\vec{p}_{\mathbf{m}''}\| \leq 2|\vec{\xi}|^{r_{n-1}(|\vec{\xi}|)}$. Using (337) with n instead of $n+1$, we easily obtain:

$$\left| \tilde{B}_{\mathbf{m}''}(\vec{\xi}) \right| = O\left(|\vec{\xi}|^{-|\vec{\xi}|^{\frac{1}{2} r_{n-2}(|\vec{\xi}|)}}\right) O\left(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|^{-|\vec{\xi} + \vec{p}_{\mathbf{m}''}|^{\frac{1}{2} r_{n-2}(|\vec{\xi} + \vec{p}_{\mathbf{m}''}|)}}\right). \quad (409)$$

Considering (302) with $n-1$ instead of n and taking into account that $\|\vec{p}_{\mathbf{m}''}\| \leq 2|\vec{\xi}|^{r_{n-1}(|\vec{\xi}|)}$, we easily get:

$$\left| \tilde{B}_{\mathbf{m}''}(\vec{\xi}) \right| = \|\vec{p}_{\mathbf{m}''}\|^{-8} O\left(\xi_*^{-\frac{1}{2} r_{n-2}(\xi_*)}\right).$$

Summarizing the last estimate for $\mathbf{m}'' \neq \mathbf{0}$ and using (409) for $\mathbf{m}'' = \mathbf{0}$, we arrive at (407). $\square$

By (407), the sequence of operators $S_n(\mathcal{G}_{\infty,\lambda})$ is a Cauchy sequence in the space of bounded operators. We denote its limit by $S_\infty(\mathcal{G}_{\infty,\lambda})$. Note that the convergence of $S_n(\mathcal{G}_{\infty,\lambda})$ to $S_\infty(\mathcal{G}_{\infty,\lambda})$ is uniform in λ when $\lambda > \lambda_*$.

Lemma 9.11. *The operator $S_\infty(\mathcal{G}_{\infty,\lambda})$ can be described by the formula*

$$(S_\infty f)(\vec{x}) = \frac{1}{2\pi} \int_{\mathcal{G}_{\infty,\lambda}} f(\vec{z}) \Psi_\infty(\vec{z}, \vec{x}) d\vec{z} \quad (410)$$

for any $f \in L_\infty(\mathcal{G}_{\infty,\lambda})$.

Proof. From Theorem 8.9 it follows that for every $f \in L_\infty(\mathcal{G}_{\infty,\lambda})$

$$\lim_{n \rightarrow \infty} \int_{\mathcal{G}_{\infty,\lambda}} f(\vec{z}) \Psi_n(\vec{z}, \vec{x}) d\vec{z} = \int_{\mathcal{G}_{\infty,\lambda}} f(\vec{z}) \Psi_\infty(\vec{z}, \vec{x}) d\vec{z} \quad (411)$$

for all $\vec{x}$. Hence, (410) holds. $\square$

Now we consider the sequence of operators $T_n(\mathcal{G}_{\infty,\lambda})$ which are given by (374) and act from $L_2(\mathbb{R}^2)$ to $L_2(\mathcal{G}_{\infty,\lambda})$. Since, $T_n = S_n^*$, the sequence has a limit T_∞ in the class of bounded operators, $T_\infty = S_\infty^*$. Note that the convergence of $T_n(\mathcal{G}_{\infty,\lambda})$ to $T_\infty(\mathcal{G}_{\infty,\lambda})$ is uniform in λ when $\lambda > \lambda_*$.

Lemma 9.12. *The operator $T_\infty(\mathcal{G}_{\infty,\lambda})$ can be described by the formula $T_\infty(\mathcal{G}_{\infty,\lambda})F = \frac{1}{2\pi}(F, \Psi_\infty(\vec{z}))$ for any $F \in C_0^\infty(\mathbb{R}^2)$.*

Proof. The lemma easily follows from Theorem 8.9 and formula (374). $\square$

Lemma 9.13. *Operators $E_n(\mathcal{G}_{\infty,\lambda})$ have a limit $E_\infty(\mathcal{G}_{\infty,\lambda})$ in the class of bounded operators in $L_2(\mathbb{R}^2)$, the convergence being uniform for $\lambda > \lambda_*$. The operator $E_\infty(\mathcal{G}_{\infty,\lambda})$ is a projection. For any $F \in C_0^\infty(\mathbb{R}^2)$ it is given by (405).*

Proof. The lemma immediately follows from convergence of sequences S_n , T_n and Lemmas 9.4, 9.11, 9.12. $\square$

Lemma 9.14. *There is a strong limit $E_\infty(\mathcal{G}_\infty)$ of the projections $E_\infty(\mathcal{G}_{\infty,\lambda})$ as λ goes to infinity.*

Corollary 9.15. *The operator $E_\infty(\mathcal{G}_\infty)$ is a projection.*

Proof. It can be easily seen from (405) that the sequence of $E_\infty(\mathcal{G}_{\infty,\lambda})$ is monotonuos in λ . It is well known that a monotone sequence of projectors has a limit. $\square$

The proofs of the next two lemmas are completely analogous to the proofs of Lemmas 10, 11 in [28].

Lemma 9.16. *Projections $E_\infty(\mathcal{G}_{\infty,\lambda})$, $\lambda \in \mathbb{R}$, and $E_\infty(\mathcal{G}_\infty)$ reduce the operator H .*

Lemma 9.17. *The family of projections $E_\infty(\mathcal{G}_{\infty,\lambda})$ is the resolution of the identity of the operator $HE_\infty(\mathcal{G}_\infty)$ acting in $E_\infty(\mathcal{G}_\infty)L_2(\mathbb{R}^2)$.*

Lemma 9.18. *Formula (406) holds, when $F \in C_0^\infty(\mathbb{R}^2)$.*

Proof. By the previous lemma, $E_\infty(\mathcal{G}_{\infty,\lambda})F \in D(H)$. It is easy to see that the r.h.s. of (405) can be differentiated with respect to $\vec{x}$ under the integral sign. Now, considering (364), we get (406). $\square$

9.4 Proof of Absolute Continuity

Now we show that the branch of spectrum (semi-axis) corresponding to $\mathcal{G}_\infty$ is absolutely continuous.

Theorem 9.19. *For any $F \in C_0^\infty(\mathbb{R}^2)$ and $0 \leq \varepsilon \leq 1$,*

$$|(E_\infty(\mathcal{G}_{\infty,\lambda+\varepsilon})F, F) - (E_\infty(\mathcal{G}_{\infty,\lambda})F, F)| \leq C_F \lambda^{-1+\frac{1}{l}} \varepsilon. \quad (412)$$

Corollary 9.20. *The spectrum of the operator $HE_\infty(\mathcal{G}_\infty)$ is absolutely continuous.*

Proof. By formula (405),

$$|(E_\infty(\mathcal{G}_{\infty,\lambda+\varepsilon})F, F) - (E_\infty(\mathcal{G}_{\infty,\lambda})F, F)| \leq C_F |\mathcal{G}_{\infty,\lambda+\varepsilon} \setminus \mathcal{G}_{\infty,\lambda}|.$$

Applying Lemmas 9.7 and 9.9, we immediately get (412). $\square$

10 Appendices

10.1 Appendix 1. Proof of Lemma 3.12

Proof.

1. The case $p_{\mathbf{m}} > 4k$. From (70) it immediately follows that $|\Im \varphi_{\mathbf{m}}^\pm| > (\cosh)^{-1} 2 > 1$. Hence, $\mathcal{W}_0 \cap \mathcal{O}_{\mathbf{m}}(k, \tau) = \emptyset$.

Further we use the Taylor series for $|\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2$ near its zeros: Noting that

$$|\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 = 2kp_{\mathbf{m}} \cos(\varphi - \varphi_{\mathbf{m}}) + p_{\mathbf{m}}^2 \quad (413)$$

and recalling that $\varphi_{\mathbf{m}}^{\pm}$ are the solutions of $|\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 = k^2$, we see:

$$\cos(\varphi_{\mathbf{m}}^{\pm} - \varphi_{\mathbf{m}}) = -\frac{p_{\mathbf{m}}}{2k}, \quad |\sin(\varphi_{\mathbf{m}}^{\pm} - \varphi_{\mathbf{m}})| = \sqrt{\left|1 - \frac{p_{\mathbf{m}}^2}{4k^2}\right|}. \quad (414)$$

Expanding (413) around $\varphi_{\mathbf{m}}^{\pm}$, we get:

$$\begin{aligned} & |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 = \\ & -2kp_{\mathbf{m}}\sin(\varphi_{\mathbf{m}}^{\pm} - \varphi_{\mathbf{m}})r_{\mathbf{m}}(1 + O(r_{\mathbf{m}}^2)) + kp_{\mathbf{m}}\cos(\varphi_{\mathbf{m}}^{\pm} - \varphi_{\mathbf{m}})r_{\mathbf{m}}^2(1 + O(r_{\mathbf{m}}^2)), \end{aligned} \quad (415)$$

where $r_{\mathbf{m}} = |\varphi - \varphi_{\mathbf{m}}^{\pm}|$.

2. In the second case we put $r_{\mathbf{m}} = \frac{\tau k^{-1-40\mu\delta}}{p_{\mathbf{m}}\sqrt{\left|1 - \frac{p_{\mathbf{m}}^2}{4k^2}\right|}}(1 + o(1))$ when $k^{-1-39\mu\delta} < p_{\mathbf{m}} < 4k$ and $\left|1 - \frac{p_{\mathbf{m}}^2}{4k^2}\right| > \tau k^{-2-40\mu\delta}$. Substituting $r_{\mathbf{m}}$ into (415), we get that the modulus of the first term is $2\tau k^{-40\mu\delta}(1+o(1))$ and that of the second term is $\frac{\tau^2 k^{-2-80\delta}}{2\left|1 - \frac{p_{\mathbf{m}}^2}{4k^2}\right|}(1+o(1))$.

Using the condition $\left|1 - \frac{p_{\mathbf{m}}^2}{4k^2}\right| > \tau k^{-2-40\mu\delta}$, one can easily see that the former is at least twice greater than the latter. Thus, we get

$$\left| |\vec{k}(\varphi) + \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^2 - k^2 \right| > \tau k^{-40\mu\delta} \quad \text{when } |\varphi - \varphi_{\mathbf{m}}^{\pm}| = r_{\mathbf{m}}. \quad (416)$$

Now, the maximum principle yield that this inequality holds everywhere outside the discs $\cup_{\pm, j \in \mathbb{Z}} (\Phi_{\mathbf{m}}^{\pm} + 2\pi j)$. Hence, $\mathcal{O}_{\mathbf{m}} \subset \cup_{\pm, j \in \mathbb{Z}} (\Phi_{\mathbf{m}}^{\pm} + 2\pi j)$.

3. In the third case we put $r_{\mathbf{m}} = 32\tau k^{-1-20\mu\delta}(1+o(1))$ and $\left|1 - \frac{p_{\mathbf{m}}^2}{4k^2}\right| < \tau k^{-2-40\mu\delta}$. This time the modulus of the second term in (415) is $64 \cdot 32\tau^2 k^{-40\mu\delta}(1+o(1))$ and that of the first is smaller than $128\tau^{3/2} k^{-40\mu\delta}(1+o(1))$. Therefore we again have (416) and $\mathcal{O}_{\mathbf{m}} \subset \cup_{\pm, j \in \mathbb{Z}} (\Phi_{\mathbf{m}}^{\pm} + 2\pi j)$.

□

10.2 Appendix 2. Proof of Lemma 3.13

Proof. The proof is by contradiction. By definition of $\mathcal{M}_2$, $J \geq 1$. Suppose $J \geq 4$. We arrive to contradiction in several steps.

1. By definition of $\mathcal{M}'(\varphi_0)$, $\varphi_0 \in \cap_{j=0}^J \mathcal{O}_{\mathbf{m}_j}(k, 1)$. This means:

$$\left| |\vec{k} + \vec{p}_{\mathbf{m}_j}|^2 - k^2 \right| \leq k^{-40\mu\delta}, \quad j = 0, \dots, J$$

It follows:

$$\left| 2 \left(\vec{k}(\varphi) + \vec{p}_{\mathbf{m}_0}, \vec{p}_{\mathbf{q}_j} \right) + p_{\mathbf{q}_j}^2 \right| < ck^{-40\mu\delta}, \quad \mathbf{q}_j = \mathbf{m}_j - \mathbf{m}_0, \quad j = 1, \dots, J, \quad (417)$$

where $\|\vec{p}_{\mathbf{q}_j}\| \leq jk^\delta$. In this part of the proof we show that no two vectors $\vec{p}_{\mathbf{q}_j}$, $j = 1, 2, 3, 4$ are colinear. Indeed, suppose $\vec{p}_{\mathbf{q}_1}$ and $\vec{p}_{\mathbf{q}_2}$ are colinear. Let $\vec{v}$ be a unit vector in the direction of $\vec{p}_{\mathbf{q}_1}$. Then the directional vector of $\vec{p}_{\mathbf{q}_2}$ is $\pm\vec{v}$, where $\pm$ means $+$ or $-$. Inequality (417) together with the estimate (28) for $p_{\mathbf{q}_j}$ yields:

$$2(\vec{k}(\varphi) + \vec{p}_{\mathbf{m}_0}, \vec{v}) + p_{\mathbf{q}_1} = O(k^{-39\mu\delta}), \quad \pm 2(\vec{k}(\varphi) + \vec{p}_{\mathbf{m}_0}, \vec{v}) + p_{\mathbf{q}_2} = O(k^{-39\mu\delta}).$$

Hence,

$$p_{\mathbf{q}_1} \mp p_{\mathbf{q}_2} = O(k^{-39\mu\delta}). \quad (418)$$

Note that $\|\vec{p}_{\mathbf{q}_1} \mp \vec{p}_{\mathbf{q}_2}\| < 8k^\delta$. Therefore, $p_{\mathbf{q}_1 \mp \mathbf{q}_2} > ck^{-\mu\delta}$. Considering that $p_{\mathbf{q}_1 \mp \mathbf{q}_2} = |p_{\mathbf{q}_1} \mp p_{\mathbf{q}_2}|$, we arrive to contradiction with (418). Hence, no two vectors $\vec{p}_{\mathbf{q}_j}$, $j = 1, 2, 3, 4$ are colinear.

2. We represent every $\vec{p}_{\mathbf{q}_j}$ in the form: $\vec{p}_{\mathbf{q}_j}/2\pi = \mathbf{s}_j + \alpha\mathbf{s}'_j$, $\mathbf{s}_j, \mathbf{s}'_j \in \mathbb{Z}^2$, $|\mathbf{s}_j|, |\mathbf{s}'_j| < 8k^\delta$. Let us show that

$$[\mathbf{s}_1, \mathbf{s}'_1] \neq 0 \text{ when } J > 1, \quad (419)$$

$[\mathbf{a}, \mathbf{b}] = a_1b_2 - a_2b_1$. Indeed, suppose that $[\mathbf{s}_1, \mathbf{s}'_1] = 0$. Then, $\mathbf{s}_1, \mathbf{s}'_1$ are colinear integer vectors. Hence, there are integers n, m and an integer vector $\mathbf{s}''_1$, such that $\mathbf{s}_1 = n\mathbf{s}''_1$ and $\mathbf{s}'_1 = m\mathbf{s}''_1$, $0 < |n| + |m| < 8k^\delta$. Therefore, $\vec{p}_{\mathbf{q}_1}/2\pi = (n + \alpha m)\mathbf{s}''_1$. Hence, $[\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_2}](2\pi)^{-2} = (n + \alpha m)([\mathbf{s}''_1, \mathbf{s}_2] + \alpha[\mathbf{s}''_1, \mathbf{s}'_2])$. It follows from (417) that the angle (modulo π) between $\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_2}$ is less than $(p_{\mathbf{q}_1} + p_{\mathbf{q}_2})k^{-1}$. Hence,

$$[\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_2}] = O(k^{-1+3\delta}). \quad (420)$$

Therefore,

$$n + \alpha m = O(k^{-1/2+3\delta/2}) \quad \text{or} \quad [\mathbf{s}''_1, \mathbf{s}_2] + \alpha[\mathbf{s}''_1, \mathbf{s}'_2] = O(k^{-1/2+3\delta/2}). \quad (421)$$

The first relation is impossible since (with the same proof as for (28)) $|n + \alpha m| > 8C_\epsilon k^{-(\mu-1+\epsilon)\delta}$ for any $\epsilon > 0$ and $(2\mu + 3)\delta < 1$. Similarly, the second relation (421) is possible if and only if $[\mathbf{s}''_1, \mathbf{s}'_2] = [\mathbf{s}''_1, \mathbf{s}_2] = 0$. Therefore $\vec{p}_{\mathbf{q}_2}$ is colinear to $\mathbf{s}''_1$, i.e. to $\vec{p}_{\mathbf{q}_1}$. This means $\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_2}$ are colinear. It cannot be the case, as we proved before. Thus, we have arrived to (419).

3. Let us consider (420). Substituting $\vec{p}_{\mathbf{q}_j} = \mathbf{s}_j + \alpha\mathbf{s}'_j$, $j = 1, 2$, we obtain:

$$n_1 + \alpha p_1 + \alpha^2 m_1 = O(k^{-1+3\delta}), \quad (422)$$

where n_1, p_1, m_1 are integers, $n_1 = [\mathbf{s}_1, \mathbf{s}_2]$, $p_1 = [\mathbf{s}_1, \mathbf{s}'_2] + [\mathbf{s}'_1, \mathbf{s}_2]$, $m_1 = [\mathbf{s}'_1, \mathbf{s}'_2]$. Obviously, $n_1, p_1, m_1 = O(k^{2\delta})$. Note that $n_1^2 + p_1^2 + m_1^2 \neq 0$, since otherwise vectors

$\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_2}$ are colinear. Next, $m_1 \neq 0$, since otherwise $O(k^{-1+3\delta}) > c(\epsilon)k^{-2(\mu-1+\epsilon)\delta}$ for any $\epsilon > 0$. This cannot be true for our choice of δ .

Suppose there is another triple (n_2, p_2, m_2) , $0 < n_2^2 + p_2^2 + m_2^2 < 10^3 k^{4\delta}$, such that (422) holds. Namely,

$$n_2 + \alpha p_2 + \alpha^2 m_2 = O(k^{-1+3\delta}). \quad (423)$$

The goal of this part is to show that such (n_2, p_2, m_2) is, in fact, a multiple of (n_1, p_1, m_1) . Indeed, $m_2 \neq 0$ for the same reason as $m_1 \neq 0$. Excluding α^2 from (425) and (423), we get

$$(n_1 m_2 - n_2 m_1) + \alpha(p_1 m_2 - p_2 m_1) = O(k^{-1+5\delta}).$$

It follows that $p_1 m_2 - p_2 m_1 = 0$, $n_1 m_2 - n_2 m_1 = 0$, since otherwise $O(k^{-1+5\delta}) > c(\epsilon)k^{-4(\mu-1+\epsilon)\delta}$ for any $\epsilon > 0$. Thus, (n_2, p_2, m_2) is colinear to (n_1, p_1, m_1) and $(n_2, p_2, m_2) = \frac{r}{s}(n_1, p_1, m_1)$, where r, s are integers, $s \neq 0$ and $r, s = O(k^{2\delta})$.

4. In this part we show that

$$\vec{p}_{\mathbf{q}_3} = \frac{r}{s}\vec{p}_{\mathbf{q}_2} + \frac{t}{s}\vec{p}_{\mathbf{q}_1}, \quad (424)$$

where s, r, t are nonzero integers, $s, r, t = O(k^{3\delta})$. Indeed, let us consider the relation $[\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_3}] = O(k^{-1+3\delta})$. This relation follows from (417) the same way as (420). Substituting $\vec{p}_{\mathbf{q}_j} = \mathbf{s}_j + \alpha \mathbf{s}'_j$, $j = 1, 3$, we obtain:

$$n_2 + \alpha p_2 + \alpha^2 m_2 = O(k^{-1+3\delta}), \quad (425)$$

where n_2, p_2, m_2 are integers, $n_2 = [\mathbf{s}_1, \mathbf{s}_3]$, $p_2 = [\mathbf{s}_1, \mathbf{s}'_3] + [\mathbf{s}'_1, \mathbf{s}_3]$, $m_2 = [\mathbf{s}'_1, \mathbf{s}'_3]$. Note that $n_2^2 + p_2^2 + m_2^2 \neq 0$, since otherwise vectors $\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_3}$ are colinear. Therefore, by part 3, $(n_2, p_2, m_2) = \frac{r}{s}(n_1, p_1, m_1)$, where r, s are integers, $s, r \neq 0$, $s, r = O(k^{2\delta})$. This yields:

$$s[\mathbf{s}_1, \mathbf{s}_3] = r[\mathbf{s}_1, \mathbf{s}_2], \quad (426)$$

$$s([\mathbf{s}_1, \mathbf{s}'_3] + [\mathbf{s}'_1, \mathbf{s}_3]) = r([\mathbf{s}_1, \mathbf{s}'_2] + [\mathbf{s}'_1, \mathbf{s}_2]), \quad (427)$$

$$s[\mathbf{s}'_1, \mathbf{s}'_3] = r[\mathbf{s}'_1, \mathbf{s}'_2]. \quad (428)$$

Note that $|\mathbf{s}_1|, |\mathbf{s}'_1| \neq 0$, since $[\mathbf{s}_1, \mathbf{s}'_1] \neq 0$ by part 2. It follows from (426) and (428) that $s\mathbf{s}_3 - r\mathbf{s}_2$ is colinear to $\mathbf{s}_1$ and $s\mathbf{s}'_3 - r\mathbf{s}'_2$ is colinear to $\mathbf{s}'_1$. Hence

$$s\mathbf{s}_3 = r\mathbf{s}_2 + t\mathbf{s}_1, \quad s\mathbf{s}'_3 = r\mathbf{s}'_2 + t'\mathbf{s}'_1, \quad (429)$$

t, t' being rational numbers. Substituting these expressions for $s\mathbf{s}_3, s\mathbf{s}'_3$ into (427) and simplifying, we obtain $(t - t')[\mathbf{s}'_1, \mathbf{s}_1] = 0$. Considering that $[\mathbf{s}'_1, \mathbf{s}_1] \neq 0$ yields $t = t'$. It easily follows from (429) that $t\mathbf{s}_1$ is an integer vector and $t\mathbf{s}_1 = O(k^{3\delta})$. Hence, t is a rational number and the denominator of t is less than $|\mathbf{s}_1|$, i.e. it is $O(k^\delta)$. Multiplying both sides of (429) by the denominator of t , we rewrite (429) with all integers r, s, t such that $r, s, t = O(k^{3\delta})$. We already showed that $s, r \neq 0$. Note that $t \neq 0$ too, since otherwise $\vec{p}_{\mathbf{q}_3}$ and $\vec{p}_{\mathbf{q}_2}$ are colinear.

5. In this part by the way of contradiction we show that $\vec{p}_{\mathbf{q}_4}$ does not exist. Indeed, suppose it does. Excluding $(\vec{k} + \vec{p}_{\mathbf{m}_0}, \vec{p}_{\mathbf{q}_1})$ and $(\vec{k} + \vec{p}_{\mathbf{m}_0}, \vec{p}_{\mathbf{q}_2})$ from relations (417) for $\vec{p}_{\mathbf{q}_1}$, $\vec{p}_{\mathbf{q}_2}$ and $\vec{p}_{\mathbf{q}_3} = \frac{r}{s}\vec{p}_{\mathbf{q}_2} + \frac{t}{s}\vec{p}_{\mathbf{q}_1}$, we obtain:

$$p_{\mathbf{q}_1}^2 \frac{t}{s} \left(\frac{t}{s} - 1 \right) + 2p_{\mathbf{q}_1} p_{\mathbf{q}_2} \frac{t}{s} \frac{r}{s} + p_{\mathbf{q}_2}^2 \frac{r}{s} \left(\frac{r}{s} - 1 \right) = O(k^{-40\mu\delta+3\delta}). \quad (430)$$

Considering as in Part 4, we obtain $\vec{p}_{\mathbf{q}_4} = \frac{\hat{t}}{\hat{s}}\vec{p}_{\mathbf{q}_2} + \frac{\hat{t}}{\hat{s}}\vec{p}_{\mathbf{q}_1}$ and

$$p_{\mathbf{q}_1}^2 \frac{\hat{t}}{\hat{s}} \left(\frac{\hat{t}}{\hat{s}} - 1 \right) + 2p_{\mathbf{q}_1} p_{\mathbf{q}_2} \frac{\hat{t}}{\hat{s}} \frac{\hat{r}}{\hat{s}} + p_{\mathbf{q}_2}^2 \frac{\hat{r}}{\hat{s}} \left(\frac{\hat{r}}{\hat{s}} - 1 \right) = O(k^{-40\mu\delta+3\delta}). \quad (431)$$

- (a) Assume first $(t-s)^2 + (\hat{t}-\hat{s})^2 \neq 0$. We multiply both parts of (430) by s^2 and both parts of (431) by $\hat{s}^2$. Excluding the terms containing $p_{\mathbf{q}_1}^2$ from the last two relations and using the estimates $s, \hat{s}, r, \hat{r}, t, \hat{t} = O(k^{3\delta})$, we obtain:

$$p_{\mathbf{q}_1} p_{\mathbf{q}_2} S = R p_{\mathbf{q}_2}^2 + O(k^{-40\mu\delta+15\delta}),$$

where

$$S = 2\hat{t}r\hat{t}(t-s) - 2tr\hat{t}(\hat{t}-\hat{s}), \quad S = O(k^{12\delta}).$$

$$R = \hat{t}r(\hat{t}-\hat{s})(r-s) - t\hat{r}(t-s)(\hat{r}-\hat{s}), \quad R = O(k^{12\delta}).$$

It follows $p_{\mathbf{q}_1} S - p_{\mathbf{q}_2} R = O(k^{-39\mu\delta+15\delta})$.

- i. Suppose $R^2 + S^2 \neq 0$. Considering that the angle (modulo π) between $\vec{p}_{\mathbf{q}_1}$ and $\vec{p}_{\mathbf{q}_2}$ is less than $k^{-1+\delta}$ and (28) for $p_{\mathbf{q}_2}$, we obtain $|\vec{p}_{\mathbf{q}_1} S \pm \vec{p}_{\mathbf{q}_2} R| = O(k^{-39\mu\delta+15\delta})$, where $\pm$ means $+$ or $-$. The last relation yields $p_{S\mathbf{q}_1 \pm R\mathbf{q}_2} = O(k^{-39\mu\delta+15\delta})$. However, it follows from estimates for S and R that we have $\|\vec{p}_{S\mathbf{q}_1 \pm R\mathbf{q}_2}\| < ck^{13\delta}$. Hence, by (28) $p_{S\mathbf{q}_1 \pm R\mathbf{q}_2} > ck^{-13\delta\mu}$. We have arrived to the contradiction.
- ii. Now we check the case $R^2 + S^2 = 0$. It was shown in Part 4 that $s, \hat{s}, t, \hat{t}, r, \hat{r} \neq 0$. The equation $S = 0$ yields

$$\hat{r}(t-s) = r(\hat{t}-\hat{s}), \quad (432)$$

both parts are nonzero by the assumption $(t-s)^2 + (\hat{t}-\hat{s})^2 \neq 0$. Next, $R = 0$ yields

$$\hat{t}(r-s) = t(\hat{r}-\hat{s}). \quad (433)$$

Let us consider (432), (433) as a linear system with respect to $s, \hat{s}$. If the determinant of this system $\hat{r}t - r\hat{t}$ is zero, then it follows $r/\hat{r} = t/\hat{t} = s/\hat{s}$. This means that $\vec{p}_{\mathbf{q}_3}$ and $\vec{p}_{\mathbf{q}_4}$ are equal. This contradicts to our initial assumption. Suppose that the determinant is not zero. Solving (432), (433) with respect to s and $\hat{s}$, we get: $s = t + r, \hat{s} = \hat{t} + \hat{r}$. Substituting

$s = t + r$ into (430), we easily obtain: $\frac{rt}{s^2} (p_{\mathbf{q}_1} - p_{\mathbf{q}_2})^2 = O(k^{-40\mu\delta+3\delta})$. Considering that $s^2 = O(k^{6\delta})$ and the angle (modulo π) between $\vec{p}_{\mathbf{q}_1}, \vec{p}_{\mathbf{q}_2}$ is $O(k^{-1+\delta})$, we obtain $|\vec{p}_{\mathbf{q}_1} \pm \vec{p}_{\mathbf{q}_2}| = O(k^{-20\mu\delta+5\delta})$, where $\pm$ means $+$ or $-$. Obviously, $p_{\mathbf{q}_1-\mathbf{q}_2} = |\vec{p}_{\mathbf{q}_1} - \vec{p}_{\mathbf{q}_2}|$. Hence, $p_{\mathbf{q}_1-\mathbf{q}_2} = O(k^{-20\mu\delta+5\delta})$. This cannot be the case, because $||\vec{p}_{\mathbf{q}_1-\mathbf{q}_2}|| < 3k^\delta$ and thus, $|p_{\mathbf{q}_1-\mathbf{q}_2}| \geq ck^{-\mu\delta}$.

- (b) It remains to consider the case $(t-s)^2 + (\hat{t}-\hat{s})^2 = 0$. If $(r-s)^2 + (\hat{r}-\hat{s})^2 \neq 0$, we exclude $p_{\mathbf{q}_2}^2$ from (430), (431) by and make considerations similar to the case $(t-s)^2 + (\hat{t}-\hat{s})^2 \neq 0$. If $(t-s)^2 + (\hat{t}-\hat{s})^2 + (r-s)^2 + (\hat{r}-\hat{s})^2 = 0$, then (430), (431) give: $p_{\mathbf{q}_1}p_{\mathbf{q}_2} = O(k^{-40\mu\delta+3\delta})$. It contradict to the inequalities (28) for $p_{\mathbf{q}_1}, p_{\mathbf{q}_2}$. Thus, $\vec{p}_{\mathbf{q}_4}$ does not exist.

□

10.3 Appendix 3

Lemma 10.1. *The equation*

$$\lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}) = k^{2l} + \varepsilon_0, \quad 0 < p_{\mathbf{m}} \leq 4k^\delta, \quad |\varepsilon_0| \leq p_{\mathbf{m}}k^\delta, \quad (434)$$

has no more than two solutions $\varphi^\pm(\varepsilon_0)$ in $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$. They satisfy the estimates:

$$|\varphi^\pm(\varepsilon_0) - \varphi_{\mathbf{m}}^\pm| < k^{-2l+1+2\delta}.$$

Proof. Let $\varphi \in \tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$. The equation (434) is equivalent to

$$\lambda^{(1)}(\vec{y}(\varphi)) = \lambda^{(1)}(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}) + \varepsilon_0, \quad \vec{y}(\varphi) = \vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}.$$

We use perturbation formula (39):

$$|\vec{y}(\varphi)|_{\mathbb{R}}^{2l} + f_1(\vec{y}(\varphi)) = |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} + f_1(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}) + \varepsilon_0,$$

where f_1 is the series in the right-hand side of (39). This equation can be rewritten as

$$\left(2(\vec{\mathcal{Z}}^{(1)}(\varphi), \vec{p}_{\mathbf{m}})_{\mathbb{R}} + p_{\mathbf{m}}^2\right) \left(|\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} + \dots + |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l-2}\right) + f_1(\vec{y}(\varphi)) - f_1(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}) = \varepsilon_0. \quad (435)$$

Using the notation $\vec{p}_{\mathbf{m}} = p_{\mathbf{m}}(\cos \varphi_{\mathbf{m}}, \sin \varphi_{\mathbf{m}})$, dividing both sides of the equation (435) by $2p_{\mathbf{m}}k \left(|\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} + \dots + |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l-2}\right)$, and considering that $\vec{y}(\varphi) = \vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}} = (k + h^{(1)})\vec{\nu} + \vec{p}_{\mathbf{m}}$, we obtain:

$$\cos(\varphi - \varphi_{\mathbf{m}}) + \frac{p_{\mathbf{m}}}{2k} - \varepsilon_0 g_1(\varphi) + g_2(\varphi) + g_3(\varphi) = 0, \quad (436)$$

where $g_1(\varphi) = (2p_{\mathbf{m}}k)^{-1} \left(|\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} + \dots + |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l-2} \right)^{-1}$ and

$$g_2(\varphi) = \frac{(\vec{h}^{(1)}(\varphi), \vec{p}_{\mathbf{m}})}{p_{\mathbf{m}}k}, \quad g_3(\varphi) = \left(f_1(\vec{y}(\varphi)) - f_1(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}) \right) g_1(\varphi), \quad \vec{h}^{(1)}(\varphi) = h^{(1)}(\varphi)\vec{v}.$$

Obviously $g_1 = O(p_{\mathbf{m}}^{-1}k^{-2l+1})$. Using Lemma 3.11 and considering that $0 < p_{\mathbf{m}} \leq 4k^\delta$, we easily obtain:

$$|g_2(\varphi)| = \left| \frac{(\vec{h}^{(1)}(\varphi), \vec{p}_{\mathbf{m}})}{p_{\mathbf{m}}k} \right| \leq \frac{2h^{(1)}}{k} = O(k^{-4l+(80\mu+6)\delta}).$$

Let us show $g_3(\varphi) = O(k^{-2l+1+\delta})$. If $p_{\mathbf{m}} \geq k^{-2l+\delta(80\mu+6)}$, then the estimate easily follows from (45) and the estimate for g_1 . Let $p_{\mathbf{m}} < k^{-2l+\delta(80\mu+6)}$. It can be easily shown that the series $f_1(\vec{y})$, $\nabla f_1(\vec{y})$ converge for all $\vec{y}$: $|\vec{y} - \vec{\mathcal{Z}}^{(1)}(\varphi)|_{\mathbb{C}^2} < k^{-1-\delta(40\mu+1)}$ and holomorphic with respect to y_1, y_2 . Using (45), we get $\nabla f_1(y) = k^{-2l+1+\delta(120\mu+7)}$. Hence,

$$|f_1(\vec{y}(\varphi)) - f_1(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}})| \leq \sup |\nabla f_1| p_{\mathbf{m}} = o(p_{\mathbf{m}}),$$

and therefore, $g_3(\varphi) = O(k^{-2l+1+\delta})$. Since $|\varepsilon_0| < p_{\mathbf{m}}k^\delta$, we obtain $\varepsilon_0 g_1(\varphi) = O(k^{-2l+1+\delta})$. Thus,

$$g_2(\varphi) + g_3(\varphi) - \varepsilon_0 g_1(\varphi) = O(k^{-2l+1+\delta}) \quad \text{when } \varphi \in \tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2}). \quad (437)$$

By definition $\varphi_{\mathbf{m}}^\pm$ satisfy the equation $\cos(\varphi - \varphi_{\mathbf{m}}) + \frac{p_{\mathbf{m}}}{2k} = 0$.

Suppose both $\varphi_{\mathbf{m}}^\pm$ are in $\tilde{\mathcal{W}}^{(1)}(k, \frac{3}{16})$. We draw two circles $C_\pm$ centered at $\varphi_{\mathbf{m}}^\pm$ with the radius $k^{-2l+1+2\delta}$. They are both inside $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$, the perturbation series converging and the estimate (437) holds. For any φ on $C_\pm$, $|\varphi - \varphi_{\mathbf{m}}^\pm| = k^{-2l+1+2\delta}$ and, therefore, $|\cos(\varphi - \varphi_{\mathbf{m}}) + \frac{p_{\mathbf{m}}}{2k}| > \frac{1}{2}k^{-2l+1+2\delta} > |g_2(\varphi) + g_3(\varphi) - \varepsilon_0 g_1(\varphi)|$ for any $\varphi \in C_\pm$. By Rouché's Theorem, there is only one solution of the equation (436) inside each $C_\pm$. Obviously, (436) does not have solutions in $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$ outside $C_\pm$.

If both $\varphi_{\mathbf{m}}^\pm$ are not in $\tilde{\mathcal{W}}^{(1)}(k, \frac{3}{16})$, then their distance to $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8})$ is at least $\frac{1}{16}k^{-40\mu\delta}$, hence $|\cos(\varphi - \varphi_{\mathbf{m}}) + \frac{p_{\mathbf{m}}}{2k}| > \frac{1}{4}k^{-2l+1+2\delta}$ in $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8})$. Therefore, equation (436) has no solution in $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$. The case, when only one $\varphi_{\mathbf{m}}^\pm$ is not in $\tilde{\mathcal{W}}^{(1)}(k, \frac{3}{16})$ is the obvious combination of the two previous situations. Thus, there are at most two solutions in $\tilde{\mathcal{W}}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$ and $|\varphi^\pm(\varepsilon_0) - \varphi_{\mathbf{m}}^\pm| < k^{-2l+1+2\delta}$. $\square$

Lemma 10.2. For any $\varphi \in \tilde{\mathcal{W}}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, 1)$ satisfying the estimate $|\varphi - \varphi_{\mathbf{m}}^\pm| < k^{-\delta}$,

$$\frac{\partial}{\partial \varphi} \lambda^{(1)}(\vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}) = \pm 2lp_{\mathbf{m}}k^{2l-1}(1 + o(1)), \quad (438)$$

Proof. First, assume φ is real. Let $\vec{y}(\varphi) = \vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}$. Using the perturbation formula (39) and Lemma 3.11, we obtain:

$$\begin{aligned} \frac{\partial}{\partial \varphi} \lambda^{(1)}(\vec{y}(\varphi)) &= \frac{\partial}{\partial \varphi} [\lambda^{(1)}(\vec{y}(\varphi)) - k^{2l}] = \frac{\partial}{\partial \varphi} [\lambda^{(1)}(\vec{y}(\varphi)) - \lambda^{(1)}(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}})] = \\ &= \left(\nabla_{\vec{y}} \lambda^{(1)}(\vec{y}(\varphi)) - \nabla_{\vec{y}} \lambda^{(1)}(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}), \frac{\partial}{\partial \varphi} \vec{y}(\varphi) \right)_{\mathbb{R}} = \\ &= \left(\nabla |\vec{y}(\varphi)|_{\mathbb{R}}^{2l} - \nabla |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l}, (k + h^{(1)})\vec{t} + (h^{(1)})'\vec{v} \right)_{\mathbb{R}} + \\ &\quad (\nabla f_1(\vec{y}(\varphi)) - \nabla f_1(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}), (k + h^{(1)})\vec{t} + (h^{(1)})'\vec{v})_{\mathbb{R}}, \end{aligned} \quad (439)$$

where $\vec{v} = (\cos \varphi, \sin \varphi)$ and $\vec{t} = \vec{v}' = (-\sin \varphi, \cos \varphi)$, f_1 is the series in the right-hand side of (39). Note that

$$\begin{aligned} &\nabla |\vec{y}(\varphi)|_{\mathbb{R}}^{2l} - \nabla |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l} = \\ &2l |\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} \vec{y}(\varphi) - 2l |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l-2} (\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}) = \\ &2l |\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} \vec{p}_{\mathbf{m}} + 2l \left(|\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} - |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l-2} \right) (\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}). \end{aligned} \quad (440)$$

Substituting (440) into (439), we get $\frac{\partial}{\partial \varphi} \lambda^{(1)}(\vec{y}(\varphi)) = T_1 + T_2 + T_3$,

$$\begin{aligned} T_1 &= 2l |\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} (\vec{p}_{\mathbf{m}}, (k + h^{(1)})\vec{t} + (h^{(1)})'\vec{v})_{\mathbb{R}}, \\ T_2 &= \left(|\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} - |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l-2} \right) (\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}, (k + h^{(1)})\vec{t} + (h^{(1)})'\vec{v})_{\mathbb{R}}, \\ T_3 &= (\nabla f_1(\vec{y}(\varphi)) - \nabla f_1(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}), (k + h^{(1)})\vec{t} + (h^{(1)})'\vec{v})_{\mathbb{R}}. \end{aligned}$$

We see that φ is close to $\varphi_{\mathbf{m}} \pm \pi/2$, since $|\varphi - \varphi_{\mathbf{m}}^{\pm}| < k^{-\delta}$ by the hypothesis of the lemma and $\varphi_{\mathbf{m}}^{\pm} = \varphi_{\mathbf{m}} \pm \pi/2 + O(k^{-1+\delta})$ when $p_{\mathbf{m}} < 4k^{\delta}$. Now we readily obtain: $(\vec{p}_{\mathbf{m}}, \vec{v})_{\mathbb{R}} = o(p_{\mathbf{m}})$, $(\vec{p}_{\mathbf{m}}, \vec{t})_{\mathbb{R}} = \pm p_{\mathbf{m}}(1 + o(1))$. Using also estimates (65), (66) for $h^{(1)}$, we get $T_1 = \pm 2l p_{\mathbf{m}} k^{2l-1}(1 + o(1))$. Note that $\vec{y}(\varphi) - \vec{p}_{\mathbf{m}} = \vec{\mathcal{Z}}^{(1)}(\varphi)$ and, hence, it is orthogonal to $\tau(\varphi)$. Using this fact, we simplify the expression for T_2 :

$$T_2 = \left(|\vec{y}(\varphi)|_{\mathbb{R}}^{2l-2} - |\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}|_{\mathbb{R}}^{2l-2} \right) (\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}, (h^{(1)})'\vec{v})_{\mathbb{R}}.$$

Using (66) for $(h^{(1)})'$, we obtain $T_2 = o(p_{\mathbf{m}} k^{2l-1})$. Let us estimate T_3 . It can be easily shown that the series $f_1(\vec{y})$, $\nabla f_1(\vec{y})$, $D^2 f_1(\vec{y})$ converge for all $\vec{y}$: $|\vec{y} - \vec{\mathcal{Z}}_1(\varphi)|_{\mathbb{C}^2} < k^{-1-\delta(40\mu+1)}$ or $|\vec{y} + \vec{p}_{\mathbf{m}} - \vec{\mathcal{Z}}_1(\varphi)|_{\mathbb{C}^2} < k^{-1-\delta(40\mu+1)}$, the series being holomorphic with respect to y_1, y_2 in these neighborhoods. Using (45), we get $\nabla f_1(\vec{y}) = k^{-2l+1+\delta(120\mu+7)}$, $D^2 f_1(\vec{y}) = k^{-2l+2+\delta(160\mu+8)}$. Let $p_{\mathbf{m}} \geq \frac{1}{2} k^{-1-\delta(40\mu+1)}$. Then, using the estimate for $\nabla f_1(\vec{y})$, we easily obtain $T_3 = k^{-2l+2+\delta(120\mu+7)} = o(k^{2l-1} p_{\mathbf{m}})$. Let $p_{\mathbf{m}} < \frac{1}{2} k^{-1-\delta(40\mu+1)}$. Then, using the estimate for the second derivative in the direction of $\vec{p}_{\mathbf{m}}$, we get $|\nabla f_1(\vec{y}(\varphi)) -$

$\nabla f_1(\vec{y}(\varphi) - \vec{p}_{\mathbf{m}}) \Big| = O(p_{\mathbf{m}}k^{-2l+2+(160\mu+8)\delta})$. Therefore, $T_3 = O(p_{\mathbf{m}}k^{-2l+3+(160\mu+8)\delta})$. Thus, $T_3 = o(k^{2l-1}p_{\mathbf{m}})$ for all $\vec{p}_{\mathbf{m}}$. Adding the estimates for T_1, T_2, T_3 , we get (438).

Since all formulas can be analytically extended to the area of non-real φ , the estimates being preserved, (438) holds for any $\varphi \in \tilde{W}^{(1)}(k, \frac{1}{4}) \cap \mathcal{O}_{\mathbf{m}}(k, 1)$. $\square$

Lemma 10.3. *Let $\tilde{\mathcal{O}}_{\mathbf{m},\varepsilon}^{\pm}$ be the open discs of the radius ε centered at $\varphi^{\pm}(0)$. For any $\varphi \in \tilde{W}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$, $\varphi \notin \tilde{\mathcal{O}}_{\mathbf{m},\varepsilon}^{\pm}$, and $0 \leq \varepsilon < k^{-2l+1+\delta}$,*

$$|\lambda^{(1)}(\vec{y}(\varphi)) - k^{2l}| \geq k^{2l-1}p_{\mathbf{m}}\varepsilon. \quad (441)$$

Proof. Suppose (441) does not hold for some $\varphi \in \tilde{W}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$, $\varphi \notin \tilde{\mathcal{O}}_{\mathbf{m},\varepsilon}^{\pm}$. This means that φ satisfies equation (434) with some ε_0 : $|\varepsilon_0| < k^{2l-1}p_{\mathbf{m}}\varepsilon (< p_{\mathbf{m}}k^{\delta})$. By Lemma 10.1, φ could be either $\varphi^+(\varepsilon_0)$ or $\varphi^-(\varepsilon_0)$. Without loss of generality, assume $\varphi = \varphi^+(\varepsilon_0)$. By Lemma 10.1, $|\varphi^+(\varepsilon_0) - \varphi_{\mathbf{m}}^{\pm}| < k^{-2l+1+2\delta}$ for $\varphi_{\mathbf{m}}^+$ or $\varphi_{\mathbf{m}}^-$. Obviously, $k^{-2l+1+2\delta}$ neighborhood of $\varphi^+(\varepsilon_0)$ satisfies conditions of Lemma 10.2. Using (438) and Rouché's theorem in the $k^{-2l+1+2\delta}$ -neighborhood of $\varphi^+(\varepsilon_0)$, we obtain that there is a point $\varphi^+(0)$ in this neighborhood and $|\varphi^+(\varepsilon_0) - \varphi^+(0)| < \varepsilon$, i.e., $\varphi \in \tilde{\mathcal{O}}_{\mathbf{m},\varepsilon}$. This contradicts the hypothesis of the lemma. $\square$

Lemma 10.4. *If $0 < p_{\mathbf{m}} \leq 4k^{\delta}$ and $\varphi \in \tilde{W}^{(1)}(k, \frac{1}{8}) \cap \mathcal{O}_{\mathbf{m}}(k, \frac{1}{2})$, then*

$$\left\| (\lambda^{(1)}(\vec{y}(\varphi)) - k^{2l}) (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - k^{2l})P_{\mathbf{m}})^{-1} \right\| \leq 8, \quad \vec{y}(\varphi) := \vec{\mathcal{Z}}^{(1)}(\varphi) + \vec{p}_{\mathbf{m}}. \quad (442)$$

Proof. Let $\tilde{C}_1$ be the circle in $\mathbb{C}$ of the radius $k^{2l-1-80\mu\delta}$ centered at $z = |\vec{y}(\varphi)|_{\mathbb{R}}^{2l}$. Using (92) and (93), we easily get:

$$|\vec{y}(\varphi) + \vec{p}_{\mathbf{q}}|_{\mathbb{R}}^{2l} - z \gtrsim \frac{1}{2}k^{2l-1-40\mu\delta}, \quad \text{when } \mathbf{q} \in \Omega(\delta), \quad z \in \tilde{C}_1.$$

Therefore,

$$\left\| (P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1} \right\| \lesssim 2k^{-2l+1+40\mu\delta}, \quad (443)$$

$$\left\| (P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1} \right\|_1 \lesssim 2k^{-2l+1+40\delta(\mu+1)}. \quad (444)$$

Next, by (89),

$$\left| \det \frac{P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}}}{P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}}} - 1 \right| < 2\|V\| \left\| (P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1} \right\|_1 \quad (445)$$

for every z on the contour $\tilde{C}_1$. Using the estimate (444), we obtain that the right-hand part of (445) is less than 1. Applying Rouché's theorem, we conclude that the determinant has the same number of zeros and poles inside $\tilde{C}_1$. Considering that the resolvent $(P_{\mathbf{m}}(H_0(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1}$ has a single pole, $z = |\vec{y}(\varphi)|_{\mathbb{R}}^{2l}$, we obtain that

$(P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1}$ has a single pole inside $\tilde{C}_1$ too. Obviously, the pole is in the point $z = \lambda^{(1)}(\vec{y}(\varphi))$. Therefore

$$(\lambda^{(1)}(\vec{y}(\varphi)) - z) (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1}$$

is a holomorphic function of z inside $\tilde{C}_1$.

Let $z \in \tilde{C}_1$. Using (45), we easily obtain: $|\lambda^{(1)}(\vec{y}(\varphi)) - z| \leq 2k^{2l-1-40\mu\delta}$. From (443) and Hilbert identity it follows that

$$\left\| (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1} \right\| \leq 4k^{-2l+1+40\mu\delta}, \text{ when } z \in \tilde{C}_1. \quad (446)$$

Multiplying the last two estimates, and using maximum principle we get

$$\left\| (\lambda^{(1)}(\vec{y}(\varphi)) - z) (P_{\mathbf{m}}(H(\vec{\mathcal{Z}}^{(1)}(\varphi)) - z)P_{\mathbf{m}})^{-1} \right\| \leq 8, \quad z \in \overline{\text{Int } \tilde{C}_1}. \quad (447)$$

Note that $z := k^{2l} \in \overline{\text{Int } \tilde{C}_1}$. Indeed, by (92), $||\vec{y}(\varphi)|_{\mathbb{R}}^{2l} - k^{2l}| < k^{2l-2-40\mu\delta}$. Substituting $z = k^{2l}$ in the last estimate, we get (442). $\square$

10.4 Appendix 4

Lemma 10.5. *Let R be the smallest positive integer for which (231) holds. We have $R > \frac{1}{64}k^{(\frac{\gamma}{2}+2\delta_0)r_1-\delta}$.*

Proof. Notice that

$$\mathcal{A}_R = \sum_{i_1, \dots, i_R=0}^{\hat{I}} \mathcal{A}_{i_1, \dots, i_R},$$

where

$$\mathcal{A}_{i_1, \dots, i_R} := P^{(int)} W(\hat{H}_0 - k^{2l})^{-1} \left[\prod_{k=1}^R (P_{i_k} W(\hat{H}_0 - k^{2l})^{-1}) \right] P^{\partial}.$$

Here we used that R is the smallest positive integer for which $\mathcal{A}_R \neq 0$.

In some sense everything is defined by the case where all i_k are equal to zero. But to include impurities of non-resonant and white clusters we need additional construction. Consider a particular $\mathcal{A}_{i_1, \dots, i_R}$. For the sequence $i_1, \dots, i_R$ we take a subsequence of *all* non-zero indices $i_{k_1}, \dots, i_{k_s}$, $1 \leq k_1 < \dots < k_s \leq R$ (this sequence can be empty). Now we construct a subsequence $j_1, \dots, j_p$ ($p \leq s$) of non-repeating indices as follows. We choose $j_1 = i_{k_1}$. If i_{k_1} is not equal to any other i_{k_t} , $t = 2, \dots, s$, then $j_2 = i_{k_2}$. If there is one or more i_{k_t} equal to i_{k_1} then we denote the segment between the first and the last i_{k_1} as I_1 . The next term after I_1 we choose to be an j_2 etc. Thus (with a slight abuse of the notation) we have

$$i_{k_1}, \dots, i_{k_s} = I_1, I_2, \dots, I_p, \quad p \leq s.$$

Now, the initial sequence $i_1, \dots, i_R$ can be represented as $I_1^0, \tilde{I}_1, I_2^0, \tilde{I}_2, \dots, \tilde{I}_p, I_{p+1}^0$. Here each I_j^0 is a sequence of only zeros (it can be empty) and $\tilde{I}_j$ is I_j with possibly some zeros inside. Put

$$P_{j_1} B P_{j_1} := P_{j_1} W(\hat{H}_0 - k^{2l})^{-1} P_{i'} W(\hat{H}_0 - k^{2l})^{-1} \dots (\hat{H}_0 - k^{2l})^{-1} P_{j_1}.$$

Here for all internal projectors $P_{i'}$ we have either $i' \in I_1$ or $i' = 0$. We notice that $P_{j_1} B P_{j_1}$ has a block form and $\|P_{j_1} B P_{j_1}\| \leq k^{-\beta}$. We can represent now $\mathcal{A}_{i_1, \dots, i_R}$ in the following form

$$\begin{aligned} \mathcal{A}_{i_1, \dots, i_R} &= P^{(int)} W(\hat{H}_0 - k^{2l})^{-1} \left[\prod_{k=1}^p \left(P_0 W(\hat{H}_0 - k^{2l})^{-1} \right)^{s_k} P_{j_k} B P_{j_k} W(\hat{H}_0 - k^{2l})^{-1} \right] \\ &\times \left(P_0 W(\hat{H}_0 - k^{2l})^{-1} \right)^{s_{p+1}} P^\partial. \end{aligned} \quad (448)$$

Here s_k is the number of elements in I_k^0 , $s_k \geq 0$; j_k is a non-zero index corresponding to I_k . Obviously, $\left(W(\hat{H}_0 - k^{2l})^{-1} \right)_{\mathbf{m}\mathbf{m}'} = 0$ if $\|\cdot\|$ -distance between the cluster containing $\vec{p}_{\mathbf{m}}$ and the cluster containing $\vec{p}_{\mathbf{m}'}$ is greater than k^δ (here, as usual, we consider the points in the range of P_0 as 1×1 clusters). Next, if P_{j_k} is the projection on a non-resonant cluster, then

$$(P_{j_k} B P_{j_k})_{\mathbf{m}\mathbf{m}'} = 0, \quad \text{for } \|\vec{p}_{\mathbf{m}-\mathbf{m}'}\| > 8k^\delta,$$

since a non-resonant cluster has the size not greater than $8k^\delta$. Let p' be the number of non-resonant projections in the sequence $\{P_{j_k}\}_{k=1}^p$. Hence, $p - p'$ is the number of white clusters. The operator $\mathcal{A}_{i_1, \dots, i_R}$ can be non-zero only if

$$\frac{D}{2} \leq k^\delta \sum_{k=1}^{p+1} (s_k + 1) + 8k^\delta p' + \sum_{m=1}^{p-p'} d_m, \quad (449)$$

where d_m is the size of a white cluster $P_{j_{k_m}}$. Next, we prove that

$$\sum_{k=1}^{p+1} (s_k + 1) + p' \geq \frac{1}{32} k^{(\frac{\gamma}{2} + 2\delta_0)r_1 - \delta}. \quad (450)$$

Assume that (450) does not hold. Then, by (449)

$$\sum_{m=1}^{p-p'} d_m \geq \frac{1}{4} k^{(\frac{\gamma}{2} + 2\delta_0)r_1}, \quad (451)$$

since $D = k^{(\frac{\gamma}{2} + 2\delta_0)r_1}$. Obviously,

$$d_m \leq n_m k^{\frac{\gamma r_1}{6}},$$

where n_m is the number of $\mathcal{M}^{(2)}$ points in the white cluster $\Pi_{j_{k_m}}$, $m = 1, \dots, p - p'$. Let ℓ be the size of a minimal box containing all white clusters $\Pi_{j_{k_m}}$. It is easy to see that

$$\ell \leq k^\delta \sum_{k=1}^{p+1} (s_k + 1) + 8k^\delta p' + \sum_{m=1}^{p-p'} d_m \leq \frac{1}{4} k^{(\frac{\gamma}{2} + 2\delta_0)r_1} + \sum_{m=1}^{p-p'} d_m \leq 2 \sum_{m=1}^{p-p'} d_m.$$

Here we also used (451) and the inequality opposite to (450). By Lemma 4.11

$$\sum_{m=1}^{p-p'} n_m \leq C\ell^{2/3}k. \quad (452)$$

Combining the last three inequalities and solving for $\sum_{m=1}^{p-p'} d_m$, we obtain: $\sum_{m=1}^{p-p'} d_m < k^{\frac{\gamma r_1}{2} + 3}$. This contradicts to (451). Thus, we proved (450). Using the obvious inequality $\sum_{k=1}^{p+1} (s_k + 1) + p' \leq R + 1$ proves the lemma.

□

10.5 Appendix 5

Lemma 10.6. *Let R be the smallest positive integer for which (231) holds in the case of a black cluster. We have $R > \frac{1}{64} k^{\gamma r_1 + \delta_0 r_1 - \delta}$.*

Proof. We again use formula (448), where P_{j_k} are projections on non-resonant, white and grey components in a component of a black region. Assume first that all components Π_{j_k} can be placed in one $\|\cdot\|$ box of the size $4k^{\gamma r_1 + \delta_0 r_1}$. Obviously,

$$\frac{1}{2} k^{\gamma r_1 + \delta_0 r_1} \leq k^\delta \sum (s_k + 1) + 8k^\delta p' + \sum_m d_m^w + \sum_{\tilde{m}} d_{\tilde{m}}^g, \quad (453)$$

where p' is the number of non-resonant components in the black component, $\sum_m d_m^w$ and $\sum_{\tilde{m}} d_{\tilde{m}}^g$ are the total lengths of white and grey components in the black component. Let us prove first that $k^\delta \sum (s_k + 1) + 8k^\delta p' + \sum_m d_m^w > \frac{1}{4} k^{\gamma r_1 + \delta_0 r_1}$. Suppose that it is not so. Then, by (453), $\sum_{\tilde{m}} d_{\tilde{m}}^g > \frac{1}{4} k^{\gamma r_1 + \delta_0 r_1}$. The $4k^{\gamma r_1 + \delta_0 r_1}$ -box containing all Π_{j_k} , consists of no more than $4^4 k^{4\delta_0 r_1}$ boxes of the size $k^{\gamma r_1}$. Since all Π_{j_k} are in white $k^{\gamma r_1}$ -boxes, the total number of points of $\mathcal{M}^{(2)}$ in these white boxes does not exceed $ck^{\frac{1}{2}\gamma r_1 + \delta_0 r_1} \cdot k^{4\delta_0 r_1}$. Since each grey box contains no less than $k^{\frac{1}{6}\gamma r_1 - \delta_0 r_1}$ points, the total number of grey boxes is less than $ck^{\frac{1}{2}\gamma r_1 + \delta_0 r_1} \cdot k^{4\delta_0 r_1} \cdot k^{-\frac{1}{6}\gamma r_1 + \delta_0 r_1} = ck^{\frac{1}{3}\gamma r_1 + 6\delta_0 r_1}$. Therefore, the total size of the grey region is less than $ck^{\frac{1}{3}\gamma r_1 + 6\delta_0 r_1} \cdot k^{\frac{1}{2}\gamma r_1 + 2\delta_0 r_1} = ck^{\frac{5}{6}\gamma r_1 + 8\delta_0 r_1}$. Since $\delta_0 < \frac{1}{48}\gamma$, it is much less than $\frac{1}{4} k^{\gamma r_1 + \delta_0 r_1}$. We have arrived to the contradiction with the assumption $\sum_{\tilde{k}} d_{\tilde{k}}^g > \frac{1}{4} k^{\gamma r_1 + \delta_0 r_1}$. Therefore, $k^\delta \sum (s_k + 1) + 8k^\delta p' + \sum_k d_k^w > \frac{1}{4} k^{\gamma r_1 + \delta_0 r_1}$. Considering again that the total number of $\mathcal{M}^{(2)}$ points in the white boxes of the $4k^{\gamma r_1 + \delta_0 r_1}$ -box does not exceed $ck^{\frac{\gamma r_1}{2} + 5\delta_0 r_1}$, we obtain $\sum_k d_k^w < ck^{\frac{\gamma r_1}{2} + 5\delta_0 r_1} \cdot 4k^{\frac{\gamma r_1}{6}} < \frac{1}{20} k^{\gamma r_1 + \delta_0 r_1}$. It follows $k^\delta \sum (s_k + 1) + 8k^\delta p' > \frac{1}{5} k^{\gamma r_1 + \delta_0 r_1}$. Hence, $R + 1 \geq \sum (s_k + 1) + p' > \frac{1}{40} k^{\gamma r_1 + \delta_0 r_1 - \delta}$.

Assume that we cannot put all the components Π_{j_k} in one $||| \cdot |||$ -box of the size $4k^{\gamma r_1 + \delta_0 r_1}$. Let us consider the box of this size around Π_{j_1} . Let K be a number such that all Π_{j_k} , $k = 1, \dots, K$ are in the box and $\Pi_{j_{K+1}}$ is not. Then, instead of (448) we consider just its piece

$$P_{j_1} W(\hat{H}_0 - k^{2l})^{-1} \left[\prod_{k=2}^K \left(P_0 W(\hat{H}_0 - k^{2l})^{-1} \right)^{s_k} P_{j_k} B P_{j_k} W(\hat{H}_0 - k^{2l})^{-1} \right] \times \left(P_0 W(\hat{H}_0 - k^{2l})^{-1} \right)^{s_{K+1}} P_{j_{K+1}}. \quad (454)$$

Further considerations are the same as in the previous case since by construction the distance between Π_{j_1} and $\Pi_{j_{K+1}}$ is at least $\frac{1}{2}k^{\gamma r_1 + \delta_0 r_1}$.

□

10.6 Appendix 6. On Application of Bezout Theorem

Let $D(\vec{\varkappa}, \lambda)$ be the determinant of the truncated operator $H(\vec{\varkappa}) - \lambda$ of the size k^{r_*} , $r_* \geq 1$. Obviously, D is the polynomial of the degree k^{4lr_*} with respect to $\varkappa_1, \varkappa_2$ and a line is not a solution of the equation $D(\vec{\varkappa}, \lambda) = 0$. Let λ be fixed, $\lambda = k^{2l}$.

Definition 10.7. We call a piece of $D(\vec{\varkappa}, \lambda) = 0$ elementary, if

- 1) it can be parameterized by $\varkappa_1$: $\varkappa_2 = \varkappa_2(\varkappa_1)$ with $|\varkappa_2'(\varkappa_1)| \leq 1$ or by $\varkappa_2$: $\varkappa_1 = \varkappa_1(\varkappa_2)$ with $|\varkappa_1'(\varkappa_2)| \leq 1$;
- 2) function $\varkappa_1 = \varkappa_1(\varkappa_2)$ (or $\varkappa_2 = \varkappa_2(\varkappa_1)$) is monotone and continuously differentiable;
- 3) it does not have inflection points inside;
- 4) it has a length not greater than 1.

We will show that the curve $D(\vec{\varkappa}, \lambda) = 0$ can be split into elementary pieces and estimate the number of such pieces. In the proof we will apply several times the following statement (which is a simplified version of Bezout Theorem).

Theorem 10.8. Let P and Q be two plane real-valued polynomials of degree p and q respectively. If P and Q do not contain common factors then the total number of points satisfying $P(\vec{\varkappa}) = 0 = Q(\vec{\varkappa})$ (i.e. number of points of intersection) does not exceed pq .

We have

Lemma 10.9. The set $D(\vec{\varkappa}, \lambda) = 0$ can be split into k^{17lr_*} or less elementary pieces.

Proof. First, $D(\vec{\varkappa}, \lambda)$ can be represented as a product of simple (i.e. irreducible) factors (counting multiplicity). The total number of factors is less than k^{4lr_*} which is also the bound for their total degree. We consider one of such simple factors P and denote by

p its degree (note that we do ignore the multiplicity of the factor). Let us consider the points

$$P(\vec{\varkappa}) = 0, \quad \frac{\partial}{\partial \varkappa_2} P(\vec{\varkappa}) = 0. \quad (455)$$

Since P is irreducible and $\frac{\partial}{\partial \varkappa_2} P(\vec{\varkappa})$ has degree less than p (we also notice that $\frac{\partial}{\partial \varkappa_2} P(\vec{\varkappa})$ is not identically zero since $D(\vec{\varkappa}, \lambda) = 0$ does not contain lines) they do not have common factors. Thus, the number of such points $\vec{\varkappa}$ does not exceed $p(p-1)$. Next, by the same reasons the number of points

$$P(\vec{\varkappa}) = 0, \quad \frac{\partial}{\partial \varkappa_1} P(\vec{\varkappa}) = 0 \quad (456)$$

does not exceed $p(p-1)$ and the number of points

$$P(\vec{\varkappa}) = 0, \quad \frac{\partial P}{\partial \varkappa_2}(\vec{\varkappa}) = \pm \frac{\partial P}{\partial \varkappa_1}(\vec{\varkappa}) \quad (457)$$

does not exceed $2p(p-1)$. We split each previous piece by such points. Thus, we have at most $4p(p-1)+1$ pieces, each end satisfying (455) or (457). The sign of $(\frac{\partial}{\partial \varkappa_2} P)^2 - (\frac{\partial}{\partial \varkappa_1} P)^2$ is constant on each piece, i.e. the piece admits parametrization as in the property 2 of Definition 10.7. Making parametrization by $\varkappa_1$ or $\varkappa_2$, depending on the sign, we obtain that the length of a piece does not exceed $\sqrt{2} \cdot 4k^{r*}$ (obviously, $|\varkappa_j| < 2k^{r*}$). Therefore the total length of the curve $P = 0$ does not exceed $18p^2k^{r*}$. Next, for each piece where $\frac{\partial}{\partial \varkappa_2} P(\vec{\varkappa}) \neq 0$ inflection points of $P(\vec{\varkappa}) = 0$ are described by the system

$$P = 0, \quad P_{\varkappa_2 \varkappa_2} (P_{\varkappa_1})^2 - 2P_{\varkappa_1 \varkappa_2} P_{\varkappa_1} P_{\varkappa_2} + P_{\varkappa_1 \varkappa_1} (P_{\varkappa_2})^2 = 0. \quad (458)$$

Again, since P is irreducible and no line is a solution, we have no common factors here and can apply Bezout Theorem. The number of points satisfying (458) does not exceed $p(2(p-1) + (p-2)) = 3p^2 - 4p$. Therefore, we have at most $12p^4$ pieces with the ends satisfying (455) or (457) or (458). At last, we split each of these concave pieces into pieces with the length not greater than 1. Considering that the total length of $P(\vec{\varkappa}) = 0$ is less than $18p^2k^{r*}$, we obtain that the total number of elementary pieces does not exceed $18p^2k^{r*} + 12p^4$. Taking the sum over all simple factors of D we prove the lemma. $\square$

10.7 Appendix 7. On the Proof of Geometric Lemmas Allowing to Deal with Clusters instead of Boxes

In the proof of Lemma 5.13 it is important that we deal with the same curve generated by the determinant and just change the argument $\vec{\varkappa}$. At the same time, a priori we have the estimates for the resolvent of the operator reduced onto a particular cluster. The form of clusters can vary which formally changes the projector and thus the determinant and the

curve. Here we explain how to deal with this situation. We will show that every cluster (white, grey or black) can be embedded into a box of the fixed size (depending on the color of the cluster) such that the estimate for the resolvent on this box is essentially the same as for the embedded cluster. We also notice that the estimate for the number of points of $\mathcal{M}^{(2)}$ inside these boxes is the same as the worst possible estimate for the corresponding cluster (see Lemmas 4.12, 4.13, 4.14). This justifies the application of Lemma 5.13 in the proof of Lemma 5.14.

By construction, white clusters are separated from each other by the distance no less than $k^{\gamma_{r_1}/6}$. Grey and black clusters are separated by the distance at least $k^{\gamma_{r_1}/2+2\delta_0 r_1}$ and $k^{\gamma_{r_1}+\delta_0 r_1}$, respectively. Consider first a white cluster. Let Π_w be a singular white cluster, namely,

$$\|(P_w(H - k^{2l})P_w)^{-1}\| > k^\xi, \quad \xi > k^{\gamma_{r_1}/6-2\delta}, \quad (459)$$

here and below $H = H(\vec{x}^{(2)}(\varphi_0))$, P_w is the projector corresponding to Π_w . By construction, Π_w belongs to a small white box and its neighbors. Let us refer to it as expanded small white box. Its size is $3k^{\gamma_{r_1}/2+2\delta_0 r_1}$ and it contains less than $k^{\gamma_{r_1}/6-\delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$.

Lemma 10.10. *If (459) holds for a white cluster Π_w then*

$$\|(P(H - k^{2l})P)^{-1}\| > ck^{k^{\gamma_{r_1}/6-2\delta}}, \quad (460)$$

P being the projector corresponding to the expanded small white box Π containing Π_w . The box Π has the size $3k^{\gamma_{r_1}/2+2\delta_0 r_1}$ and contains less than $k^{\gamma_{r_1}/6-\delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$.

Proof. Assume (459) holds, but (460) does not. Let $f \in P_w \ell^2$ be such that $\|f\| = 1$, $P_w(H - k^{2l})f = o(k^{-\xi})$, $\xi = k^{\gamma_{r_1}/6-2\delta}$. Let us define

$$g := f - (P(H - k^{2l})P)^{-1}(P - P_w)Vf.$$

Now we have:

$$\begin{aligned} P(H - k^{2l})Pg &= P(H - k^{2l})f - (P - P_w)Vf = \\ P_w(H - k^{2l})f + (P - P_w)(H - k^{2l})f - (P - P_w)Vf &= \\ P_w(H - k^{2l})P_w f + (P - P_w)(H_0 - k^{2l})P_w f &= P_w(H - k^{2l})P_w f = o(k^{-\xi}). \end{aligned} \quad (461)$$

If we show that

$$\|(P(H - k^{2l})P)^{-1}(P - P_w)Vf\| = o(1), \quad (462)$$

which means $\|g\| = 1 + o(1)$, then the lemma easily follows by the way of contradiction. Thus, it remains to prove (462). Denote $\tilde{f} := (P - P_w)Vf$. Let $\tilde{H}_w^{(2)}$ be the operator

consisting of k^δ -clusters in Π . Namely, $\tilde{H}_w^{(2)} = \sum_i P_i H P_i$, P_i being projectors onto k^δ -clusters. Formally,

$$\begin{aligned} (P(H - k^{2l})P)^{-1} \tilde{f} &= \sum_{r=0}^{r_0} (\tilde{H}_w^{(2)} - k^{2l})^{-1} \left(-(H - \tilde{H}_w^{(2)})(\tilde{H}_w^{(2)} - k^{2l})^{-1} \right)^r \tilde{f} + \\ & (P(H - k^{2l})P)^{-1} \left(-(H - \tilde{H}_w^{(2)})(\tilde{H}_w^{(2)} - k^{2l})^{-1} \right)^{r_0+1} \tilde{f}, \quad r_0 = \lceil k^{\gamma r_1/6-\delta} \rceil - 1. \end{aligned} \quad (463)$$

Some of k^δ -clusters $P_i H P_i$ are resonant. However, their distance to the boundary of any white cluster is greater than $k^{\gamma r_1/6}$. Using this fact and considering as in the proof of (122), we obtain

$$\left\| \left(-(H - \tilde{H}_w^{(2)})(\tilde{H}_w^{(2)} - k^{2l})^{-1} \right)^r \tilde{f} \right\| < ck^{-\beta r} \quad \text{when } r \leq r_0 + 1,$$

since $(\tilde{f})_{\mathbf{m}} \neq 0$ only near the boundary of a white cluster. Hence, the right hand part of (463) is well defined. Now, substituting the last estimate into (463), applying the same arguments as in the proof of Theorem 4.1 and using the estimate opposite to (460) we obtain (462). $\square$

For singular grey and black clusters the proof is very similar. So, we just introduce corresponding objects and formulate the results.

Let Π_g be a singular grey cluster, i.e.

$$\|(P_g(H - k^{2l})P_g)^{-1}\| > k^\xi, \quad \xi > k^{\gamma r_1/2+2\delta_0 r_1-2\delta}, \quad (464)$$

P_g being the projector corresponding to Π_g . By construction, Π_g belongs to a big white box and its neighbors. We refer to it as expanded big white box. Its size is $3k^{\gamma r_1}$ and it contains less than $k^{\gamma r_1/2+\delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$.

Lemma 10.11. *If (464) holds for a grey cluster Π_g , such that all the white clusters imbedded into it do not satisfy (460), then*

$$\|(P(H - k^{2l})P)^{-1}\| > ck^{k^{\gamma r_1/2+2\delta_0 r_1-2\delta}}, \quad (465)$$

P being the projector corresponding to the expanded big white box Π containing Π_g . The box Π has the size $3k^{\gamma r_1}$ and it contains less than $k^{\gamma r_1/2+\delta_0 r_1}$ elements of $\mathcal{M}^{(2)}$.

The proof is analogous to that of Lemma 10.10 up to the obvious changes: instead of P_w we take P_g , $r_0 = k^{\frac{1}{2}\gamma r_1+2\delta_0 r_1-\delta}$ and $\tilde{H}_w^{(2)}$ is replaced by $\tilde{H}_g^{(2)}$ which consists of k^δ non-resonance clusters and white clusters, which do not satisfy (460).

Let Π_b be a singular black cluster, i.e.

$$\|(P_b(H - k^{2l})P_b)^{-1}\| > k^\xi, \quad \xi > k^{\gamma r_1+\delta_0 r_1-2\delta}, \quad (466)$$

P_b being the projector corresponding to Π_b . By Lemma 4.12 any black cluster can be covered by a box of the size $ck^{3\gamma r_1/2+3}$ containing less than $ck^{\gamma r_1+3}$ elements of $\mathcal{M}^{(2)}$. We refer to it as expanded black box.

Lemma 10.12. *If (466) holds for a black cluster Π_b , such that all the white and grey clusters imbedded into it do not satisfy (460), (465), then*

$$\|(P(H - k^{2l})P)^{-1}\| > ck^{k^{\gamma r_1 + \delta_0 r_1 - 2\delta}}, \quad (467)$$

P being the projector corresponding to the expanded black box containing Π_b . The box Π has the size $ck^{3\gamma r_1/2+3}$ and it contains less than $ck^{\gamma r_1+3}$ elements of $\mathcal{M}^{(2)}$.

References

- [1] E. I. Dinaburg, Ya. Sinai, *The One-dimensional Schrödinger Equation with a Quasiperiodic Potential*, Funct. Anal. Appl. **9** (1975), 279–289.
- [2] H. Rüssmann, *On the one dimensional Schrödinger equation with a quasi-periodic potential*, Ann. N. Y. Acad. Sci. **357** (1980), 90–107.
- [3] R. Johnson, J. Moser, *The rotation number for almost periodic potentials*, Commun. Math. Phys. **84** (1982), 403–438.
- [4] J. Moser, J. Püschel, *An extension of a result by Dinaburg and Sinai on quasi-periodic potentials*, Comment. Math. Helvetic **59** (1984), 39–85.
- [5] H. L. Cycon, R. G. Froese, W. Kirsch, B. Simon, *Schrödinger Operators*, Berlin: Springer Verlag, 1987, corrected and extended 2nd printing, Springer Verlag, 2008.
- [6] L. Pastur, A. Figotin, *Spectra of Random and Almost-Periodic Operators*, Springer-Verlag, 1992, 583 pp.
- [7] L. H. Eliasson, *Floquet Solutions for the One-dimensional Quasi-periodic Schrödinger Equation*, Comm. Math. Phys. **146** (1992), no. 3, 447–482.
- [8] S. Jitomirskaya, *Metal-Insulator Transition for the Almost Mathieu Operator*, Ann. of Math. **150** (1999), 1159–1175.
- [9] D. Damanik, R. Killip, and D. Lenz, *Uniform spectral properties of one-dimensional quasicrystals. III. Alpha-continuity*, Commun. Math. Phys. **212** (2000), 191–204.
- [10] B. Simon, *Schrödinger Operators in the Twentieth Century*, J. Math. Phys. **41** (2000), no. 6, 3523–3355.
- [11] A. Fedotov, F. Klopp, *On the Singular Spectrum for Adiabatic Quasi-periodic Schrödinger Operators on the real line*, Ann. Henri Poincaré **5** (2004), no. 5, 929–978.

- [12] A. Fedotov, F. Klopp, *On the Absolutely Continuous Spectrum of One-dimensional Quasi-periodic Schrödinger operators in the Adiabatic Limit*, Trans. Amer. Math. Soc. **357** (2005), no. 11, 4481–4516.
- [13] A. Fedotov, F. Klopp, *Strong Resonant Tunneling, Level Repulsion and Spectral Type for One-dimensional Adiabatic Quasi-periodic Schrödinger Operators*, Ann. Sci. École Norm. Sup. (4) **38** (2005), no. 6, 889–950.
- [14] M. A. Shubin, *Density of States for Selfadjoint Elliptic Operators with Almost Periodic Coefficients*, Trudy sem. Petrovskii (Moscow University) **3** (1978), 243–281.
- [15] J. Bellissard, R. Lima, and E. Scoppola, *Localization in n -dimensional incommensurable structures*, Commun. Math. Phys. **88**, (1983) 465–477.
- [16] A. L. Figotin, L. A. Pastur, *An Exactly Solvable Model of a Multidimensional Incommensurate Structure*, Commun. Math. Physics **95** (1984), 401–425.
- [17] J. Bellissard, *Almost periodicity in solid state Physics and C^* -algebras*, Mat.-Fys. Medd danske Vid. Selsk. **42** (1989), no. 3, 35–75.
- [18] V. Chulaevsky, E. I. Dinaburg, *Methods of KAM theory for Long-Range Quasiperiodic Potentials on $\mathbb{Z}^{\vec{\nu}}$. Pure Point Spectrum*, Commun. Math. Physics **153** (1993), no. 3, 559–577.
- [19] J. Bourgain, M. Goldstein, *On Nonperturbative Localization with Quasi-Periodic Potential*, Ann. of Math. **152** (2000), no. 3, 835–879.
- [20] J. Bourgain, M. Goldstein, W. Schlag, *Anderson Localization on $\mathbb{Z}^2$ with Quasi-Periodic Potential*, Acta Math. **188** (2002), 41–87.
- [21] J. Bourgain, *On Quasi-Periodic Lattice Schrödinger Operators*, Discrete and Continuous Dynamical Systems **10** (2004), no. 1&2, 75–88.
- [22] D. Damanik, *A Version of Gordon’s Theorem for Multidimensional Schrödinger Operators*, Trans. Amer. Math. Soc. **356** (2004), 495–507.
- [23] L. Parnowski, R. Shterenberg, *Complete Asymptotic Expansion of the Integrated Density of States of Multidimensional Almost-periodic Schrödinger Operators*, Ann. of Math. **176** (2012), no. 2, 1039–1096.
- [24] S. Morozov, L. Parnowski, R. Shterenberg, *Complete Asymptotic Expansion of the Integrated Density of States of Multidimensional Almost-periodic Pseudo-differential Operators*, preprint arXiv: 1204.1076.

- [25] V. N. Popov, M. M. Skriganov, *Remark on the Structure of the Spectrum of a Two-Dimensional Schrödinger Operator with Periodic Potential*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **109** (1981), 131–133; English transl.: J. Soviet Math. **24** (1984), no. 2, 239–240.
- [26] M. M. Skriganov, A. V. Sobolev, *On the Spectrum of a Limit-Periodic Schrödinger Operator*, Algebra i Analiz, **17** (2005), no. 5; Engl. Transl.: St. Petersburg Math. J. **17** (2006), 815–833.
- [27] Yu. Karpeshina, Y.-R. Lee, *Spectral properties of polyharmonic operators with limit-periodic potential in dimension two*, D’Analyse Mathématique **102** (2007), 225–310.
- [28] Yu. Karpeshina, Y.-R. Lee, *Absolutely Continuous Spectrum of a Polyharmonic Operator with a Limit Periodic Potential in Dimension Two*, Communications in Partial Differential Equations **33** (2008), no. 9, 1711–1728.
- [29] Yu. Karpeshina, Y.-R. Lee, *Spectral properties of the Schrödinger operator with limit-periodic potential in dimension two*, (arXiv:1008.4632v1 [math-ph], pp 1-89), submitted for publication.
- [30] G. Gallavotti, *Perturbation Theory for Classical Hamiltonian Systems*, in *Scaling and Self-Similarity in Physics* edited by J. Froehlich, Birkhäuser, Basel, Switzerland, 1983, 359–424.
- [31] L.E. Thomas, S.R. Wassel, *Stability of Hamiltonian systems at high energy*, J. Math. Phys. **33**(10), (1992), 3367–3373.
- [32] L. E. Thomas and S. R. Wassel, *Semiclassical Operators at High Energy*, Lecture Notes in Physics **403**, edited by E. Balslev, Springer-Verlag, 1992, 194–223.
- [33] J. Frölich, T. Spencer, *Absence of Diffusion in the Anderson Tight Binding Model for Large Disorder and Low Energy*, Commun. Math. Physics **88** (1983), no. 2, 151–184.
- [34] J. Bourgain, *Quasiperiodic Solutions of Hamilton Perturbations of 2D Linear Schrödinger Equation*, Ann. of Math. **148** (1998), no. 2, 363–439.
- [35] I. M. Gel’fand, *Expansion in Eigenfunctions of an Equation with Periodic Coefficients*, Dokl. Akad. Nauk SSSR **73** (1950), 1117–1120 (in Russian).
- [36] M. Reed, B. Simon, *Methods of Modern Mathematical Physics*, Vol IV, Academic Press, 3rd ed., New York-San Francisco-London (1987), 396 pp.